

## A Multi-Wavelength Simultaneous Study of the Composition of the Halley Family Comet 8P/Tuttle

E. Jehin · D. Bockelée-Morvan · N. Dello Russo · J. Manfroid ·  
D. Hutsemékers · H. Kawakita · H. Kobayashi · R. Schulz ·  
A. Smette · J. Stüwe · M. Weiler · C. Arpigny · N. Biver ·  
A. Cochran · J. Crovisier · P. Magain · H. Rauer · H. Sana ·  
R. J. Vervack · H. Weaver · J.-M. Zucconi

Received: 26 January 2009 / Accepted: 8 May 2009 / Published online: 30 May 2009  
© Springer Science+Business Media B.V. 2009

**Abstract** We report on simultaneous optical and infrared observations of the Halley Family comet 8P/Tuttle performed with the ESO Very Large Telescope. Such multi-wavelength and coordinated observations are a good example of what can be done to support space missions. From high resolution optical spectroscopy of the CN (0,0) 388 nm and NH<sub>2</sub> (0,9,0) 610 nm bands using UVES at UT2 we determined  $^{12}\text{C}/^{13}\text{C} = 90 \pm 10$  and  $^{14}\text{N}/^{15}\text{N} = 150 \pm 20$  in CN and we derived a nuclear spin temperature of NH<sub>3</sub> of  $29 \pm 1$  K. These values are similar to those found in Oort-Cloud and Jupiter Family comets. From low resolution long slit spectroscopy with FORS1 at UT2 we determined the CN, C<sub>3</sub> and C<sub>2</sub> production rates and the parent and daughter scale lengths up to  $5.2 \cdot 10^5$  km

---

Based on observations collected at the European Southern Observatory, Paranal, Chile (ESO Prog. 080.C-0615 and 280.C-5053).

---

E. Jehin (✉) · J. Manfroid · D. Hutsemékers · C. Arpigny · P. Magain  
Institut d'Astrophysique et de Géophysique de l'Université de Liège, Allée du 6 août 17, 4000 Liège,  
Belgium  
e-mail: ejehin@ulg.ac.be

D. Bockelée-Morvan · M. Weiler · N. Biver · J. Crovisier  
LESIA, Observatoire de Paris, 92195 Meudon, France

N. Dello Russo · R. J. Vervack · H. Weaver  
The Johns Hopkins University Applied Physics Laboratory, Laurel, MD 2072, USA

A. Smette · H. Sana  
ESO, Vitacura Casilla, 19001 Santiago, Chile

H. Kawakita · H. Kobayashi  
Kyoto Sangyo University, Motoyama, Kamigamo, Kita-ku, Kyoto 603-8555, Japan

A. Cochran  
Department of Astronomy and McDonald Observatory, University of Texas, Austin 1400, USA

R. Schulz  
ESA/RSSD, ESTEC, 2200 AG Noordwijk, The Netherlands

tailward. From high resolution IR spectroscopy with CRIRES at UT1 we measured simultaneously the production rates and mixing ratios of H<sub>2</sub>O, HCN, C<sub>2</sub>H<sub>2</sub>, CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, and CH<sub>3</sub>OH.

**Keywords** Comets · 8P/Tuttle · Spectroscopy · Isotope ratios · NH<sub>2</sub> OPR · Production rates · Mixing ratios

## 1 Introduction

Determining the composition of cometary nuclei is essential for understanding the formation and evolution of volatile material within our solar system. Observational evidence supports chemical diversity among comets that may reflect the diversity of conditions in comet-forming regions in the solar nebula.

However, comets experienced different processing histories whose importance has to be properly investigated both theoretically and observationally. The orbits of Halley-family comet (HFCs) have evolved to periods less than 200 years, making them the most processed Oort-cloud comets. However, HFCs are underrepresented in compositional surveys.

The 2008 apparition of comet 8P/Tuttle was an excellent opportunity to perform a detailed investigation of the composition of a HFC with modern techniques.

Observations during the 1980 apparition showed 8P/Tuttle to be typical in C<sub>2</sub> abundance relative to H<sub>2</sub>O (C<sub>2</sub>/H<sub>2</sub>O  $\sim 4.1 \times 10^{-3}$ ), with an inferred H<sub>2</sub>O production rate between  $\sim 1.6$  and  $5.7 \times 10^{28}$  mol s<sup>-1</sup> at a heliocentric distance (R<sub>h</sub>) between 1.02 and 1.16 AU (A'Hearn et al. 1995).

## 2 Observing Circumstances

8P/Tuttle made a close approach to Earth at 0.25 AU on Jan 2, 2008. On Jan 16, 28 and Feb 4 UT, we undertook simultaneous spectroscopic observations in the near-infrared and the visible range using the CRIRES, FORS1 and UVES instruments installed at the UT1 and UT2 units of the Very Large Telescope (VLT) of the European Southern Observatory (ESO). The observing circumstances are given in Table 1. The weather was cloudy on the visitor mode night of Jan 16 and only UVES spectra could be used. On the two other nights (service mode) simultaneous observations with the three instruments were performed, the weather was clear and the seeing was 0.5–1.5".

---

J. Stüwe  
Leiden Observatory, 2300 RA Leiden, The Netherlands

J.-M. Zucconi  
Observatoire de Besançon, 25010 Besançon, France

H. Rauer  
Institute of Planetary Research, DLR, 12489 Berlin, Germany

**Table 1** Observing circumstances

Date	$R_h$ (AU)	$V_{Rh}$ (km/s)	$\Delta$ (AU)	$V_\Delta$ (km/s)	P.A. ( $^\circ$ )	$\alpha$ ( $^\circ$ )
16/01/08	1.04	-4.32	0.36	21.75	67.5	70.9
28/01/08	1.03	0.38	0.52	24.79	74.7	70.5
04/02/08	1.03	3.13	0.62	24.22	80.1	67.9


$R_h$  and  $V_{Rh}$  are the heliocentric distance and velocity,  $\Delta$  and  $V_\Delta$  are the geocentric distance and velocity, respectively. P.A. stands for position angle of the antisolar direction,  $\alpha$  is the phase angle

### 3 C and N Isotope Ratios, $NH_2$ OPR


The high resolution spectroscopic observations in the optical domain ( $\lambda/\Delta\lambda = 80000$ , [303–1004 nm]) had several objectives, among which measurements of carbon and nitrogen isotopic ratios and the ortho-to-para (OPR) ratio in  $NH_2$ .

Synthetic spectra of  $^{12}C^{14}N$ ,  $^{13}C^{14}N$ ,  $^{12}C^{15}N$  (0,0) bands at 388 nm were computed using a fluorescence model (Zucconi and Festou 1985). The isotope mixture was adjusted to best fit the observed continuum subtracted spectrum (Arpigny et al. 2003). This is illustrated in Fig. 1. The carbon isotopic ratio found in 8P/Tuttle ( $^{12}C/^{13}C = 90 \pm 20$ ) is in agreement with other comets and with the terrestrial and solar value (89). The  $^{15}N$  enrichment ( $^{14}N/^{15}N = 150 \pm 25$ ) with respect to the terrestrial atmospheric value (272) is consistent with the value observed in about 20 comets of the Oort-Cloud and Jupiter-Family types (Manfroid et al. 2009, in preparation).

The red part of the UVES spectra gives access to the  $NH_2$  (0,9,0) band at 610 nm, which was used to measure the OPR of  $NH_2$  on the basis of a fluorescence excitation model (Kawakita et al. 2001) (Fig. 2). A mean OPR of  $3.29 \pm 0.07$  was derived, providing a nuclear spin temperature of  $NH_3$  of  $T_{spin} = 29 \pm 2$  K in agreement with measurements in other comets (27–35 K). If  $T_{spin}$  is correlated with the location of formation of the comet in



**Fig. 1** A small part of the CN (0,0) mean spectrum (black line). The positions of the  $^{13}C^{14}N$  (red ticks) and  $^{12}C^{15}N$  (blue ticks) isotopic lines are indicated. The synthetic spectrum with  $^{12}C/^{13}C = 90$  and  $^{14}N/^{15}N = 150$  is superimposed (blue line). The red dotted line is the synthetic spectrum without  $^{13}C^{14}N$  and  $^{12}C^{15}N$  (Color figure online)



**Fig. 2** Observed and modelled spectrum of the  $\text{NH}_2$  (0,9,0) band at 610 nm on Feb 04. Strong ortho- and para- $\text{NH}_2$  lines are marked

the solar nebula, the Halley type comet 8P/Tuttle formed in a similar environment as most of the Oort-Cloud and Jupiter-family comets.


#### 4 CN, $\text{C}_3$ and $\text{C}_2$ Density Profiles, Production Rates and Scale Lengths

Long slit spectroscopy ( $\lambda/\Delta\lambda = 440$ , [330–660 nm]) was performed to measure the radial distributions of the CN,  $\text{C}_2$ ,  $\text{C}_3$  radicals and their parent production rates, in order to constrain their origin. The slit was put along the tail and offsets were applied in order to cover continuously the coma up to  $5.2 \cdot 10^5$  km in the tail direction. Spatial profiles of the gaseous emissions of CN [3830–3905 Å],  $\text{C}_3$  [3975–4150 Å] and  $\text{C}_2$  [4860–5210 Å] were extracted by integrating the flux densities over the above cited wavelength ranges (Fig. 3). The fluxes were then converted to column densities.

The Haser model was used to fit the CN,  $\text{C}_3$  and  $\text{C}_2$  column density profiles and to derive production rates, parent ( $l_p$ ) and daughter ( $l_d$ ) scale lengths (Table 2). For CN, the spatial coverage was not sufficient on the sunward side, so that the scale lengths were fixed to the values measured tailward. A parent and daughter expansion velocity of 1.0 km/s was assumed.

$\text{C}_2/\text{CN}$  and  $\text{C}_3/\text{CN}$  production rate ratios are consistent with values measured by narrowband photometry (Schleicher IAUC8903, pers.com.) and fall near the mean values for typical comets according to A'Hearn et al. (1995) taxonomy (as already observed at previous 8P apparitions).

For CN the sunward side is not well fitted with the ( $l_p$ ,  $l_d$ ) parameters measured tailward. The CN parent scale length is not fully in agreement with that one expected from the photodissociation of HCN. Using the Haser-equivalent model, the photodissociation rates of  $1.55 \cdot 10^{-5} \text{ s}^{-1}$  (HCN),  $3.2 \cdot 10^{-6} \text{ s}^{-1}$  (CN) (Bockelée-Morvan and Crovisier 1985, Huebner et al. 1992) and a CN ejection velocity of 0.86 km/s (Fray et al. 2005) one gets  $l_p = 4.5 \cdot 10^4$  km and  $l_d = 4.0 \cdot 10^5$  km to be compared with the measured values of  $3.5 \cdot 10^4$  km and  $1.9 \cdot 10^5$  km, respectively. The comparison of the CN and HCN



**Fig 3** Spatial profile of the gaseous emission of CN [3830–3905 Å], C<sub>3</sub> [3975–4150 Å] and C<sub>2</sub> [4860–5810 Å]

**Table 2** CN, C<sub>3</sub>, C<sub>2</sub> production rates and their parent (lp) and daughter (ld) scale lengths for the sunward and tailward sides using a Haser model

Radical	Date	Production rate (mol/s)	lp (km)		ld (km)	
			Tail/Sun	Tail/Sun	Tail/Sun	Tail/Sun
CN	28/01	$1.4/1.0 \times 10^{26}$	$3.6 \times 10^4$	$1.6 \times 10^5$		
	04/02	$1.3/1.1 \times 10^{26}$	$3.4 \times 10^4$	$2.1 \times 10^5$		
C <sub>3</sub>	28/01	$8.1/4.8 \times 10^{24}$	$5.6/3.9 \times 10^3$	$3.5/2.7 \times 10^4$		
	04/02	$10.2/4.8 \times 10^{24}$	$8.0/3.4 \times 10^3$	$3.3/4.7 \times 10^4$		
C <sub>2</sub>	28/01	$1.9/1.2 \times 10^{26}$	$6.5/5.1 \times 10^4$	$6.7/5.1 \times 10^4$		
	04/02	$2.0/1.4 \times 10^{26}$	$5.8/4.9 \times 10^4$	$5.8/4.8 \times 10^4$		


production rates presented in the next section suggests that HCN is not the major source of CN in this comet. A possible candidate is C<sub>2</sub>N<sub>2</sub> (Bockelée-Morvan and Crovisier 1985; Festou et al. 1998; Fray et al. 2005). However, using the C<sub>2</sub>N<sub>2</sub> photodissociation rate and CN ejection velocity of 2–3 km s<sup>-1</sup> given by Bockelée-Morvan and Crovisier (1985), we infer an equivalent parent scale length lp = 0.6–1.2 × 10<sup>4</sup> km much shorter than observed.

The fits for C<sub>2</sub> are the poorest. A complex chemistry model is needed to reproduce the C<sub>2</sub> and C<sub>3</sub> profiles as these species are produced in more than one reaction step from several parent species (Helbert et al. 2005).

### 5 Mixing Ratios and Production Rates of Parent Molecules

Infrared high resolution spectroscopy observations ( $\lambda/\Delta\lambda = 50000$ , [2.9–3.3 μm]) with CRIRES were conducted to measure the production rates and abundance of several parent molecules (Fig. 4). An important objective was to compare HCN and C<sub>2</sub>H<sub>2</sub> production rates to those of their daughter products CN and C<sub>2</sub> observed simultaneously with FORS 1 (Table 3).

The mixing HCN/H<sub>2</sub>O ratio is lower, by a factor of 3, than the mean value of 0.26% (Disanti and Mumma 2008) measured in Oort-Cloud comets from IR observations. A similar conclusion is reached from millimetre observations (Biver et al. 2008) and from the



**Fig. 4** Spectrum of H<sub>2</sub>O and OH observed on January 28 with CRIRES

**Table 3** Mixing ratios and production rates of parent molecules (Kobayashi et al. 2009, in preparation)

Molecules	Trot (K)	Production rate (mol/s)	Mixing ratio (%)
<i>Jan 28</i>			
H <sub>2</sub> O	70 ± 15	(4.6 ± 0.4) × 10 <sup>28</sup>	100
HCN	54 ± 9	(3.4 ± 1.1) × 10 <sup>25</sup>	0.07 ± 0.02
C <sub>2</sub> H <sub>2</sub>	(70)	(2.1 ± 0.7) × 10 <sup>25</sup>	0.05 ± 0.02
<i>Feb 4</i>			
H <sub>2</sub> O	65 +15/-13	(3.0 ± 0.2) × 10 <sup>28</sup>	100
CH <sub>4</sub>	(70)	(1.7 ± 1.1) × 10 <sup>26</sup>	0.6 ± 0.4
C <sub>2</sub> H <sub>6</sub>	(70)	(6.7 ± 1.3) × 10 <sup>25</sup>	0.23 ± 0.04
CH <sub>3</sub> OH	(70)	(9.7 ± 1.1) × 10 <sup>27</sup>	3.3 ± 0.3

NIR observations of Bonev et al. (2008) and Boehnhardt et al. (2008). Strong depletions of C<sub>2</sub>H<sub>2</sub> and C<sub>2</sub>H<sub>6</sub> are also observed, compared to the mean value in Oort-Cloud comets but the CH<sub>3</sub>OH and CH<sub>4</sub> mixing ratios are typical.

The CN/HCN ratio is ~3, suggesting that there is another source of CN in the coma. This has been already noticed for a number of comets (Fray et al. 2005).

A similar conclusion is reached for C<sub>2</sub>, as the C<sub>2</sub>/C<sub>2</sub>H<sub>2</sub> ratio is ~8.

## References

- M. A'Hearn et al., *Icarus* **118**, 223 (1995). doi:[10.1006/icar.1995.1190](https://doi.org/10.1006/icar.1995.1190)  
 C. Arpigny et al., *Science* **301**, 1522 (2003). doi:[10.1126/science.1086711](https://doi.org/10.1126/science.1086711)  
 N. Biver et al., in *Asteroids, Comets, Meteors*, LPI Contribution No. 1405, Baltimore, MD, 14–18 July 2008  
 D. Bockelée-Morvan, J. Crovisier, *Astron. Astrophys.* **151**, 90 (1985)  
 H. Boehnhardt et al., *Astrophys. J.* **683**, L71 (2008)  
 B.P. Bonev et al., *Astrophys. J.* **680**, L61 (2008)  
 M.A. DiSanti, M.J. Mumma, *Space Sci. Rev.* **138**, 127 (2008)

- M. Festou et al., *BAAS* **30**, 1089 (1998)  
N. Fray et al., *P&SS* **53**, 1243 (2005)  
J. Helbert et al., *Astron. Astrophys.* **442**, 1107 (2005)  
W.F. Huebner et al., *Ap&SS* **195**, 1 (1992)  
H. Kawakita et al., *Science* **294**, 1089 (2001)  
J.-M. Zucconi, M. Festou, *Astron. Astrophys.* **150**, 180 (1985)