

Erratum to: Antifungal Susceptibility Profile of Cryptic Species of *Aspergillus*

Ana Alastruey-Izquierdo · Laura Alcazar-Fuoli · Manuel Cuenca-Estrella

Published online: 9 December 2014
© Springer Science+Business Media Dordrecht 2014

**Erratum to: Mycopathologia (2014) 178:427–433
DOI 10.1007/s11046-014-9775-z**

Unfortunately, there are some errors in Table 2:

TRB values are MIC and not MEC (the line for MEC in the headings should be only for echinocandins: CPF, MCF and ANF).

GMs for *A. fumigatiafinnis* for ICZ: 5.04 (Instead of 2.17), VCZ: 3.17 (Instead of 1.83), PCZ: 0.44

(Instead of 0.27), TRB: 0.89 (Instead of 0.65), CPF: 0.22 (Instead of 0.27).

PCZ range for *A. hiratsukae* is 0.03–16.00 (Instead of 0.03–6.00).

CPF range for *A. udagawae* is 0.03–2.00 (Instead of 0.25–2.00).

The corrected table is given here.

The online version of the original article can be found under doi:[10.1007/s11046-014-9775-z](https://doi.org/10.1007/s11046-014-9775-z).

A. Alastruey-Izquierdo (✉)
Mycology Reference Laboratory, National Centre for Microbiology, Spanish Network for the Research in Infectious Diseases (REIPI RD12/0015), Instituto de Salud Carlos III, Majadahonda, Madrid, Spain
e-mail: anaastruey@isciii.es

L. Alcazar-Fuoli · M. Cuenca-Estrella
Mycology Reference Laboratory, National Centre for Microbiology, Instituto de Salud Carlos III, Majadahonda, Madrid, Spain

Table 2 Susceptibility profile of cryptic *Aspergillus* species isolated in the Mycology Reference Laboratory of Spain

Species (no.)	MIC (mg/L)						MEC (mg/L)			
	AMB	ICZ	VCZ	PCZ	TRB	CPF	MCF	ANF		
<i>A. alliacetus</i> (30)	GM	19.25	0.20	0.52	0.12	0.44	12.15	3.78	1.98	
	Range	0.5–32.00	0.03–16.00	0.12–16.00	0.015–16.00	0.03–32.00	0.12–32.00	0.03–32.00	0.03–32.00	
	MIC ₅₀	32.00	0.12	0.50	0.06	0.25	32.00	32.00	32.00	
	MIC ₉₀	32.00	2.00	2.00	0.50	32.00	32.00	32.00	32.00	
<i>A. calidoustus</i> (19)	GM	0.96	8.61	6.20	6.86	0.72	0.52	0.04	0.04	
	Range	0.12–32.00	1.00–16.00	4.00–16.00	1.00–16.00	0.032–32.00	0.03–32.00	0.03–1.00	0.03–0.50	
	MIC ₅₀	1.00	16.00	8.00	16.00	0.50	1.00	0.03	0.03	
	MIC ₉₀	16.00	16.00	8.00	16.00	4.00	4.00	0.06	0.25	
<i>A. fumigatiifinis</i> (6)	GM	4.76	5.04	3.17	0.44	0.89	0.22	0.04	0.05	
	Range	2.00–16.00	0.25–16.00	2.00–8.00	0.12–4.00	0.25–4.00	0.03–0.50	0.03–0.06	0.03–0.25	
	MIC ₅₀	16.00	16.00	4.00	0.50	1.00	0.50	0.03	0.03	
	MIC ₉₀	16.00	16.00	8.00	4.00	3.00	0.50	0.06	0.25	
<i>A. hirsutiae</i> (5)	GM	1.74	0.87	1.15	0.16	2.30	0.29	0.06	0.12	
	Range	0.50–32.00	0.12–16.00	0.25–16.00	0.03–16.00	0.12–32.00	0.03–4.00	0.03–0.50	0.03–2.00	
	MIC ₅₀	1.00	0.50	1.00	0.06	1.00	0.50	0.03	0.03	
	MIC ₉₀	32.00	16.00	16.00	16.00	32.00	4.00	0.50	2.00	
<i>A. lentulus</i> (26)	GM	2.98	1.45	3.41	0.22	1.05	1.74	0.10	0.09	
	Range	0.25–32.00	0.12–16.00	0.25–16.00	0.03–1.00	0.12–32.00	0.03–32.00	0.03–32.00	0.03–32.00	
	MIC ₅₀	4.00	0.50	4.00	0.25	0.75	2.00	0.03	0.03	
	MIC ₉₀	16.00	16.00	8.00	0.50	16.00	32.00	32.00	32.00	
<i>A. tubingensis</i> (29)	GM	0.10	0.63	0.91	0.12	0.31	0.28	0.04	0.03	
	Range	0.06–2.00	0.12–32.00	0.25–2.00	0.03–0.50	0.03–2.00	0.06–1.00	0.03–0.25	0.03–0.06	
	MIC ₅₀	0.12	0.50	1.00	0.12	0.50	0.50	0.03	0.03	
	MIC ₉₀	0.12	2.00	2.00	0.25	1.00	1.00	0.12	0.03	
<i>A. udagawae</i> (5)	GM	2.00	0.57	2.30	0.22	0.66	0.33	0.04	0.03	
	Range	0.50–4.00	0.25–1.00	2.00–4.00	0.12–0.25	0.25–2.00	0.03–2.00	0.03–0.12	0.03–0.06	
	MIC ₅₀	2.00	0.50	2.00	0.25	0.50	0.50	0.03	0.03	
	MIC ₉₀	4.00	1.00	4.00	0.25	2.00	2.00	0.12	0.05	

GM geometric mean, MEC minimal effective concentration, MIC₅₀ MIC causing inhibition of 50 % of the isolates, MIC₉₀ MIC causing inhibition of 90 % of the isolates, AMB amphotericin B, ICZ itraconazole, VCZ voriconazole, PCZ posaconazole, TRB terbinafine, CPF caspofungin, MCF micafungin, ANF anidulafungin