Correction to: Influence of heterozygosity and competition on morphological tree characteristics of *Quercus rubra* L.: a new single-tree based approach

Katharina Burkardt¹ · Tim Pettenkofer² · Christian Ammer¹ · Oliver Gailing² · Ludger Leinemann² · Dominik Seidel¹ · Torsten Vor¹

Published online: 5 August 2021
© The Author(s) 2021

Correction to: New Forests (2021) 52:679–695

<https://doi.org/10.1007/s11056-020-09814-1>

The article "Influence of heterozygosity and competition on morphological tree characteristics of *Quercus rubra* L.: a new single-tree based approach" written by Katharina Burkardt · Tim Pettenkofer · Christian Ammer · Oliver Gailing · Ludger Leinemann · Dominik Seidel · Torsten Vor was originally published electronically on the publisher's internet portal on 3rd November 2020 without open access. The article is funded under Open Access funding enabled and organized by Projekt DEAL. Hence the article is forthwith distributed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0>.

The funding note has been added in the original publication.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long

---

The original article can be found online at <https://doi.org/10.1007/s11056-020-09814-1>.

✉ Katharina Burkardt
katharina.burkardt@uni-goettingen.de

¹ Department of Silviculture and Forest Ecology of the Temperate Zones, Faculty of Forest Sciences and Forest Ecology, University of Göttingen, Büsgenweg 1, 37077 Göttingen, Germany

² Department of Forest Genetics and Forest Tree Breeding, Faculty of Forest Sciences and Forest Ecology, University of Göttingen, Büsgenweg 2, 37077 Göttingen, Germany

as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.