Correction to: Capsaicin Exerts Anti-convulsant and Neuroprotective Effects in Pentylenetetrazole-Induced Seizures

Omar M. E. Abdel-Salam¹ · Amany A. Sleem² · Marawan Abd El Baset Mohamed Sayed² · Eman R. Youness³ · Nermeen Shaffie⁴

Published online: 25 February 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

Correction to: Neurochemical Research

<https://doi.org/10.1007/s11064-020-02979-3>

The original version of this article unfortunately contains an error in the Y axis units in Fig. 1b, c (the symbol μ is not clear: $\mu\text{mol/g.tissue}$).

This has been corrected by publishing this erratum.
The correct version of the Fig. 1 is given below. The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s11064-020-02979-3>.

✉ Omar M. E. Abdel-Salam
omasalam@hotmail.com

¹ Department of Toxicology and Narcotics, National Research Centre, Tahrir Street, Dokki, Cairo, Egypt

² Department of Pharmacology, Tahrir Street, Dokki, Cairo, Egypt

³ Department of Medical Biochemistry, Tahrir Street, Dokki, Cairo, Egypt

⁴ Department of Pathology, National Research Centre, Tahrir Street, Dokki, Cairo, Egypt

Fig. 1 a–d Effect of capsaicin, phenytoin or capsaicin/phenytoin on **a** brain malondialdehyde (MDA), **b** nitric oxide, **c** reduced glutathione (GSH) and **d** paraoxonase-1 (PON-1) activity in pentylenetetrazole (PTZ)-kindled rats. * $p < 0.05$ vs. vehicle-treated group and between different groups as shown in the figure. $^+p < 0.05$ vs. PTZ control group. $^{\#}p < 0.05$ vs. PTZ + capsaicin 1 mg/kg. $^{\textcircled{a}}p < 0.05$ vs. PTZ + capsaicin 1 or 2 mg/kg