Correction to: Intrathecal Epigallocatechin Gallate Treatment Improves Functional Recovery After Spinal Cord Injury by Upregulating the Expression of BDNF and GDNF

Wei Tian¹ · Xiao-Guang Han¹ · Ya-Jun Liu¹ · Guo-Qing Tang² · Bo Liu¹ · Yong-Qing Wang¹ · Bin Xiao¹ · Yun-Feng Xu¹

Published online: 22 April 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

Correction to: Neurochemical Research (2013) 38:772–779
<https://doi.org/10.1007/s11064-013-0976-5>

Since the publication of our article [1] it has come to our attention that there was an error in Figure 4 in which the bottom left immunohistochemistry panel Control/Bax was a duplication of the bottom right immunohistochemistry panel EGCG/GDNF in Figure 3.

A corrected Fig. 4 is given below.

We apologize to readers for this error.

The original article can be found online at <https://doi.org/10.1007/s11064-013-0976-5>.

✉ Wei Tian
jstspine@126.com

¹ Department of Spine Surgery, Beijing Jishuitan Hospital, No. 31, Xijiekou East Street, Xicheng District, Beijing, People's Republic of China

² Department of Orthopedics, Kunshan Chinese Traditional Medical Hospital, Kunshan, People's Republic of China

Fig. 4 Effect of EGCG treatment on changes in the expression of Bcl-2 and Bax in animals with SCI. **a** The immunohistochemistry outcome for Bcl-2 and Bax in the different groups. The positive cells were brown due to the DAB reaction. **b** The expression of Bcl-2 and Bax protein after SCI were detected by Western blot analysis using samples from each group. **c** The intensity of each band was estimated by densitometric analysis. b-actin was used as a loading control. * $p < 0.05$, ** $p < 0.01$ compared with the control group, ## $p < 0.01$, represents the comparison between the 10 and 20 mg/kg EGCG groups. All experiments were repeated 3 times and showed similar results