Correction to: Detectable Neuropsychological Differences in Early Preclinical Alzheimer's Disease: a Meta-Analysis

S. Duke Han^{1,2,3,4} · Caroline P. Nguyen¹ · Nikki H. Stricker⁵ · Daniel A. Nation³

Published online: 23 December 2017
© Springer Science+Business Media, LLC, part of Springer Nature 2017

Correction to: Neuropsychol Rev

<https://doi.org/10.1007/s11065-017-9345-5>

Errors were discovered in the reporting of processing speed data that do not impact the interpretation of findings.

Corrections to the text on page 8 should read as follows:

1. Under the Amyloid Positive Versus Negative section, following presentation of visuospatial ability results: “processing speed (12 studies: $H_g = -0.247$, $SE_g = 0.080$, 95% CI = -0.404 to -0.089 , $p = 0.002$)”
2. Under the Amyloid Positive Versus Negative section, following presentation of memory heterogeneity results: “processing speed ($Q = 13.657$, $p = 0.253$, $I^2 = 19.454$, $\tau^2 = 0.121$, and $\tau^2 = 0.015$)”

Also, below is a corrected Figure 6:

The online version of the original article can be found at <https://doi.org/10.1007/s11065-017-9345-5>

✉ S. Duke Han
Duke.Han@med.usc.edu

¹ Department of Family Medicine, USC Keck School of Medicine, 1000 S. Fremont Avenue, Unit 22, HSA Building A-6, 4th Floor, Room 6437A, Alhambra, CA 91803, USA

² Department of Neurology, USC Keck School of Medicine, Los Angeles, CA, USA

³ Department of Psychology, USC Dornsife College, Los Angeles, CA, USA

⁴ USC School of Gerontology, Los Angeles, CA, USA

⁵ Department of Psychiatry and Psychology, Mayo Clinic, Rochester, MN, USA

Fig. 6 Forest plot for processing speed for amyloid positive and negative. Difference in means reflects amyloid positive minus amyloid negative. Rhombus midpoint is the standard difference in means and the left and right points span the lower and upper limit