CORRECTION

Correction to: Comparative Study of the Dose-Dependence of OATPIB Inhibition by Rifampicin Using Probe Drugs and Endogenous Substrates in Healthy Volunteers

Issey Takehara^{1,2} • Takashi Yoshikado^{3,4} • Keiko Ishigame⁴ • Daiki Mori² • Ken-ichi Furihata⁵ • Nobuaki Watanabe⁶ • Osamu Ando⁶ • Kazuya Maeda² • Hiroyuki Kusuvara⁴ • Yuichi Sugiyama²

Published online: 21 February 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to: Pharm Res (2018) 35: 138

<https://doi.org/10.1007/s11095-018-2416-3>

There was a miscalculation of coproporphyrin I AUC_{0-24h} in the published article (Volume 35, Number 7). After the correction of AUC_{0-24h} , AUC ratio and R-square were recalculated. Then, following corrections were made in the abstract, the body, Fig. 3, Fig. 4 and Table 2 in this article. The interpretation of data in this article was not influenced.

The corrected sentences are below.

Page 1.

Abstract

line 20: coproporphyrin I ($r^2 = 0.78$).

line 23: ($r^2 = 0.48-0.70$).

Page 5.

Effect of Rifampicin on the Plasma Concentrations of Coproporphyrin I.

line 4: 2.3- and 3.5-times higher.

Correlations between the AUC_{0-24h} of Endogenous OATPIB Substrates, and of these with the AUC_{0-24h} of Atorvastatin.

line 6: coproporphyrin I ($r^2 = 0.78$).

line 8: atorvastatin (r^2 of 0.65, 0.70 and 0.48, respectively).

Issey Takehara and Takashi Yoshikado contributed equally to this work.

The online version of the original article can be found at <https://doi.org/10.1007/s11095-018-2416-3>

✉ Hiroyuki Kusuvara
kusuvara@mol.f.u-tokyo.ac.jp

¹ Biomarker Department, Daiichi Sankyo Co. Ltd., Tokyo, Japan

² Laboratory of Molecular Pharmacokinetics, Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

³ Present address: Laboratory of Clinical Pharmacology, Yokohama University of Pharmacy, 601 Matano-cho, Totsuka-ku, Yokohama-shi, Kanagawa 245-0066, Japan

⁴ Sugiyama Laboratory, RIKEN Innovation Center, RIKEN, Yokohama, Japan

⁵ P-One Clinic, Keikokai Medical Corp, Tokyo, Japan

⁶ Drug Metabolism & Pharmacokinetics Research Laboratories, Daiichi Sankyo Co., Ltd., Tokyo, Japan

The corrected Figures and Table appears below.

Page 9.

Fig. 3e

Page 10

Table 2, line 6 Coproporphyrin I.

Probe drug	Control	+P	+RIF300	Fold change ^a	+RIF600	Fold change ^a
Coproporphyrin I	nd	16.8 ± 1.6	38.1 ± 2.7**	2.3 ± 0.1	57.8 ± 3.8**,††	3.5 ± 0.1

Fig. 4 **a** Correlation between AUC_{0-24h} of GCDCA-S and CP-I. **b** Correlation between AUC_{0-24h} of CP-I and atorvastatin

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.