

## Retraction Note: Use of antidepressants and the risk of type 2 diabetes mellitus: a nested case–control study

Star Khoza · Jamie C. Barner · Thomas M. Bohman ·  
Karen Rascati · Kenneth Lawson · James P. Wilson

Published online: 21 August 2012  
© Springer Science+Business Media B.V. 2012

**Retraction to: Int J Clin Pharm (2012) 34(3):432–438  
DOI 10.1007/s11096-012-9608-4**

The International Journal of Clinical Pharmacy has been alerted to a case of duplicate publication in the following papers.

Int J Clin Pharm. 2012 Jun;34(3):432-8. Epub 2012 Jan 18.

Use of antidepressants and the risk of type 2 diabetes mellitus: a nested case–control study.

Khoza S, Barner JC, Bohman TM, Rascati K, Lawson K, Wilson JP.

Eur J Clin Pharmacol. 2011 Nov 26. [Epub ahead of print].

Use of antidepressant agents and the risk of type 2 diabetes.

Khoza S, Barner JC, Bohman TM, Rascati K, Lawson K, Wilson JP.

The International Journal of Clinical Pharmacy is retracting the paper. We sincerely apologize to the scientific community for any inconvenience that this might have caused.

---

The online version of the original article can be found under doi:[10.1007/s11096-012-9608-4](https://doi.org/10.1007/s11096-012-9608-4).

---

S. Khoza (✉)  
Department of Clinical Pharmacology, College of Health Sciences, University of Zimbabwe, PO Box A 178, Avondale, Harare, Zimbabwe  
e-mail: [skhoza53@gmail.com](mailto:skhoza53@gmail.com)

J. C. Barner · K. Rascati · K. Lawson · J. P. Wilson  
College of Pharmacy, The University of Texas at Austin, Austin, TX, USA

T. M. Bohman  
School of Social Work, The University of Texas at Austin, Austin, TX, USA