



Flavonoids of the Caryophyllaceae

Katarzyna Jakimiuk · Michael Wink · Michał Tomczyk



Received: 1 December 2020 / Accepted: 9 April 2021 / Published online: 17 April 2021
© The Author(s) 2021

Abstract The plant family Caryophyllaceae, commonly known as the pink family, is divided into 3 subfamilies and contains over 80 genera with more than 2600 species that are widely distributed in temperate climate zones. Plants belonging to this family produce a variety of secondary metabolites important in an ecological context; however, some of these metabolites also show health-promoting activities. The most important classes of phytochemicals include saponins, phytoecdysteroids, other sterols, flavonoids, lignans, other polyphenols, essential oils, and *N*-containing compounds such as vitamins, alkaloids or cyclopeptides. Flavonoids are polyphenolic compounds that remain one of the most extensively studied constituents of the Caryophyllaceae family. Numerous structurally diverse aglycones, including flavones, flavonols, flavonones (dihydroflavones),

flavonols, isoflavones, and their *O*- or *C*-glycosides, exhibit multiple interesting biological and pharmacological activities, such as antioxidant, anti-inflammatory, anti-oedemic, antimicrobial, and immunomodulatory effects. Thus, this review analysed the flavonoid composition of 26 different genera and more than 120 species of Caryophyllaceae for the first time.

Keywords Caryophyllaceae · Phytochemistry · Flavonoids · Secondary metabolites

Introduction

The Caryophyllaceae family, commonly known as the pink family, contains over 80 genera with more than 2600 species. The pink family is divided into 3 subfamilies, Paronychioideae, Alsinoideae, and Caryophylloideae, according to the presence or absence of stipules as well as the type of calyx and corolla. Plants of the Caryophyllaceae family are erect, prostrate, annual or perennial herbs or shrubs with simple cross-opposite leaves and swollen nodes. Tetramerous or pentamerous flowers are frequently gathered in panicle, raceme, or capitulum inflorescences (Hegnauer 1964; Kubitzki 1993; Schweingruber 2007).

K. Jakimiuk · M. Tomczyk (✉)
Department of Pharmacognosy, Faculty of Pharmacy with
the Division of Laboratory Medicine, Medical University
of Białystok, ul. Mickiewicza 2a, 15-230 Białystok,
Poland
e-mail: michal.tomczyk@umb.edu.pl

K. Jakimiuk
e-mail: katarzyna.jakimiuk@umb.edu.pl

M. Wink
Institute of Pharmacy and Molecular Biotechnology,
Heidelberg University, INF 364, 69120 Heidelberg,
Germany
e-mail: wink@uni-heidelberg.de

The subfamily Paronychoideae, containing the genera *Spergula* L., *Spergularia* Presl., *Polycarpon* L. *Herniaria* L., and *Paronychia* Mill., occurs mostly in warm and tropical parts of the world. The characteristic attributes of these plants are leaves with stipules and visible separation of calyx from the corolla. The lack of stipules and the unique corolla are typical for members of the subfamily Alsinoideae: The genera *Scleranthus* L., *Arenaria* L., *Sagina* L., *Cerastium* L., *Minuartia* L., *Stellaria* L., and *Colobanthus* Bartl. are widespread on all continents and are even present in Antarctica. Several species of the subfamily Caryophylloideae are field weeds that inhabit northern temperate climate regions. The specific structures of this subfamily are long calyx tubes that occur in *Agrostemma* L., *Maleandrium* Roehl., *Silene* Mill., *Gypsophila* L., and *Dianthus* L. (the largest genus). A great number of Caryophyllaceae species are grown as decorative landscape plants. Furthermore, many members of this family produce secondary metabolites with medicinal properties (Brockington et al. 2011; Volodin and Volodina 2015).

Diversity of phytochemicals in Caryophyllaceae

Caryophyllaceae are known to be a rich source of pharmacologically active secondary metabolites spanning several structural chemical classes. Secondary metabolites are important for plants as protective chemicals against herbivores (insects, molluscs, vertebrates) and microbial pathogens (fungi, bacteria, viruses), UV light, and other plants competing for light, water, and nutrients. In addition, many secondary metabolites serve as signalling compounds to attract pollinating and seed-dispersing animals and provide communication signals among plants and symbiotic microbes (Wink 2011).

The main secondary metabolites of Caryophyllaceae are saponins, phytoecdysteroids, other sterols, flavonoids, lignans, other polyphenols, essential oils, and *N*-containing compounds such as vitamins, alkaloids and cyclic peptides.

Methodology

The search strategy helps to define appropriate search string and identify the relevant thematic databases to collect the relevant scientific literature. The search databases for this review were SCOPUS, PubMed/MEDLINE, Web of Science (SCI-EXPANDED), Wiley Online Library, Taylor & Francis Online, Google Scholar, REAXYS Database, Science Direct/ELSEVIER, and EBSCO Discovery Service (EDS). They have been searched systematically for articles published from 1950 until 2020. The following syntax was used: TITLE-ABS-KEY as additional search engine in combinations of the above keywords like “Caryophyllaceae”, OR “genus” (each genus from the Caryophyllaceae family was introduced), OR “phenolic compounds”, OR “flavonoids”, OR “flavones”, OR “flavonols”, OR “flavonones”, OR “isoflavones”, OR *C*-flavonoids”, OR “Caryophyllaceae”, OR “saponins”, OR “phytoecdysteroids”, OR “essential oils”, OR “volatile compounds”, OR “sterols”, OR “*N*-containing compounds”, OR “alkaloids”, OR “cyclic peptides”, OR “vitamins”, OR “lignans”, OR “bioavailability”, OR “metabolism”, OR “biological activity”. Search terms had run in separate or with limited combinations that considered the requirements, or limitations, of the database used. Additionally, based on USDA Plant Database and Kew Science (Royal Botanic Gardens), we have been ascertaining the genera belonging to the Caryophyllaceae family (USDA Plant Database 2020; Kew Science 2020).

Triterpene saponins

Triterpene saponins constitute the greatest proportion of all phytochemicals known to be present in Caryophyllaceae. The structure of Caryophyllaceae saponins may vary with respect to genera within a family, as well as to plant organs. Oleanane-type saponins, such as gypsogenin, gypsogenic acid, quillaic acid (Fig. 1), 16 α -hydroxygypsogenic acid or their derivatives, constitutes the main group of saponins in these plants (Hegnauer 1989; Vincken et al. 2007; Böttger et al. 2011; Cheikh-Ali et al. 2019). For example, this class of compounds is synthesized in *Gypsophila altissima* (Chen et al. 2010a, b), *Gypsophila glomerata* (Gevrenova et al.




Fig. 1 The chemical structures of triterpene saponins in plants of Caryophyllaceae family. Gypsogenin (a), quillaic acid (b) succulentoside A (c), gypsophilin (d)

2018), *Gypsophila capillaris* (Elgamal et al. 1995), *Saponaria officinalis* (Koike et al. 1999), *Silene vulgaris* (Kim et al. 2015), *Vaccaria segetalis* (Koike et al. 1998), *Dianthus versicolor* (Ma et al. 2009), *Silene cucubalus* (Larhsini et al. 2003), *Paronychia chionaea* (Avunduk et al. 2007) and many other species (Hegnauer 1964; Böttger and Melzig 2011). Moreover, among triterpene saponins from Caryophyllaceae, ursane-type, hopane-type, and lupane-type saponins have also been reported (Vincken et al. 2007). For instance, succulentoside A (Fig. 1) and B, which are hopane-type saponins, were isolated from *Polycarpon succulentum* (Meselhy and Aboutabl 1997). Gypsophilin (Fig. 1), its glucosyl ester gypsophilinoside and sulfated lupane triterpenes were detected in *Gypsophila repens* (Elbandy et al. 2007).

Phytoecdysteroids

Phytoecdysteroids, structural analogues of the insect moulting hormone ecdysone, are another group of compounds commonly found in Caryophyllaceae. Several *Silene* Mill. species, e.g., *S. guntensis* (Mamadaliyeva et al. 2011), *S. antirrhina*, *S. chlorifolia*, *S.*

cretica, *S. disticha*, *S. echinata*, *S. italica*, *S. portensis*, *S. pseudotites*, *S. radicata*, *S. regia* (Meng et al. 2001), *S. viridiflora*, *S. linicola* (Mamadaliyeva et al. 2004), *S. nutans*, *S. otites*, and *S. tatarica* (Bathori et al. 1990), are rich sources of 20-hydroxyecdysone (Fig. 2). Along with 20-hydroxyecdysone, in the genus *Silene* Mill., a notably large number of structurally various phytoecdysteroids have been observed (Mamadaliyeva et al. 2014). Furthermore, plants of the genus *Coronaria* L. are potential producers of ecdysteroid compounds such as viticosterone E, α -ecdysone, taxisterone (Fig. 2), polypodine B, 20,26-dihydroxyecdysone, 2-deoxyecdysterone, and 20-hydroxyecdysone (Mamadaliyeva et al. 2008). Several ecdysteroids were also established in *Silene flos-cuculi* (syn. *Lychnis flos-cuculi*) (Báthori et al. 2001; Dinan et al. 2020). Based on TLC and HPLC analyses, the biotechnological regenerated shoots and roots of *L. flos-cuculi*, reveals the ability to accumulate 20-hydroxyecdysone and polypodine B (Thiem et al. 2016; Maliński et al. 2019).




Fig. 2 The chemical structures of phytoecdysteroids in Caryophyllaceae species. 20-Hydroxyecdysone (a), taxisterone (b) α -ecdysone (c), viticosterone E (d)

Essential oils and volatile compounds

Essential oils are widely distributed in the plant kingdom. This finding suggests that essential oils are also produced in flowering parts of taxa in the pink family. As essential oils are isolated by distillation, they contain a variety of volatile molecules—terpenes and terpenoids, phenol-derived aromatic components, and aliphatic constituents. Components of volatile oils isolated from *Dianthus acicularis* are chiefly 2-pentadecanone (Fig. 3) and 2-tridecanone, which are presumed to be responsible for the insect repellent activity of this plant (Kirillov et al. 2017). According to analyses of the major constituents of *Dianthus calocephalus* and *Dianthus carmelitarum* essential oils, the presence of heneicosane, docosane, tetracosane, phytol, 4,4-dimethyl-2-pentene, pentacosane, and hexahydrofarnesyl acetone (Yücel and Yayli 2018). Additionally, floral fragrance compounds were also established in other *Dianthus* L. species and *Saponaria officinalis* with the largest amounts of benzenoids, phenyl propanoids, and isoprenoids (Jürgens et al. 2003). Gas chromatography and gas chromatography combined with mass spectrometry (GLC-MS) examinations of aerial parts of *Silene morganae* revealed the presence of over 30




Fig. 3 The chemical structures of the main components of essential oils from Caryophyllaceae plants. 2-Pentadecanone (a), 2-tridecanone (b)

compounds with the highest content of monoterpene hydrocarbons being of terpenoids (Azadi and Sohrabi 2014). Furthermore, benzenoids followed by FADs seems to be the dominating compound classes of aromatic compounds in night-blooming or moth-pollinated flowers of *Silene* Mill. species (Jürgens et al. 2002; Jürgens 2004). Essential oils and their volatile components were also observed in *Minuartia recurva* (Jovanović et al. 2009), *Dianthus caryophyllus* (Nerio et al. 2010), *Dianthus cruentus* (Radulović et al. 2018), some *Silene* species (Dötterl and Jürgens 2005; Mamadalieva et al. 2014; Mihaylova et al. 2018), *Gypsophila bicolor* (Shafaghat and Shafaghatlonbar 2011), and two hermaphroditic *Schiedea* species (Powers et al. 2020).

Sterols

Sterols seem to be useful chemotaxonomic markers at the species level within families of the order Caryophyllales. Atypical for higher plants but predominant in the pink family, the sterol-type class of compounds Δ^7 -sterols represented by 22-dihydrospinasterol (Fig. 4) occur in *Gypsophila perfoliata* (Schmidt et al. 1996), *Gypsophila paniculata*, *Silene cucubalus*, *Arenaria serpyllifolia*, *Cerastium vulgarum*, *Cerastium arvense*, *Myosoton aquaticum*, *Minuartia caroliniana*, *Spergula arvensis*, *Saponaria officinalis*, *Dianthus armeria*, *Lychnis alba*, *Paronychia virginica* and *Scleranthus annuus* (Salt and Adler 1986). Recent research revealed the presence of the α -spinasterol 3-*O*- β -D-glucoside in the roots of *Psammosilene tunicoides* (Zhou et al. 2013) and the roots/rhizomes of *Silene tatarinowii* (Liang et al. 2019).




Fig. 4 The chemical structures of sterols in plant of Caryophyllaceae family. 22-Dihydrospinasterol (a), α -spinasterol 3-O- β -D-glucoside (b)

Cyclic peptides

Cyclic peptides, consisting of a maximum of 14 amino acid residues, are typical *N*-containing secondary metabolites from Caryophyllaceae (Ma et al. 2006). Genera containing cyclopeptides as major phytochemicals among all plants from this family seem to be *Dianthus* L., *Gypsophila* L., *Stellaria* L., and *Vaccaria* Mill. For example, the cyclic peptides gypsophins A–F were isolated from the roots of *Gypsophila oldhamiana* (Wang et al. 2013); the hexapeptides dianthins E, G, and H were found in the aerial parts of *Dianthus superbus* (Tong et al. 2012); and diandrines A–D (Fig. 5) and drymarins A–B occur in *Drymaria diandra* (Hsieh et al. 2004a, b; Ding et al. 2000). According to available data, seeds of *Vaccaria segetalis* are a valuable source of the penta- and hexapeptides segetalin B and segetalin A, respectively

(Itokawa et al. 1995; Wang et al. 2011). It is worth mentioning that this group of compounds is present in taxa of the subfamily Alsinoideae, which grow in Antarctica (Jia et al. 2004).

Alkaloids

Another group of nitrogen-containing secondary metabolites are alkaloids, which also occur in Caryophyllaceae to some degree. In particular, alkaloids belonging to the β -carboline group have been described (Dai et al. 2018). For instance, siliendines A–D were isolated from the aerial parts of *Silene seoulensis* (Seo et al. 2020), drymaritin from the whole plant material of *Drymaria diandra* (Hsieh et al. 2004a, b), oldhamiaines A and B from the roots of *Gypsophila oldhamiana* (Zhang et al. 2015), and




Fig. 5 The chemical structures of cyclic peptides in Caryophyllaceae species. Diandrine A (a), diandrine B (b) diandrine C (c)

arenarines A-D from *Arenaria kansuensis* (Wu et al. 1989; Bracher and Puzik 2004). Phytochemical investigation of the roots of *Stellaria dichotoma* led to the isolation of 23 various β -carboline-type alkaloids, including stellarines A-B, dichotomides I-XIV, dichotomines A, B, E (Fig. 6), and K, L, glucodichotomine B and 1-acetyl-3-methoxycarbonyl- β -carboline (Chen et al. 2010a, b; Luo et al. 2012). *Brachystemma calycinum* also produces alkaloids: Brachystemidines A-E were isolated from the roots of this plant (Cheng et al. 2002). Superbusines A and B, which are quinolone alkaloids, were detected in *Dianthus superbus* (Sun et al. 2019).

Vitamins

Analysis of plant-derived vitamins showed the presence of four tocopherols (α , β , γ , δ) with a different number of methyl substitutions in *Silene vulgaris* as well as vitamin C and phylloquinone, known as vitamin K1 (Fig. 7). Upon examination of *S. vulgaris*, the presence of the antioxidant β -carotene, a provitamin of vitamin A, was also reported (Vardavas et al. 2006; Morales et al. 2012; Mamadalieva et al. 2014). Moreover, β -carotene was reported in other Caryophyllaceae plants, e.g., in *Stellaria media* whose seeds contain vitamin B2 (riboflavin), vitamin B3 (niacin) and vitamin E (Slavokhotova et al. 2011; Taskin and Bitis 2013).

Phenolic compounds

Phenolic compounds constitute a large proportion of secondary metabolites in Caryophyllaceae plants. Phenolic acids are the main polyphenols produced by plants. However, only a few publications report on




Fig. 7 The chemical structures of vitamins present in Caryophyllaceae plants. Vitamin B2 (a), vitamin B3 (b), vitamin E (c)

phenolic compound isolation and identification in Caryophyllaceae. For instance, caffeic acid was obtained from aerial parts of *Silene* (syn. *Lychnis*) *flos-cuculi* (Tomczyk 2008), *p*-coumaric acid, dihydroferulic acid, and syringic acid were identified in the ground roots, stems, leaves, and flowers of *Gypsophila paniculata* (Chou et al. 2008); and *Dianthus* species are a source of gentisic acid, a commonly reported aromatic acid in green plants (Griffiths 1959). Fractionation of a *Gypsophila sphaerocephala* extract resulted in the isolation of 3,4-dihydroxybenzoic acid, syringic acid, *p*-hydroxybenzoic acid, and rosmarinic acid (Fig. 8) from the methanol extract and rosmarinic acid and syringic acid from the water extract (Altay et al. 2018). Additionally, the *Silene* Mill. genus is also known as a source of phenolic acids (Mamadalieva et al. 2014). Derivatives of cinnamic acid or benzoic acid and aromatic amino acids (anthranilic acid), so-called anthranilamides with phytoalexin-related activity, are commonly found in parts of *Dianthus caryophyllus* infected by pathogens (Niemann 1993).

Catechins (flavanol derivatives) are similar in structure to flavonols, except for the lack of a carbonyl




Fig. 6 The chemical structures of alkaloids in Caryophyllaceae plants. Dichotomine A (a), dichotomine B (b) dichotomine E (c)




Fig. 8 The chemical structures of phenolic acids in plants of Caryophyllaceae family. Rosmarinic acid (a), 3,4-dihydroxybenzoic acid (b) syringic acid (c), p-hydroxybenzoic acid (d)

group in the pyran ring (Heim et al. 2002). Plants of the Caryophyllaceae family were also screened for flavanols, but only a few species, including *Herniaria fontanessii* (Mbark et al. 1999) and *Arenaria kansuensis* (Liu et al. 2018), contained this group of compounds. The major flavanols, catechin, and epicatechin, act as strong antioxidant agents similar to other polyphenols (Iacopini et al. 2008).

Lignans, insoluble elements of certain cell walls, are rather uncommon phytochemicals in Caryophyllaceae, except for *Pteranthus dichotomus*, which contains 8-oxo-pinoresinol (Allaoua et al. 2016).

Unlike the many taxa of the order Caryophyllales that produce betalains as coloured flower pigments, Caryophyllaceae produce anthocyanins: cyanidin glycoside derivatives were identified in *Silene dioica* (Kamsteeg et al. 1976; Kamsreoo et al. 1980) and *S. armeria* (Mamadalieva et al. 2014). Cyclic malyl anthocyanins were isolated from deep pink and red-purple *Dianthus caryophyllus* flower petals (Nakayama et al. 2000). Moreover, the genus *Lychnis* is a source of the anthocyanin aglycones named

anthocyanidins, such as cyanidin, peonidin, and pelargonidin (Fig. 9), as well as their glycoside derivatives (Kuwayama et al. 2005).

Among the many polyphenolic phytoconstituents occurring in this family, tannins are also present and have physiological activity against herbivores. Tannins were detected in some *Minuartia* species (Zaychenko and Zernov 2017), *Stellaria laeta* (Jung et al. 1979), *Polycarpaea corymbosa* (Balamurugan et al. 2013), *Drymaria cordata* (Baruah et al. 2009), *Silene vulgaris* (Kim et al. 2015), *Silene compacta* (Bakroglu et al. 2014), and *Spergula fallax* (Aldhebiani and Mufarah 2017).

However, flavonoid compounds remain one of the most extensive groups of polyphenols in Caryophyllaceae, and novel compounds are yet to be identified. The aglycones and their glycosides are probably present in almost all plants.

Flavonoids of the Caryophyllaceae and their main biological activities

Flavonoids are low-molecular-weight secondary plant metabolites composed of two benzene rings and one heterocyclic pyran ring that are chemically divided into groups according to their chemical substitutions. Flavonoid moieties can be modified by glycosylation, hydrogenation, hydroxylation, and methylation as well as malonylation and sulfatation. The chemical and biological activities of flavonoids and their derivatives are connected with their structure and the position of various substitutions on the molecule. The general activity of polyphenols concerns the reactivity of their phenolic OH groups. The hydroxyl groups can dissociate under physiological conditions to negatively charged phenolate ions. Thus, polyphenols can interact with proteins by forming hydrogen bridges and, more importantly, ionic bonds with positively charged amino groups. As a consequence, the




Fig. 9 The chemical structures of the anthocyanins in Caryophyllaceae. Cyanidin (a), peonidin (b) pelargonidin (c)

bioactivity of proteins can be directly changed when a polyphenol binds to a receptor side or active centre of an enzyme. Polyphenols, especially those with several phenolic OH groups (such as rosmarinic acid or tannins), can change the 3D structure of proteins and impair their bioactivity. Because of these interactions, polyphenols affect many proteins in the human body and in microbes that are medically relevant. This is the mechanism by which plant polyphenols are medically active (Wink 2015; van Wyk and Wink 2017).

The biological activities of flavonoids may be also connected with their metabolites, which are produced in vivo. The gastrointestinal tract reveals primary role in the absorption, distribution, metabolism and excretion of flavonoids, which are substrates for conjugating and hydrolyzing enzymes in the small intestine, liver, and colon to *O*-glucuronides, *O*-methyl and sulfate esters. Firstly, metabolism of flavonoids take place in the small intestine followed by the liver where they are transformed and then produced glucuronides and sulfate derivatives. Flavonoid compounds that reach the colon are catabolized to low molecular weight phenolic acids by the intestinal microflora (Thilakarathna and Rupasinghe 2013). An anaerobic bacteria found in human gastrointestinal tract (e.g. *Eubacterium ramulus*) splits the ring structures of several flavonols and flavones leading to the formation of aglycones and common phenolics intermediates consisting of hydroxyphenylacetic, hydroxyphenylpropionic, acetate, and butyrate acids with varying degrees of hydroxylation (Blaut et al. 2003; Serra et al. 2012; Pei et al. 2020). The amount of urinary excretion demonstrates that the colonic catabolites are absorbed into the portal vein and this way run over the body in the circulatory system (Crozier et al. 2010). The flavonoid glucuronides and sulfate derivatives facilitate their excretion through urine and bile (Thilakarathna and Rupasinghe 2013). Urinary excretion of < 1.0% confirms that *C*-flavones are poorly absorbed, and 10–88% recovery from feces indicates that they may be resistant to degradation by gut bacteria in rats (Ma et al. 2010). As with flavone *O*-glycosides, the *C*-glycosides are less bioavailable in humans than in rats. Nevertheless, it is known that the absorption of dietary flavonoids may be affected by the food matrix, the metabolic processes mediated by the liver, intestine, kidneys, as well as colon microbiota (Hollman 2004; Viskupičová et al. 2008;

Hostetler et al. 2017; Cosme et al. 2020; Di Lorenzo et al. 2021).

To the best of our knowledge, apigenin, found in 28 species, is the major flavone in Caryophyllaceae plants. The apigenin exhibits cancer chemopreventive activity such as antiproliferative effects on human breast cancer cells, inhibition of cell growth by apoptosis in cervical carcinoma, or selective apoptotic effects in monocytic and lymphocytic leukaemias (Shukla and Gupta 2010; Imran et al. 2020). A similar number of species contain another widely distributed aglycone—luteolin. As with many other polyphenols, luteolin is a powerful antioxidant that can prevent inflammation and allergies and suppress the expression of cancer-promoting proteins (Imran et al. 2019a, b). Other important flavones are the luteolin 8-*C*-glucoside and apigenin 8-*C*-glucoside, orientin and vitexin, respectively. Plants rich in orientin are often used in traditional medicine for the treatment of respiratory disorders, pharyngitis, skin disorders, common cold, and mild anxiety (Grundmann et al. 2008; Lam et al. 2016). In addition, luteolin 8-*C*-glucoside acts as an antioxidant, antiaging, anti-inflammatory, cardioprotective, radioprotective, and neuroprotective agent (Uma Devi et al. 2000; Praveena et al. 2014; Lam et al. 2016). Vitexin, successfully isolated from Caryophyllaceae, exhibits various medicinal properties, such as fat reduction, improved glucose metabolism, hepatoprotection, neuroprotection, cardioprotection, and even anticancer activity (Ganesan and Xu 2017; Peng et al. 2020).

Kaempferol exhibits multiple biological effects, such as antioxidant, anti-inflammatory, antidiabetic, antiaging, and antimicrobial effects, and is being applied in the chemotherapy of skin, liver, and colon tumours (Zhu et al. 2018; Cho and Park 2013; Imran et al. 2019a, b). Furthermore, kaempferol can be used in the treatment of cardiovascular diseases, degenerative disorders, diabetes, and microbial contamination diseases (Imran et al. 2018, 2019a). The flavonol aglycone, quercetin can function as an antioxidant as well as a blood pressure-lowering and anticancer agent (Kukongviriyapan et al. 2012; Egert et al. 2009). Moreover, quercetin can decrease the levels of proinflammatory cytokines, e.g., interleukin 6, 8, 1 β , and TNF α (Wang et al. 2016). Rutin, a 3-*O*-rutinoside derivative of quercetin established in 18 different species of the Caryophyllaceae, is also often used in studies due to its extensive therapeutic properties: The

health-promoting effects of rutin are linked with antioxidant, cytoprotective, neuroprotective, vasoprotective, and cardioprotective activities (Kim et al. 2009; Ganeshpurkar and Saluja 2017). The results from some studies also indicated a positive effect of rutin on Parkinson's and Alzheimer's diseases (Gullón et al. 2017). The main strategies for a neurodegenerative disease therapy involves the reduction of reactive oxygen species and amyloid beta-protein production, and the activation of mechanisms of neuronal death (de Andrade Teles et al. 2018).

The main biological activities of hesperidin isolated from *Herniaria hemistemon* (Elhagali et al. 2019) are chemotherapeutic, antiallergic, anti-inflammatory, endocrine, cardiovascular, and organ-protective effects (Kumar et al. 2008; Zanotti et al. 2013; Ganeshpurkar and Saluja 2019). The aglycone naringenin exhibited multiple therapeutic effects associated with its free radical-scavenging properties. Depending on the concentration and method of administration, naringenin can be useful in the treatment of viral, bacterial, and inflammatory diseases and obesity (Ke et al. 2016; Kozłowska et al. 2017; Salehi et al. 2019). In addition, naringenin was tested for its potential anticancer activity and as a cardioprotective agent (Salehi et al. 2019). A wide range of therapeutic properties of naringenin, a 7-hesperidoside derivative of naringenin, include the treatment of metabolic syndrome, oxidative stress, and conditions of the central nervous system (Sachdeva et al. 2014; Dhanya et al. 2015; Chen et al. 2016).

A medically useful group of flavonoids are isoflavones, which are also known as phytoestrogens (Heim et al. 2002). These compounds can bind to receptors of oestrogen and oestrogen hormone binding protein and inhibit an important enzyme of angiogenesis and tumour formation, tyrosinase (Wink 2015). It was concluded that plants rich in isoflavones are effective in treating cardiovascular and osteoporosis disorders as well as in reducing postmenopausal symptoms (Clarkson 2002; Atkinson et al. 2004; Vitale et al. 2013). To date, the distribution of genistein and daidzein is common in several legumes of the Fabaceae family, such as soybean (Bustamante-Rangel et al. 2018). However, there are reports of the presence of genistein in a species of the Caryophyllaceae family, e.g., *Stellaria dichotoma* or *Stellaria holostea* (Mikšátková et al. 2014).

Our approach included screening for flavonoid aglycones and their highly glycosylated derivatives within Caryophyllaceae family (Cook and Samman 1996). The flavonoid aglycone and glycoside group remains one of the most extensive groups of polyphenols in Caryophyllaceae. Most of these compounds occur in *Silene* L., *Dianthus* L. (Obmann et al. 2011a, b; Boguslavskaya et al. 1983), *Gypsophila* L. (Zhang et al. 2011a, b; Zheleva-Dimitrova et al. 2018), *Stellaria* L. (Mikšátková et al. 2014), *Spergularia* Presl. (Ferrerres et al. 2011) and *Herniaria* L. (Elhagali et al. 2019; El Mabruki et al. 2014). Nevertheless, flavonoid compounds are probably present in almost all plants. We assembled information regarding their presence in 26 genera and over 120 species of the Caryophyllaceae family (see Table 1).

Flavones

One of the most pharmacologically valuable flavonoid classes present is that comprising flavones, which can be synthesized by various pathways, depending on whether they contain *C*- or *O*-glycosylation, *O*-methylation acylation, and hydroxylated B-ring. These compounds undergo characteristic reactions ascribed to three functional structures—hydroxyl and carbonyl groups and a double bond (Singh et al. 2014; Panche et al. 2016). Their natural distribution is demonstrated for almost all plant tissues (Figs. 10, 11, 12).

Flavonols

An additional class of flavonoids commonly found in Caryophyllaceae is that comprising flavonols, including kaempferol, quercetin and its glycoside rutin (quercetin 3-*O*-rutinoside). Flavonols, compared to flavones, carry an additional hydroxyl group in the pyran ring (Panche et al. 2016) (Fig. 13).

Flavonones (dihydroflavones)

Flavonones (dihydroflavones) differ from flavones by the lack of a double bond in the pyran ring. Hesperidin, naringenin, and its glycoside naringin (naringenin 7-hesperidoside) are commonly found in citrus fruits (Panche et al. 2016), but they can also be found in certain species of the pink family.

Table 1 Flavonoids compounds of the Caryophyllaceae family

Genus	Compounds	References
<i>Agrostemma githago</i>	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	Richardson (1978)
	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	
<i>Alsinidendron trinerve</i>	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77)	
	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	
<i>Arenaria kansuensis</i>	apigenin (48)	Liu et al. (2018)
	luteolin 3'-methyl ether (chrysoeriol) (128)	
	luteolin 7- <i>O</i> - β -D-glucoside (cynaroside) (37)	
	tricin 7- <i>O</i> - β -D-glucoside (160)	
	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77)	
	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	
	chrysoeriol 8- <i>C</i> - β -D-glucoside (scoparin) (129)	
	luteolin (1)	Liu et al. (2018), Cui et al. (2017b)
	homoeriodictyol (217)	
	kaempferol (172)	Liu et al. (2018), Tong et al. (2014)
	quercetin (189)	
	tricin (159)	Wu et al. (1990), Liu et al. (2018), Cui et al. (2017a), Cui et al. (2017b), Cui et al. (2018)
	chrysoeriol 6- <i>C</i> - β -D-glucoside (isoscoparin) (141)	
	tricin 4'- <i>O</i> -(<i>C</i> -veratrolyglycol) ether (151)	Cui et al. (2019)
<i>Arenaria saxatilis</i>	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77)	Darmograi (1979)
<i>A. serpyllifolia</i>	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	
<i>A. stenophyla</i>		
<i>A. juncea</i>	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	
<i>A. lychmidea</i>	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	

Table 1 continued

Genus	Compounds	References
	isomollupentin 7,2''-di- <i>O</i> -glucoside (102)	Dubois et al. (1983)
	isomollupentin 7- <i>O</i> -glucoside-2''- <i>O</i> -arabinoside (103)	
	isomollupentin 7- <i>O</i> -glucoside-2''- <i>O</i> -xyloside (104)	
	isovitexin 7- <i>O</i> -glucoside-2''- <i>O</i> -arabinoside (122)	
	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	Dubois et al. (1985)
<i>Dianthus acieularis</i>	apigenin 6- <i>C</i> - <i>anti</i> - α -D-glucoside (isoneoavroside) (79)	Boguslavskaya et al. (1983)
	apigenin 6- <i>C</i> - <i>syn</i> - α -D-glucoside (neoavroside) (80)	
<i>Dianthus anatoficus</i>	kaempferol (172)	Richardson (1978)
	quercetin (189)	
	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	
	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	
<i>Dianthus japonicus</i>	isovitexin 7- <i>O</i> - β -D-glucoside (saponarin) (105)	Nakano et al. (2011)
	saponarin 2''- <i>O</i> - α -L-rhamnoside-6'''- <i>O</i> -7,8-dihydroferulate (106)	
<i>Dianthus hoeltzeri</i>	apigenin 6- <i>C</i> - <i>anti</i> - α -D-glucoside (isoneoavroside) (79)	Boguslavskaya et al. (1983), Obmann et al. (2011a, b)
	apigenin 6- <i>C</i> - <i>syn</i> - α -D-glucoside (neoavroside) (80)	
	chrysoeriol 6- <i>C</i> - <i>syn</i> - α -D-glucoside (142)	
<i>Dianthus dicolor</i>		
<i>Dianthus squarrosus</i>	isovitexin 4'- <i>O</i> - β -D-glucoside (isosaponarin) (124)	Boguslavskaya et al. (1983)
<i>Dianthus superbus</i>	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	Seraya et al. (1978), Obmann et al. (2011a, b)
	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	
<i>Dianthus platycodon</i>	kaempferol 3- <i>O</i> -(β -D-glucosyl- β -D-glucoside) (173)	Boguslavskaya (1976), Obmann et al. (2011a, b)
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
<i>Dianthus ramosissimus</i>	chrysoeriol 6- <i>C</i> - <i>anti</i> - α -D-glucoside (143)	Obmann et al. (2011a, b)
<i>Dianthus pseudosquarrosus</i>	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77)	
	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	
	isovitexin 4'- <i>O</i> - β -D-glucoside (isosaponarin) (124)	
	luteolin 5- <i>O</i> -glucoside (36)	
	luteolin 7- <i>O</i> -diglucoside (38)	
	luteolin 7- <i>O</i> - β -D-glucoside (cynaroside) (37)	

Table 1 continued

Genus	Compounds	References
<i>Dianthus versicolor</i>	apigenin (48)	Obmann et al. (2011a, b), Obmann et al. (2012)
	apigenin 4'-methyl ether (acacetin) (151)	
	luteolin (1)	
	luteolin 3'-methyl ether (chrysoeriol) (128)	
	luteolin 4'-methyl ether (diosmetin) (154)	
	isoorientin 7-O-galactoside (22)	
	isoorientin 7-O-rhamnosyl-galactoside (23)	
	isoorientin 7-O-rutinoside (24)	
	isoscoparin 7-O-galactoside (147)	
	isoscoparin 7-O-rhamnosyl-galactoside (148)	
	isoscoparin 7-O-rutinoside (149)	
	isovitexin 7-O- β -D-glucoside (saponarin) (105)	
	isovitexin 7-O-rhamnosyl-galactoside (107)	
	isovitexin 7-O-rutinoside (109)	
	apigenin C-hexosyl-O-hexoside maly ester (110)	
	luteolin C-hexosyl-O-hexoside maly ester (39)	
	apigenin 6-C- β -D-glucoside (isovitexin) (77)	
	apigenin 7-O- β -D-glucoside (cosmosiin) (111)	
	isovitexin 2''-O-rhamnoside (99)	
luteolin 7-O- β -D-glucoside (cynaroside) (37)		
<i>Dianthus caryophyllus</i>	kaempferide 3-O- β -D-glucosyl-(1 \rightarrow 2)-O-[α -L-rhamnosyl-(1 \rightarrow 6)]- β -D-glucoside (188)	Curir et al. (2001), Obmann et al. (2011a, b)
	kaempferide 3-O-[2 ^G - β -D-glucosyl]- β -rutinoside (187)	Curir et al. (2005), Obmann et al. (2011a, b)
	quercetin 3-[6-O-(α -L-arabinosyl)- β -D-glucoside] (peltatoside) (194)	Curir et al. (2003), Obmann et al. (2011a, b) Al-Snafi (2017)
	apigenin 6-C-glucosyl-7-O-(6-maly-glucoside) (112)	Fukui et al. (2003), Obmann et al. (2011a, b)
	kaempferol 3-O-[6'''-rhamnosyl-2'''-(6-maly-glucosyl)]-glucoside (174)	
	kaempferol 3-O-(6'''-rhamnosyl-2'''-glucosyl)-glucoside (175)	
	kaempferol 3-O-(β -D-glucosyl- β -D-glucoside) (173)	Ogata et al. (2004), Obmann et al. (2011a, b), Stich et al. (1992)
	apigenin 6,8-di-C- β -D-glucoside (vicenin-2) (53)	Galeotti et al. (2008a, b), Obmann et al. (2011a, b)
	kaempferol 3-O- β -D-rutinoside (nicotiflorin) (176)	
	kaempferol 3-O- β -D-glucosyl-(1''' \rightarrow 2'')-O-[α -L-rhamnosyl-(1''' \rightarrow 6''')]- β -D-glucoside (178)	Galeotti et al. (2008a, b), Galeotti et al. (2008a), Obmann et al. (2011a, b), Iwashina et al. (2010), Al-Snafi (2017)
	kaempferol 3-O- β -D-glucosyl-(1''' \rightarrow 2'')-O- β -D-glucosyl-(1 ^V \rightarrow 2''')-O-[α -L-rhamnosyl-(1 ^{IV} \rightarrow 6'')]- β -D-glucoside (177)	Galeotti et al. (2008a), Obmann et al. (2011a, b)
	kaempferol (172)	Stich et al. (1992)
	naringenin (220)	
	kaempferol 4'-methyl ether (kaempferide) (186)	Martinetti et al. (2010)
	kaempferol 3-O-neohesperidoside (179)	Iwashina et al. (2010)
	kaempferol 3-O-sophoroside (sophoraflavonololide) (185)	

Table 1 continued

Genus	Compounds	References	
<i>Dianthus deltoides</i>	luteolin (1)	Obmann et al. (2011a, b)	
	apigenin 6- <i>C-anti-α</i> -D-glucoside (isoneoavroside) (79)		
	apigenin 6- <i>C-syn-α</i> -D-glucoside (neoavroside) (80)		
	chrysoeriol 4'- <i>O-β</i> -D-glucoside (150)		
	luteolin 3'-methyl ether (chrysoeriol) (128)		
<i>Dianthus arenarius</i>	luteolin 4'- <i>O-β</i> -D-glucoside (40)	Boguslavskaya et al. (1983)	
	apigenin 4'- <i>O</i> -glucoside (125)		
<i>D. crinitus</i>	apigenin 6- <i>C-β</i> -D-glucoside (isovitexin) (77)	Richardson (1978)	
<i>D. tetralapsis</i>	apigenin 8- <i>C-β</i> -D-glucoside (vitexin) (49)		
	luteolin 4'- <i>O-β</i> -D-glucoside (40)		
<i>Dianthus grandiflora</i>	luteolin 6- <i>C-β</i> -D-glucoside (isorientin) (18)	Richardson (1978)	
	luteolin 8- <i>C-β</i> -D-glucoside (orientin) (2)		
<i>Drymaria diandra</i>	kaempferol (172)	Ding et al. (1999), Brahmachari and Gorai (2006)	
	quercetin (189)		
	drymariatin A (164)	Ding et al. (2005)	
	6- <i>trans</i> -{2''- <i>O</i> -(rhamnosyl)}-ethenyl-5,7,4'-trihydroxyflavone (170)		
	drymariatin B (165)	Nono et al. (2016)	
	drymariatin C (166)		
	drymariatin D (167)		
	5,4'-dihydroxy-7-methoxyflavone-6- <i>C</i> -(2''- <i>O-α</i> -L-rhamnosyl)- <i>β</i> -D-glucoside (87)		
<i>Drypis spinosa</i>	5,7,3',4'-tetrahydroxyflavone-6- <i>C</i> -(2''- <i>O-α</i> -L-rhamnosyl)- <i>β</i> -D-glucoside (45)	Hsieh et al. (2004a, b), Mandal et al. (2009)	
	apigenin 6- <i>C-β</i> -D-glucoside (isovitexin) (77) diandraflavone (168)		
	torosaflavone A (169)		
<i>Gymnocarpus decander</i>	naringenin (220)	Kremer et al. (2021)	
	quercetin (189)		
	quercetin 3- <i>O-α</i> -L-rutinoside (rutin) (190)		
<i>Gymnocarpus decander</i>	isorhamnetin 3- <i>O</i> -[(2'''- <i>O</i> -acetyl - <i>β</i> -D-xylosyl-(1 → 6)-[<i>β</i> -D-apiofuranosyl-(1 → 2)]- <i>β</i> -D-glucoside (212)	Bechlem et al. (2017)	
	isorhamnetin 3- <i>O</i> -2'''- <i>O</i> -acetyl - <i>β</i> -D-xylosyl-(1 → 6)- <i>β</i> -D-glucoside (213)		
	quercetin 3- <i>O</i> -(2'''- <i>O</i> -acetyl - <i>β</i> -D-xylosyl-(1 → 6)- <i>β</i> -D-glucoside (198)		
	apigenin (48)		Zitouni (2017)
	kaempferol (172)		
	luteolin (1)		
	myricetin 3'- <i>O</i> -methyl ether (laricitrin) (214)		
	naringenin (220)		
	kaempferol 3- <i>O-β</i> -D-rutinoside (nicotiflorin) (176)		
	luteolin 7- <i>O-β</i> -D-glucoside (cynaroside) (37)		
	quercetin 3- <i>O-β</i> -D-galactoside (hyperoside) (191)		
quercetin 3- <i>O-α</i> -L-rutinoside (rutin) (190)			

Table 1 continued

Genus	Compounds	References
	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	El-Hawary et al. (2020)
	quercetin (189)	Zitouni (2017), Mubarek (2019), El-Hawary et al. (2020)
	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77)	Mubarek (2019)
	rivularin (162)	
	quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192)	Zitouni (2017), Mubarek (2019)
	quercetin 3'-methyl ether (isorhamnetin) (204)	
<i>Gypsophila altissima</i>	isovitexin 7- <i>O</i> - β -D-glucoside (saponarin) (105)	Zdraveva et al. (2015)
<i>Gypsophila arrosti</i>	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	Altay et al. (2019)
<i>Gypsophila aucheri</i>	naringenin (220)	Altay (2018)
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
<i>Gypsophila repens</i>	luteolin 7- <i>O</i> - α -L-arabinosyl-6- <i>C</i> - β -glucoside (25)	Elbandy et al. (2007)
<i>Gypsophila elegans</i>	isoorientin 2''- <i>O</i> -arabinoside (19)	Zhang et al. (2011a, b), Huang et al. (2012), Lin et al. (2016)
	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	Lin et al. (2015), Tu et al. (2019)
	isovitexin 7- <i>O</i> - β -D-glucoside (saponarin) (105)	Zhang et al. (2011a, b), Zdraveva et al. (2015)
	apigenin 6- <i>C</i> -[β -D-xylosyl-(1''' \rightarrow 2'')- β -D-galactoside]-7- <i>O</i> - β -D-glucoside (113)	Zhang et al. (2011a, b)
	apigenin 7- <i>O</i> -sophoroside (114)	
	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	
	isovitexin 2''- <i>O</i> -glucoside (meloside A) (85)	
<i>Gypsophila eriocalyx</i>	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	Altay et al. (2019)
<i>Gypsophila trichotoma</i>	isovitexin 7- <i>O</i> - β -D-glucoside (saponarin) (105)	Vitcheva et al. (2011), Simeonova et al. (2014), Zheleva-Dimitrova et al. (2018)
	quercetin 3- <i>O</i> - β -D-galactoside (hyperoside) (191)	Krasteva et al. (2008)
	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	Krasteva et al. (2008), Zheleva-Dimitrova et al. (2018)
	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	
	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	
	apigenin (48)	Zheleva-Dimitrova et al. (2018)
	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77)	
	isorhamnetin 3- <i>O</i> - β -D-glucoside (205)	
	luteolin 2''- <i>O</i> -pentosyl-6- <i>C</i> -hexoside (29)	
	luteolin 4'-methyl ether (diosmetin) (154)	
	luteolin 7- <i>O</i> - β -D-glucoside (cynaroside) (37)	

Table 1 continued

Genus	Compounds	References
<i>Gypsophila glomerata</i>	apigenin (48) apigenin 2''-O-acetylpentosyl-6-C-hexoside (88) apigenin 2''-O-diacetylpentosyl-6-C-hexoside (89) apigenin 2''-O-pentosyl-6-C-hexoside (90) apigenin 6-C-β-D-glucoside (isovitexin) (77) apigenin 8-C-β-D-glucoside (vitexin) (49) diosmetin 2''-O-acetylpentosyl-6-C-hexoside (155) isorhamnetin 3-O-β-D-glucoside (205) isovitexin 7-O-β-D-glucoside (saponarin) (105) kaempferol 3-O-β-D-glucoside (astragalin) (180) kaempferol 3-O-β-D-rutinoside (nicotiflorin) (176) luteolin 2''-O-pentosyl-6-C-hexoside (29) luteolin 4'-methyl ether (diosmetin) (154) luteolin 6-C-β-D-glucoside (isoorientin) (18) luteolin 7-O-β-D-glucoside (cynaroside) (37) luteolin 8-C-β-D-glucoside (orientin) (2) luteolin methyl-2''-O-pentosyl-6-C-hexoside (30)	Zheleva-Dimitrova et al. (2018)
<i>Gypsophila tuberculosa</i>	quercetin 3-O-α-L-rutinoside (rutin) (190)	Altay et al. (2019)
<i>Gypsophila sphaerocephala</i>	quercetin 3-O-α-L-rutinoside (rutin) (190)	Altay et al. (2018)
<i>Gypsophila paniculata</i>	isovitexin 7-O-β-D-glucoside (saponarin) (105)	Zdraveva et al. (2015)
<i>Gypsophila perfoliata</i>	apigenin (48) apigenin 2''-O-hexosyl-6-C-hexoside (91) apigenin 2''-O-pentosyl-6-C-hexoside (90) apigenin 6,8-di-C-β-D-glucoside (vicenin-2) (53) apigenin 6-C-β-D-glucoside (isovitexin) (77) apigenin 6-C-hexosyl-8-C-pentoside (56) apigenin 8-C-β-D-glucoside (vitexin) (49) diosmetin 6-C-hexosyl-8-C-pentoside (156) isorhamnetin 3-O-β-D-glucoside (205) isovitexin 4'-O-β-D-glucoside (isosaponarin) (124) luteolin 2''-O-hexosyl-6-C-hexoside (31) luteolin 4'-methyl ether (diosmetin) (154) luteolin 6-C-hexosyl – 8-C-pentoside (32) luteolin 6-C-β-D-glucoside (isoorientin) (18) luteolin 7-O-β-D-glucoside (cynaroside) (37) luteolin 8-C-β-D-glucoside (orientin) (2)	Zheleva-Dimitrova et al. (2018)

Table 1 continued

Genus	Compounds	References
<i>Herniaria hirsuta</i>	quercetin (189)	Richardson (1978)
	isorhamnetin 3- <i>O</i> -rutinoside (narcissin) (206)	Van Dooren et al. (2016)
	quercetin 3- <i>O</i> -(2''- <i>O</i> - α -L-rhamnosyl)- β -D-glucuronoside (195)	
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
<i>Herniaria fontanessii</i>	apigenin (48)	Tlili et al. (2019)
	kaempferol (172)	
	naringenin (220)	
	quercetin (189)	
	quercetin 3- <i>O</i> - β -D-galactoside (hyperoside) (191)	
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
	isorhamnetin 3- <i>O</i> -robinobioside (207)	Mbark et al. (1999)
<i>Herniaria glabra</i>	isorhamnetin 3'- <i>O</i> -feruloyl-3- <i>O</i> -robinobioside (208)	
	apigenin (48)	El Mabruki et al. (2014)
	quercetin 3- <i>O</i> - β -D-galactoside (hyperoside) (191)	
	luteolin (1)	Maleš et al. (2013)
	isorhamnetin 3- <i>O</i> -rutinoside (narcissin) (206)	Kozachok et al. (2018)
	kaempferol 3- <i>O</i> - β -D-rutinoside (nicotiflorin) (176)	
	quercetin 3- <i>O</i> -[(D-apio- β -D-furanosyl-(1 \rightarrow 2)- <i>O</i> -[α -L-rhamnosyl-(1 \rightarrow 6)]- β -D-glucoside (apiorutin) (200)	
	quercetin (189)	Kulevanova et al. (2003), El Mabruki et al. (2014)
quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192)		

Table 1 continued

Genus	Compounds	References
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	Maleš et al. (2013), Kozachok et al. (2018), El Mabruki et al. (2014)
<i>Herniaria hemistemon</i>	apigenin (48)	Elhagali et al. (2019)
	kaempferol (172)	
	naringenin (220)	
	quercetin (189)	
	apigenin 4'-methyl ether (acacetin) (151)	
	apigenin 6- <i>C</i> - α -L-arabinosyl-8- <i>C</i> - β -D-galactoside (57)	
	apigenin 6- <i>C</i> -glucosyl-8- <i>C</i> -rhamnoside (58)	
	apigenin 6-rhamnosyl-8-glucoside (59)	
	apigenin 7- <i>O</i> - β -D-glucoside (cosmosiin) (111)	
	apigenin 7- <i>O</i> -neohesperidoside (rhoifolin) (115)	
	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	
	cyanidanon 4'-methyl ether (hesperetin) (218)	
	hesperetin 7- <i>O</i> - α -L-rutinoside (hesperidin) (219)	
	kaempferol 3,7-dirhamnoside (kaempferitrin) (181)	
	kaempferol 3- <i>O</i> -glucoside-2''- <i>p</i> -coumaroyl (182)	
	kaempferol 4'-methyl ether (kaempferide) (186)	
	kaempferol 7- <i>O</i> -hesperidoside (183)	
	luteolin 6- <i>C</i> -arabinosyl-8- <i>C</i> -glucoside (3)	
	luteolin 6- <i>C</i> -glucosyl-8- <i>C</i> -arabinoside (21)	
	naringenin 7- <i>O</i> - α -L-hesperidoside (naringin) (221)	
	quercetin 3- <i>O</i> -glucoside-7- <i>O</i> -rhamnoside (201)	
	quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192)	
	quercetin 3- <i>O</i> - α -L-rhamnoside (quercetrin)	
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
	quercetin 7-methyl ether (rhamnetin) (202)	
<i>Herniaria polygama</i>	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	Boguslavskaya et al. (1985a, b)
<i>Herniaria ciliolata</i>	isorhamnetin 3- <i>O</i> -rutinoside (narcissin) (206)	Królikowska et al. (1983)
	quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192)	
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
	rhamnazin 3- <i>O</i> -glucoside (209)	
	rhamnazin 3- <i>O</i> -rutinoside (polygalacin) (210)	
	rhamnetin 3- <i>O</i> -glucoside (193)	

Table 1 continued

Genus	Compounds	References
<i>Herniaria mauritanica</i>	kaempferol 3- <i>O</i> - β -D-glucoside (astragalin) (180) quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192)	Cheriti and Sekkoum (1996)
<i>Illecebrum verticillatum</i>	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2) luteolin 6- <i>C</i> - β -D-glucoside (isorientin) (18)	Richardson (1978)
<i>Lychnis senno</i>	chrysoeriol 6- <i>C</i> - β -D-glucoside (isoscoparin) (141) isoorientin 2''- <i>O</i> -rhamnoside (20) isovitexin 2''- <i>O</i> -rhamnoside (99) isovitexin 5- <i>O</i> -acetyl-2'- α -rhamnoside (92)	Shinjiro et al. (2009), Devkota et al. (2013); Maliński et al. (2014)
<i>Lychnis coronaria</i>	chrysoeriol 6- <i>C</i> - β -D-glucoside (isoscoparin) (141) tricin 7- <i>O</i> - β -D-glucoside (160)	Bahar et al. (2008), Maliński et al. (2014)
<i>Lychnis chalconica</i>	apigenin 6,8-di- <i>C</i> - β -D-glucoside (vicenin-2) (53) apigenin 8- <i>C</i> - α -D-glucoside (neovitexin) (60)	Smolyakova et al. (2010), Amosova et al. (2019)
<i>Minuartia rossi</i>	apigenin 6- <i>C</i> -arabinosyl-diglucoside (93)	Wolf et al. (1979)
<i>M. elegans</i>	apigenin 6- <i>C</i> -glucoside (isovitexin) (77)	
<i>M. austromontana</i>	apigenin 6- <i>C</i> -triglucoside (94) kaempferol 3- <i>O</i> - β -D-sophoroside (sophoraflavonolioside) (185) kaempferol 3- <i>O</i> -glucoside-2''- <i>p</i> -coumaroyl (182) quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192) quercetin 3- <i>O</i> - β -D-glucosyl- <i>O</i> -galactoside (196) quercetin 3- <i>O</i> - β -D-sophoroside (197)	
<i>Paronychia argentea</i>	isorhamnetin 3- <i>O</i> - β -D-glucoside (205) nepetin (163) quercetin 3- <i>O</i> -[(2'''-acetyl)- β -D-glucosyl]-(1 \rightarrow 6)- β -D-galactoside (199) quercetin 3- <i>O</i> - β -D-galactoside (hyperoside) (191) quercetin 3- <i>O</i> - β -D-glucosyl-(1 \rightarrow 6)- β -D-galactoside 7-(β -D-glucosyl)-4',5-dihydroxy-3',6-dimethoxyflavone (jaceoside) (146) isorhamnetin 3- <i>O</i> -dihexoside (211) quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192) quercetin 3'-methyl ether (isorhamnetin) (204) luteolin (1) quercetin (189)	Braca et al. (2008) Braca et al. (2008), Sait et al. (2015) Sait et al. (2015) Rizk (1986), Sait et al. (2015), Adjadj et al. (2015)
<i>Petrorhagia velutina</i>	isoorientin 2''- <i>O</i> -rhamnoside (20) luteolin 6- <i>C</i> -[2''- <i>O</i> - α -L-rhamnosyl-(1''' \rightarrow 2'')]- α -L-arabioside (33) luteolin 6- <i>C</i> -[2''- <i>O</i> - α -L-rhamnosyl-(1''' \rightarrow 2'')]- β -D-xyloside (34) luteolin 6- <i>C</i> - β -D-glucoside (isorientin) (18) naringenin 8- <i>C</i> - α -L-arabiosyl-7- <i>O</i> - β -D-glucoside (223) scoparin 2''- <i>O</i> -rhamnoside (140)	Pacifico et al. (2010)

Table 1 continued

Genus	Compounds	References
<i>Petrorhagia glumacea</i>	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77)	Richardson (1978)
<i>P. nanteuilli</i>	apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	
<i>P. prolifera</i>		
<i>P. velutina</i>		
<i>Petrorhagia saxifrage</i>	luteolin 6- <i>C</i> - β -D-glucoside (isorientin) (18) luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	
<i>Polycarpon tetraphyllum</i>	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77) apigenin 8- <i>C</i> - β -D-glucoside (vitexin) (49)	
<i>Psammosilene tunicoidea</i>	tectorigenin 7- <i>O</i> - β -D-glucoside (tectoridin) (228)	Liu et al. (2007)
<i>Pteranthus dichotomus</i>	apigenin (47) apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77) kaempferol (172) luteolin (1) kaempferol 3- <i>O</i> -rhamnoside-7- <i>O</i> -glucouronic acid (184) luteolin 6- <i>C</i> - β -D-glucoside (isorientin) (18) luteolin 6- <i>C</i> -rhamnosyl-(1''' \rightarrow 4'')- <i>O</i> -rhamnoside (35) myricetin 3- <i>O</i> -glucoside (215) orientin 7-methyl ether (4) quercetin 7- <i>O</i> - β -D-glucoside (203) quercetin (189)	Allaoua et al. (2016) Atta et al. (2013)
<i>Scleranthus uncinatus</i>	5,7,4'-trihydroxy-3'-methoxyflavone-8- <i>C</i> - β -D-xylosyl-2''- <i>O</i> -glucoside (138) 5,7-dihydroxy-3'-methoxy-4'-acetoxyflavone-8- <i>C</i> - β -D-xyloside-2''- <i>O</i> -glucoside (139)	Yayli et al. (2001), Yayli et al. (2002)
<i>Scleranthus annuus</i>	kaempferol (172) luteolin (1) apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77) luteolin 6- <i>C</i> - β -D-glucoside (isorientin) (18) quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192) quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190) vitexin 4'- <i>O</i> - α -L-rhamnoside (61)	Zdraveva et al. (2004)
<i>Scleranthus perennis</i>	5,7-dihydroxy-3'-methoxy-4'-acetoxyflavone-8- <i>C</i> - β -D-xylosyl-2''- <i>O</i> -glucoside (139)	Jakimiuk et al. (2020)
<i>Sagina japonica</i>	apigenin 6,8-di- <i>C</i> - β -D-glucoside (vicenin-2) (53) apigenin 6- <i>C</i> - β -D-arabinosyl-8- <i>C</i> - β -D-glucoside (vicenin-1) (62) apigenin 6- <i>C</i> - β -D-(<i>O</i> -rhamnosyl)-glucoside (95)	Zhuang (1983)
<i>Saponaria ocyroides</i>	quercetin (189)	Richardson (1978)
<i>Saponaria vaccaria</i>	quercetin (189) kaempferol (172) apigenin 6- <i>C</i> -[α -L-arabinosyl-(1''' \rightarrow 2'')- β -D-glucosyl]-7- <i>O</i> - β -D-glucoside (vaccarin) (126)	Kumar and Khanna (1994) Balsevich et al. (2011)

Table 1 continued

Genus	Compounds	References
<i>Saponaria officinalis</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77) apigenin 8-C- β -D-glucoside (vitexin) (49)	Cambie (1959)
<i>Silene alba</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77) apigenin 8-C- β -D-glucoside (vitexin) (49) isovitexin 2''-O-glucoside (meloside A) (85) vitexin 2''-O-glucoside (63)	Heinsbroek et al., (1980), Mamadalieva et al. (2014)
<i>Silene brachuica</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77) apigenin 8-C- β -D-glucoside (vitexin) (49)	Mamadalieva et al. (2014)
<i>Silene armeria</i>	luteolin 8-C- β -D-glucoside (orientin) (2) luteolin 6-C- β -D-glucoside (isorientin) (18) luteolin 8-C-(2''-O-xylosyl)- β -D-glucoside (adonivernite) (10) apigenin 6-C- β -D-glucoside (isovitexin) (77) apigenin 8-C- β -D-glucoside (vitexin) (49) luteolin 6-C-(2''-O-xylosyl)- β -D-glucoside (homoadonivernite) (26) isovitexin 4'-O- β -D-glucoside (isosaponarin) (124)	Richardson (1978), Darmograi (1977), Mamadalieva et al. (2014) Darmograi (1977), Mamadalieva et al. (2014)
<i>Silene boissieri</i>	apigenin 6,8-di-C- β -D-glucoside (vicenin-2) (53) apigenin 6-C- β -D-arabinosyl-8-C- β -D-glucoside (vicenin-1) (62) apigenin 6-C- β -D-glucoside (isovitexin) (77) apigenin 8-C- β -D-glucoside (vitexin) (49) luteolin 8-C- β -D-glucoside (orientin) (2) luteolin 6-C- β -D-glucoside (isorientin) (18)	
<i>Silene bupleuroides</i>	luteolin 8-C-(2''-O-xylosyl)- β -D-glucoside (adonivernite) (10)	
<i>S. chlorifolia</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77)	
<i>S. compacta</i>	apigenin 8-C- β -D-glucoside (vitexin) (49)	
<i>S. cretacea</i>	luteolin 6-C-(2''-O-xylosyl)- β -D-glucoside (homoadonivernite) (26)	
<i>S. cubanensis</i>		
<i>S. polaris</i>	isovitexin 4'-O- β -D-glucoside (isosaponarin) (124) luteolin 8-C- β -D-glucoside (orientin) (2) luteolin 6-C- β -D-glucoside (isorientin) (18)	
<i>Silene chlorantha</i>	apigenin 6,8-di-C- β -D-glucoside (vicenin-2) (53) apigenin 6-C- β -D-arabinosyl-8-C- β -D-glucoside (vicenin-1) (62)	
<i>S. commutate</i>		
<i>S. cyri</i>	apigenin 6-C- β -D-glucosyl-8-C- β -D-xyloside (vicenin-3) (64)	
<i>S. foliosa</i>		
<i>S. graminifolia</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77)	
<i>S. italica</i>	apigenin-8-C- β -D-glucoside (vitexin) (49)	
<i>S. jenissensis</i>	luteolin 6-C-glucoside (isorientin) (18)	
<i>S. macrostyla</i>	luteolin 8-C- β -D-glucoside (orientin) (2)	
<i>S. nutans</i>		
<i>S. wolgensis</i>		

Table 1 continued

Genus	Compounds	References
<i>Silene conoidea</i>	orientin 4'-methoxy-4''- α -L-rhamnoside (9) vitexin 4''-O-rhamnoside (65) diosmetin 8-C-(4''-O- α -L-rhamnosyl)- β -D-glucoside (153)	Ali et al. (1999), Mamadalieva et al. (2014), Ullah et al. (2019) Ahmad et al. (1998), Mamadalieva et al. (2014), Ullah et al. (2019)
<i>Silene diclinis</i>	kaempferol (172) apigenin 6-C- β -D-glucoside (isovitexin) (77) apigenin 8-C- β -D-glucoside (vitexin) (49)	Richardson (1978), Mamadalieva et al. (2014)
<i>Silene flos-cuculi</i> (syn. <i>Lychnis flos-cuculi</i>)	apigenin (47) luteolin (1) apigenin 8-C- β -D-glucoside (vitexin) (49) luteolin 8-C- β -D-glucoside (orientin) (2)	Tomczyk (2008)
<i>Silene dioica</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77)	Mamadalieva et al. (2014)
<i>Silene rubella</i>	apigenin (47) luteolin (1) luteolin 4'-methyl ether (diosmetin) (154) kaempferol (172) quercetin (189) apigenin 6-C- β -D-glucoside (isovitexin) (77) apigenin 6,8-di-C- β -D-glucoside (vicenin-2) (53) quercetin 3-O- α -L-rutinoside (rutin) (190) naringenin 7-O- α -L-hesperidoside (naringin) (221)	Hussein et al. (2019)
<i>Silene littorea</i>	kaempferol (172) luteolin 6-C- β -D-glucoside (isorientin) (18) quercetin (189)	Richardson (1978) Richardson (1978), del Valle et al. (2015)
<i>Silene macrostyla</i>	apigenin (47) luteolin (1) quercetin (189) apigenin 6-C- β -D-glucoside (isovitexin) (77) quercetin 3-O- α -L-rutinoside (rutin) (190)	del Valle et al. (2015)
<i>Silene montbretiana</i>	kaempferol 6,8-dihydroxy-3-O- α -L-rhamnoside	Kılınc et al. (2019)
<i>Silene pratensis</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77) isovitexin 7-O-galactoside-6''-O-arabinoside (108)	van Brederode et al. (1982) Niemann (1984)
<i>Silene saxatilis</i>	apigenin (46) apigenin 6,8-di-C- β -D-glucoside (vicenin-2) (53) apigenin 8-C- β -D-glucoside (vitexin) (49) luteolin 6-C- β -D-glucoside (isorientin) (18) luteolin 8-C- β -D-glucoside (orientin) (2)	Zemtsova et al. (1976), Mamadalieva et al. (2014)
<i>Silene schafta</i>	apigenin 6-C- β -D-glucosyl-8-C- α -L-arabinoside (shaftoside)	Chopin et al. (1974), Mamadalieva et al. (2014)
<i>Silene multifida</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77)	Darmograi (1977), Mamadalieva et al. (2014)
<i>S. supina</i>	apigenin 8-C- β -D-glucoside (vitexin) (49)	
<i>S. turgida</i>		

Table 1 continued

Genus	Compounds	References
<i>Silene viscariaopsis</i>	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	Richardson (1978)
<i>Silene vulgaris</i>	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	Richardson (1978), Mamadalieva et al. (2014)
<i>Silene repens</i>	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	Olennikov (2020)
	apigenin 6,8-di- <i>C</i> - β -D-glucoside (vicenin-2) (53)	
	apigenin 6- <i>C</i> - β -D-glucosyl-8- <i>C</i> - β -D-xyloside (vicenin-3) (64)	
	apigenin 6- <i>C</i> - β -D-glucoside (isovitexin) (77)	
	apigenin 6- <i>C</i> - β -glucosyl-8- <i>C</i> - α -arabinoside (schaftoside) (52)	
	luteolin 6- <i>C</i> - β -glucosyl-8- <i>C</i> -arabinoside (carlinoside, lucenin-5) (21)	
	chrysoeriol 6- <i>C</i> - β -D-glucoside (isoscoparin) (141)	
	genkwanin 6,8-di- <i>C</i> -glucoside (46)	
	genkwanin 6- <i>C</i> -glucosyl-8- <i>C</i> -arabinoside (47)	
	isomollupentin 7- <i>O</i> -glucoside-2''- <i>O</i> -arabinoside (103)	
	isoorientin 2''- <i>O</i> -arabinoside (19)	
	apigenin 6- <i>C</i> - α -arabinosyl-8- <i>C</i> - β -glucoside (isoschaftoside) (66)	
	isovitexin 2''- <i>O</i> -arabinoside (83)	
	isovitexin 2''- <i>O</i> -glucoside (meloside A) (85)	
	isovitexin 2''- <i>O</i> -xyloside (86)	
	isovitexin 7- <i>O</i> - β -D-glucoside (saponarin) (105)	
	luteolin 3'- <i>O</i> -arabinosyl-6- <i>C</i> -glucoside (lucenin-3) (44)	
	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	
	swertisin 2''- <i>O</i> -arabinoside (96)	
	swertisin 2''- <i>O</i> -glycoside (spinosin) (97)	
	isoorientin 7,3'-dimethyl ether (27)	
	silenerepin (171)	
<i>Silene schimperiana</i>	apigenin (48)	Hussein et al. (2020)
	luteolin (1)	
	luteolin 4'-methyl ether (diosmetin) (154)	
	kaempferol (172)	
	quercetin (189)	
	cyanidanon 4'-methyl ether (hesperetin) (218)	
	hesperetin 7- <i>O</i> - α -L-rutinoside (hesperidin) (219)	
	kaempferol 3- <i>O</i> - β -D-rutinoside (nicotiflorin) (176)	
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
<i>Spergularia diandra</i>	tricin (159)	El-Dien et al. (2013)

Table 1 continued

Genus	Compounds	References
<i>Spergularia marina</i>	tricin (159)	
	apigenin 6- <i>C</i> - β -D-(2 ^{'''} - <i>O</i> -feruloyl)glucosyl-8- <i>C</i> - β -D-glucoside (70)	Cho et al. (2016)
	apigenin 6- <i>C</i> - β -D-glucosyl-8- <i>C</i> - β -D-(2 ^{'''} - <i>O</i> -feruloyl)glucoside (67)	
	luteolin 6- <i>C</i> - β -D-(2 ^{''} - <i>O</i> -feruloyl)glucosyl-8- <i>C</i> - β -D-glucoside (8)	
<i>Spergularia rubra</i>	luteolin 6- <i>C</i> - β -D-glucosyl-8- <i>C</i> - β -D-(2- <i>O</i> ^{'''} -feruloyl)glucoside (11)	
	apigenin 6,8-di- <i>C</i> - β -D-glucoside (vicenin-2) (53)	Zoll et al. (1974), Bouillant et al. (1979), Ferreres et al. (2011)
	apigenin 6- <i>C</i> -arabinoside (isomollupentin) (81)	Bouillant et al. (1979)
	apigenin 6- <i>C</i> - β -glucosyl-8- <i>C</i> - α -arabinoside (schaftoside) (52)	Bouillant et al. (1979), Ferreres et al. (2011)
<i>Spergularia salina</i>	chrysoeriol 6,8-di- <i>C</i> -glucoside (stellarin-2) (130)	Zoll and Nouvel (1974)
	apigenin 6,8-di- <i>C</i> -(6 ^{'''} -malonyl, feruloyl)glucoside (68)	
	apigenin 6,8-di- <i>C</i> -(6 ^{'''} -malonyl, sinapoyl)glucoside (69)	
	apigenin 6- <i>C</i> -(2 ^{''} -feruloyl)glucosyl-8- <i>C</i> -glucoside (70)	
	apigenin 6- <i>C</i> -(4 ^{''} -malonyl)glucosyl-8- <i>C</i> -glucoside (71)	
	apigenin 6- <i>C</i> -glucosyl-8- <i>C</i> -(2 ^{'''} -feruloyl)glucoside (67)	
	apigenin 6- <i>C</i> -glucosyl-8- <i>C</i> -(2 ^{'''} -sinapoyl)glucoside (72)	
	chrysoeriol 6,8-di- <i>C</i> -(6 ^{''} -malonyl, sinapoyl)glucoside (135)	
	chrysoeriol 6,8-di- <i>C</i> -(6 ^{''} -malonyl, feruloyl)glucoside (136)	
	chrysoeriol 6,8-di- <i>C</i> -glucoside (stellarin-2) (130)	
	chrysoeriol 6- <i>C</i> -(4 ^{''} -malonyl)glucosyl-8- <i>C</i> -glucoside (137)	
	chrysoeriol 6- <i>C</i> -arabinosyl-8- <i>C</i> -glucoside (131)	
	chrysoeriol 6- <i>C</i> -glucosyl-8- <i>C</i> -(2 ^{'''} -feruloyl)glucoside (133)	
	chrysoeriol 6- <i>C</i> -glucosyl-8- <i>C</i> -(2 ^{'''} -sinapoyl)glucoside (134)	
	chrysoeriol 6- <i>C</i> -glucosyl-8- <i>C</i> -arabinoside (132)	
	chrysoeriol 7- <i>O</i> -glucosyl-6- <i>C</i> -(2 ^{''} -malonyl)-arabinosyl-8- <i>C</i> -arabinoside (144)	
	chrysoeriol 7- <i>O</i> -glucosyl-6- <i>C</i> -arabinosyl-8- <i>C</i> -(6 ^{'''} -malonyl)arabinoside (145)	
	luteolin 6,8-di- <i>C</i> -(2 ^{''} -malonyl, feruloyl)glucoside (12)	
	luteolin 6,8-di- <i>C</i> -glucoside (lucenin-2) (5)	
	luteolin 6- <i>C</i> -(2 ^{''} -feruloyl)glucosyl-8- <i>C</i> -glucoside (8)	
luteolin 6- <i>C</i> -(6 ^{''} -acetyl)glucosyl-8- <i>C</i> -glucoside (6)		
luteolin 6- <i>C</i> -(6 ^{''} -malonyl)glucosyl-8- <i>C</i> -glucoside (7)		
luteolin 6- <i>C</i> -glucosyl-8- <i>C</i> -(4 ^{''} -malonyl)glucoside (13)		
luteolin 6- <i>C</i> -glucosyl-8- <i>C</i> -(2 ^{'''} -dihydroferuloyl)glucoside (14)		
luteolin 6- <i>C</i> -glucosyl-8- <i>C</i> -(2 ^{'''} -feruloyl)glucoside (15)		
luteolin 6- <i>C</i> -glucosyl-8- <i>C</i> -(2 ^{'''} - <i>p</i> -coumaroyl)glucoside (16)		
luteolin 6- <i>C</i> -glucosyl-8- <i>C</i> -(2 ^{'''} -sinapoyl)glucoside (17)		
luteolin 6- <i>C</i> -glucosyl-8- <i>C</i> -arabinoside (21)		

Table 1 continued

Genus	Compounds	References
	luteolin 7- <i>O</i> -glucosyl-6- <i>C</i> -glucosyl-8- <i>C</i> -(2''-feruloyl)glucoside (41)	Ferreres et al. (2011), Vinholes et al. (2011)
	luteolin 7- <i>O</i> -glucosyl-6,8- <i>C</i> -diglucoside (42)	
	luteolin 7- <i>O</i> -glucosyl-6- <i>C</i> -glucosyl-8- <i>C</i> -(2''sinapoyl)glucoside (43)	
<i>Stellaria dichotoma</i>	apigenin (48)	Mikšátková et al. (2014)
	formononetin (224)	
	genistein (229)	
	glycitein (227)	
	isoformononetin (226)	
	kaempferol (172)	
	luteolin (1)	
	naringenin (220)	
	quercetin (189)	
	tectorigenin 7- <i>O</i> - β -D-glucoside (tectoridin) (228)	
	apigenin 7- <i>O</i> - β -D-glucoside (cosmosiin) (111)	
	formononetin 7- <i>O</i> - β -D-glucoside (ononin) (225)	
	genistein 7- <i>O</i> - β -D-glucoside (genistin) (230)	
	luteolin 7- <i>O</i> - β -D-glucoside (cynaroside) (37)	
	naringenin 7- <i>O</i> - β -D-glucoside (prunin) (222)	
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
	apigenin 6,8-di- <i>C</i> - β -D-glucoside (vicenin-2) (53)	Yasukawa et al. (1981), Sharma and Arora (2012)
	isoscutellarein 6- <i>C</i> -galactoside (78)	
<i>Stellaria graminea</i>	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	Richardson (1978)
	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	
<i>Stellaria holostea</i>	apigenin (48)	Mikšátková et al. (2014)
	genistein (229)	
	kaempferol (172)	
	luteolin (1)	
	naringenin (220)	
	quercetin (189)	
	apigenin 7- <i>O</i> - β -D-glucoside (cosmosiin) (111)	
	daidzein 7- <i>O</i> - β -D-glucoside (daidzin) (231)	
	formononetin 7- <i>O</i> - β -D-glucoside (ononin) (225)	
	genistein 7- <i>O</i> - β -D-glucoside (genistin) (230)	
	luteolin 7- <i>O</i> - β -D-glucoside (cynaroside) (37)	
	naringenin 7- <i>O</i> - β -D-glucoside (prunin) (222)	
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
	apigenin 6- <i>C</i> - β -glucosyl-8- <i>C</i> - α -arabinoside (schaftoside) (52)	Ancheeva et al. (2015)
	diosmetin 6- <i>C</i> - β -glucoside (158)	
	3,5,7-trihydroxy-3',5'-dimethoxyflavone (216)	

Table 1 continued

Genus	Compounds	References
	apigenin 6,8-di- <i>C</i> - β -D-glucoside (vicenin-2) (53)	Sharma and Arora (2012)
	chrysoeriol 6,8-di- <i>C</i> -glucoside (stellarin-2) (130)	
	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	Richardson (1978), Boguslavskaya et al. (1985a, b), Ancheeva et al. (2015)
	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	
<i>Stellaria media</i>	apigenin (48)	Kitanov (1992), Sharma and Arora (2012), Mikšátková et al. (2014), Rogowska et al. (2017); Melnyk et al. (2018)
	luteolin (1)	Mikšátková et al. (2014), Melnyk et al. (2018)
	quercetin 3- <i>O</i> - α -L-rutinoside (rutin) (190)	
	genistein (229)	Kitanov (1992), Sharma and Arora (2012)
	apigenin 6,8-di- <i>C</i> - β -D-glucoside (vicenin-2) (53)	
	apigenin 6- <i>C</i> - α -L-arabinosyl-8- <i>C</i> - β -D-galactoside (57)	Dong et al. (2007)
	apigenin 6- <i>C</i> - β -D-galactosyl-8- <i>C</i> - α -L-arabinoside (74)	
	apigenin 6- <i>C</i> - β -D-galactosyl-8- <i>C</i> - β -L-arabinoside (75)	
	apigenin 6- <i>C</i> - β -D-glucosyl-8- <i>C</i> - β -D-galactoside (50)	
	apigenin 6,8-di- <i>C</i> - α -L-arabinoside (73)	
	luteolin 8- <i>C</i> - β -D-glucoside (orientin) (2)	Richardson (1978)
	luteolin 6- <i>C</i> - β -D-glucoside (isoorientin) (18)	
	quercetin 3- <i>O</i> - β -D-glucoside (isoquercitrin) (192)	Melnyk et al. (2018)
<i>Stellaria media</i>	formononetin (224)	Dong et al. (2007), Mikšátková et al. (2014)
	glycitein (227)	
<i>S. nemorum</i>	kaempferol (172)	
	naringenin (220)	
	quercetin (189)	
	apigenin 7- <i>O</i> - β -D-glucoside (cosmosiin) (111)	
	formononetin 7- <i>O</i> - β -D-glucoside (ononin) (225)	
	genistein 7- <i>O</i> - β -D-glucoside (genistin) (230)	
	luteolin 7- <i>O</i> - β -D-glucoside (cynaroside) (37)	
	naringenin 7- <i>O</i> - β -D-glucoside (prunin) (222)	

Table 1 continued

Genus	Compounds	References
<i>Stellaria nemorum</i>	apigenin 6-C-[(α -arabinosyl)-(1 \rightarrow 2)- <i>O</i> - β -xyloside]	Mikšátková et al. (2014), Ancheeva et al. (2015)
	apigenin 6-C-[(α -arabinosyl)-(1 \rightarrow 2)- <i>O</i> - β -glucoside]	
	apigenin 6-C- β -galactosyl-8-C- β -glucoside	
	apigenin 6-C- β -glucosyl-8-C- α -arabinoside (schaftoside) (52)	
	apigenin 6-C- β -glucosyl-8-C- β -xyloside (55)	
<i>Telephium imperati</i>	apigenin 6-C- β -D-glucoside (isovitexin) (77)	Richardson (1978)
	apigenin 8-C- β -D-glucoside (vitexin) (49)	
<i>Vaccaria segetalis</i>	apigenin (48)	Baeva et al. (1975)
	apigenin 6-C- β -D-glucoside (isovitexin) (77)	
	apigenin 6-C-arabinosyl- <i>O</i> -glucoside (116)	
	apigenin 6-C-glucosyl- <i>O</i> -glucoside	
	apigenin 6-C-[α -L-arabinosyl-(1''' \rightarrow 2'')- β -D-glucosyl]-7- <i>O</i> - β -D-glucoside (vaccarin) (126)	
	isovitexin 4'- <i>O</i> - β -D-glucoside (isosaponarin) (124)	
	apigenin 6-C-[α -L-arabinopyranosyl-(1''' \rightarrow 2'')- β -D-glucopyranosyl]-7- <i>O</i> - β -D-glucoside (vaccarin) (126)	
	apigenin 6-C-[α -L-arabinosyl-(1''' \rightarrow 2'')- β -D-glucosyl]-7- <i>O</i> -(6'''- <i>O</i> -dihydroferuloyl)- β -D-glucoside (vaccarin-E) (118)	
	apigenin 6-C- β -D-glucosyl-7- <i>O</i> -(6'''- <i>O</i> -dihydroferuloyl)- β -D-glucoside (vaccarin-F) (119)	
	isovitexin 7- <i>O</i> - β -D-glucoside (saponarin) (105)	
	isovitexin 2''- <i>O</i> -arabinoside (83)	
	isovitexin 4'- <i>O</i> - β -D-glucoside (isosaponarin) (124)	
	isovitexin-2''- <i>O</i> - α -L-arabinosyl-4'- <i>O</i> -(6'''- <i>O</i> -dihydroferuloyl)- β -D-glucoside (vaccarin-H) (127)	
<i>Vaccaria pyramidata</i>	vitexin 7- <i>O</i> - β -D-glucoside (117)	Said et al. (2019)
	vitexin 2''- <i>O</i> - α -L-rhamnoside (76)	

Isoflavones


Phytoestrogens are non-steroidal polyphenolic compounds occurring in plants and can be chemically divided into two main groups: flavonoids (isoflavones) and non-flavonoids (lignans). The structure of isoflavone aglycone consists of a 3-phenylchroman ring that is substituted with hydroxyl groups in the positions C4' and C7 (Bustamante-Rangel et al. 2018; Křížová et al. 2019) (Fig. 14).

Because flavonoids are widely distributed in the plant kingdom and their presence in Caryophyllaceae plants has not been published until now, the authors of

the article summarized the phytochemistry of 26 flavonoid-producing genera and relevant species. The flavonoid compounds occurring in Caryophyllaceae, the corresponding species and the literature references are summarized in Table 1.

Conclusions

The Caryophyllaceae family contains a large number of genera and species that are widely distributed over different climate zones. It is evident that the plants from this family produce a wide range of




	R	R ₁	R ₂	R ₃	R ₄	R ₅
1	OH	H	OH	H	OH	OH
2	OH	H	OH	<i>β</i> -D- <i>glc</i>	OH	OH
3	OH	<i>ara</i>	OH	<i>glc</i>	OH	OH
4	OH	H	OCH ₃	<i>β</i> -D- <i>glc</i>	OH	OH
5	OH	<i>glc</i>	OH	<i>glc</i>	OH	OH
6	OH	<i>glc</i> -(6''-O-acetyl)	OH	<i>glc</i>	OH	OH
7	OH	<i>glc</i> -(6''-O-malonyl)	OH	<i>glc</i>	OH	OH
8	OH	<i>glc</i> -(2''-O-feruloyl)	OH	<i>glc</i>	OH	OH
9	OH	H	OH	<i>α</i> -L- <i>glc</i> -(4''-O- <i>rha</i>)	OH	OCH ₃
10	OH	H	OH	<i>β</i> -D- <i>glc</i> -(2''-O- <i>xyl</i>)	OH	OH
11	OH	<i>β</i> -D- <i>glc</i>	OH	<i>β</i> -D- <i>glc</i> -(2''-O-feruloyl)	OH	OH
12	OH	<i>glc</i> -(2''-O-malonyl, feruloyl)	OH	<i>glc</i> -(2''-O-malonyl, feruloyl)	OH	OH
13	OH	<i>glc</i>	OH	<i>glc</i> -(4''-O-malonyl)	OH	OH
14	OH	<i>glc</i>	OH	<i>glc</i> -(2''-O-dihydroferuloyl)	OH	OH
15	OH	<i>glc</i>	OH	<i>glc</i> -(2''-O-feruloyl)	OH	OH
16	OH	<i>glc</i>	OH	<i>glc</i> -(2''-O- <i>p</i> -coumaroyl)	OH	OH
17	OH	<i>glc</i>	OH	<i>glc</i> -(2''-O-sinapoyl)	OH	OH
18	OH	<i>β</i> -D- <i>glc</i>	OH	H	OH	OH
19	OH	<i>β</i> -D- <i>glc</i> -(2''-O- <i>ara</i>)	OH	H	OH	OH
20	OH	<i>β</i> -D- <i>glc</i> -(2''-O- <i>rha</i>)	OH	H	OH	OH
21	OH	<i>β</i> -D- <i>glc</i>	OH	<i>ara</i>	OH	OH
22	OH	<i>β</i> -D- <i>glc</i>	O- <i>gal</i>	H	OH	OH
23	OH	<i>β</i> -D- <i>glc</i>	O- <i>rha</i> -O- <i>gal</i>	H	OH	OH
24	OH	<i>β</i> -D- <i>glc</i>	O- <i>rut</i>	H	OH	OH
25	OH	<i>β</i> -D- <i>glc</i>	O- <i>α</i> -L- <i>ara</i>	H	OH	OH
26	OH	<i>β</i> -D- <i>glc</i> -(2''-O- <i>xyl</i>)	OH	H	OH	OH
27	OH	<i>β</i> -D- <i>glc</i>	OCH ₃	H	OCH ₃	OH
28	OH	<i>glc</i> -(2''-O-feruloyl)	OH	H	OH	O- <i>glc</i>
29	OH	<i>hex</i> -(2''-O- <i>pen</i>)	OH	H	OH	OH
30	OH	<i>hex</i> -(2''-O- <i>pen</i>)	OH	H	OH/OCH ₃	OH/OCH ₃
31	OH	<i>hex</i> -(2''-O- <i>hex</i>)	OH	H	OH	OH
32	OH	<i>hex</i>	OH	<i>hex</i>	OH	OH
33	OH	<i>ara</i> -[2''-O- <i>rha</i> -(1'''→2'')]]	OH	H	OH	OH
34	OH	<i>xyl</i> -[2''-O- <i>rha</i> -(1'''→2'')]]	OH	H	OH	OH
35	OH	<i>rha</i> -[4''-O- <i>rha</i> (1'''→4'')]]	OH	OH	OH	OH
36	O- <i>β</i> -D- <i>glc</i>	H	OH	H	OH	OH
37	OH	H	O- <i>β</i> -D- <i>glc</i>	H	OH	OH
38	OH	H	<i>glc</i> - <i>glc</i>	H	OH	OH
39	OH	<i>hex</i>	O- <i>hex</i> - <i>malyl</i>	H	OH	OH
40	OH	H	OH	H	OH	O- <i>β</i> -D- <i>glc</i>
41	OH	<i>glc</i>	O- <i>glc</i>	<i>glc</i> -(2'''-O-feruloyl)	OH	OH
42	OH	<i>glc</i>	O- <i>glc</i>	<i>glc</i>	OH	OH
43	OH	<i>glc</i>	O- <i>glc</i>	<i>glc</i> -(2'''-O-sinapoyl)	OH	OH
44	OH	<i>β</i> -D- <i>glc</i>	OH	H	O- <i>ara</i>	OH
45	OH	<i>β</i> -D- <i>glc</i> -(2''-O- <i>α</i> -L- <i>rha</i>)	OH	H	OH	OH
46	OH	<i>glc</i>	OCH ₃	<i>glc</i>	OH	OH
47	OH	<i>glc</i>	OCH ₃	<i>ara</i>	OH	OH

Fig. 10 The chemical structures of the luteolin and its derivatives identified in species of Caryophyllaceae family

pharmaceutically promising, interesting, and valuable flavonoids and other secondary metabolites. Phytochemical data of flavonoids in plants of this family have not been published until now. Despite the dominant proportion of triterpene saponins among all phytoconstituents, polyphenols, including

flavonoid compounds, remain a large group of compounds with health-related activity, such as antioxidant, anti-inflammatory, antimicrobial, organoprotective, and even anticancer effects (van Wyk and Wink 2017; Imran et al. 2019a; Ganeshpurkar and Saluja 2019). Our approach involved screening

Fig. 11 The chemical structures of the apigenin and its derivatives identified in species of Caryophyllaceae family



	R	R ₁	R ₂	R ₃	R ₄
48	OH	H	OH	H	OH
49	OH	H	OH	β -D-glc	OH
50	OH	β -D-glc	OH	β -D-gal	OH
51	OH	xyl	OH	ara	OH
52	OH	β -D-glc	OH	α -L-ara	OH
53	OH	β -D-glc	OH	β -D-glc	OH
54	OH	gal	OH	ara	OH
55	OH	β -D-glc	OH	β -D-glc	OH
56	OH	hex	OH	pen	OH
57	OH	α -L-ara	OH	β -D-gal	OH
58	OH	glc	OH	rha	OH
59	OH	rha	OH	glc	OH
60	OH	H	OH	α -D-glc	OH
61	OH	H	OH	β -D-glc	O- α -L-rha
62	OH	β -D-ara	OH	β -D-glc	OH
63	OH	H	OH	β -D-glc (2'-O-glc)	OH
64	OH	β -D-glc	OH	β -D-xyl	OH
65	OH	H	OH	β -D-glc-(4'-O-rha)	OH
66	OH	α -ara	OH	β -D-glc	OH
67	OH	glc	OH	glc-(2'-O-feruloyl)	OH
68	OH	glc-(6''-O-malonyl, feruloyl)	OH	glc-(6''-O-malonyl, feruloyl)	OH
69	OH	glc-(6''-O-malonyl, sinapoyl)	OH	glc-(6''-O-malonyl, sinapoyl)	OH
70	OH	glc-(2''-O-feruloyl)	OH	glc	OH
71	OH	glc-(2''-O-malonyl)	OH	glc	OH
72	OH	glc	OH	glc-(2'-O-sinapoyl)	OH
73	OH	α -L-ara	OH	α -L-ara	OH
74	OH	β -D-gal	OH	α -L-ara	OH
75	OH	β -D-gal	OH	β -L-ara	OH
76	OH	H	OH	β -D-glc-(2'-O- α -L-rha)	OH
77	OH	β -D-glc	OH	H	OH
78	OH	gal	OH	ara	OH
79	OH	anti- α -D-glc	OH	H	OH
80	OH	syn- α -D-glc	OH	H	OH
81	OH	ara	OH	H	OH
82	OH	xyl	OH	H	OH
83	OH	β -D-glc-(2'-O-ara)	OH	H	OH
84	OH	β -D-glc-(2'-O-feruloyl)	OH	H	OH
85	OH	β -D-glc-(2'-O-glc)	OH	H	OH
86	OH	β -D-glc-(2'-O-xyl)	OH	H	OH
87	OH	β -D-glc-(2'-O- α -L-rha)	OCH ₃	H	OH
88	OH	hex-(2''-O-acetyl-pent)	OH	H	OH
89	OH	hex-(2''-O-diacetyl-pent)	OH	H	OH
90	OH	hex-(2'-O-pent)	OH	H	OH
91	OH	hex-(2'-O-hex)	OH	H	OH
92	COCH ₃	β -D-glc-(2'-O-rha)	OH	H	OH
93	OH	ara-(O-diglc)	OH	H	OH
94	OH	triglc	OH	H	OH
95	OH	β -D-glc-(O-rha)	OH	H	OH
96	OH	β -D-glc (2'-O-ara)	OCH ₃	H	OH
97	OH	β -D-glc -2'-O-glc)	OCH ₃	H	OH
98	OH	ara(2'-O-glc)	OH	H	OH
99	OH	β -D-glc-(2'-O-rha)	OH	H	OH
100	OH	ara	OH	xyl	OH
101	OH	xyl	C-glu	H	OH
102	OH	ara(2'-O-glc)	C-glu	H	OH
103	OH	ara(2'-O-ara)	C-glu	H	OH
104	OH	ara(2''-O-xyl)	C-glu	H	OH
105	OH	β -D-glc	O- β -D-glc	H	OH
106	OH	β -D-glc-[2'-O-rha-(6''-O-7,8-dihydroferuloyl)]	O- β -D-glc	H	OH
107	OH	β -D-glc	O-gal-(6''-O-ara)	H	OH
108	OH	β -D-glc	C-gal	H	OH
109	OH	β -D-glc	O-rut	H	OH
110	OH	hex	O-hex-malyl	H	OH
111	OH	H	O- β -D-glc	H	OH
112	OH	β -D-glc	O-glc-(6''-O-malyl)	H	OH
113	OH	β -D-glc-[xyl-(1'' \rightarrow 2'')-O-gal]	O- β -D-glc	H	OH
114	OH	H	O- β -D-glc-[(1'' \rightarrow 2'')-O-glc]	H	OH
115	OH	H	β -D-O-glc-[(1'' \rightarrow 2'')-O-rha]	H	OH
116	OH	ara-(O-glc)	O-glc	H	OH
117	OH	H	β -D-O-glc	β -D-glc	OH
118	OH	β -D-glc-[(1'' \rightarrow 2'')- α -L-ara]	O- β -D-glc-(6''-O-dihydroferuloyl)	H	OH
119	OH	β -D-glc	O- β -D-glc-(6''-O-dihydroferuloyl)	H	OH
120	OH	ara	OH	H	O-glu
121	OH	β -D-glc-(2'-O-feruloyl)	OH	H	O-glu
122	OH	β -D-glc-(2'-O-ara)	OH	H	O-glc
123	OH	β -D-glc-(2'-O-glc)	OH	H	O-glc
124	OH	β -D-glc	OH	H	O- β -D-glu
125	OH	H	OH	H	O-glu
126	OH	β -D-glc-[2''-O- α -L-ara-(1''' \rightarrow 2''')]]	OH	H	O-glu
127	OH	β -D-glc-[(1'' \rightarrow 2'')- α -L-ara]	OH	H	O- β -D-glc-(6''-O-dihydroferuloyl)




Fig. 12 The chemical structures of the chrysoeriol (a), acacetin (b), diosmetin (c), tricrin (d), and their derivatives identified in species of Caryophyllaceae family

flavonoid-containing species, including those containing aglycones and their glycoside derivatives, which could be identified in 26 genera and more than 120 species within the Caryophyllaceae.

To the best of our knowledge, apigenin is the most common aglycone in this family and can be found in 28 different species, such as *Vaccaria segetalis* (Baeva et al. 1975), *Stellaria media* (Melnik et al. 2018), *Silene saxatilis* (Zemtsova et al. 1975), *Pteranthus dichotomus* (Allaoua et al. 2016), *Silene* (*Lychnis*)

flos-cuculi (Tomczyk 2008), *Herniaria glabra* (El Mabruki et al. 2014) and others. Furthermore, the C-bonded apigenin glucoside isovitexin has been isolated from more than 70 plants, making it the predominant flavonoid within this family. On the basis of the data collected in Table 1, it was concluded that the highly glycosylated C- and O-flavonoids (apigenin, luteolin, chrysoeriol, kaempferol, quercetin, formononetin, genistein, myricetin, tectorigenin) with either one, two or three sugar moieties, as presented in

		R	R₁	R₂	R₃	
a		172	OH	H	OH	OH
		173	<i>O</i> -β-D- <i>glc</i> -(<i>O</i> -β-D- <i>glc</i>)	H	OH	OH
		174	<i>O</i> - <i>glc</i> -[2''- <i>O</i> -(6- <i>malyl</i>)- <i>glc</i> -6''- <i>rha</i>]	H	OH	OH
		175	<i>O</i> - <i>glc</i> -[2''- <i>O</i> - <i>glc</i> -6''- <i>rha</i>]	H	OH	OH
		176	<i>O</i> -β-D- <i>rut</i>	H	OH	OH
		177	<i>O</i> -β-D- <i>glc</i> -[<i>O</i> -β-D- <i>glc</i> (1''→2'')- <i>O</i> -α-L- <i>rha</i> (1''→2'')- <i>O</i> -β-D- <i>glc</i> (1''→6'')]	H	OH	OH
		178	<i>O</i> -β-D- <i>glc</i> -[<i>glc</i> (1''→2'')- <i>O</i> -α-L- <i>rha</i> (1''→6'')]	H	OH	OH
		179	<i>O</i> -β-D- <i>glc</i> -(<i>O</i> - <i>rha</i>)	H	OH	OH
		180	<i>O</i> - <i>glc</i>	H	OH	OH
		181	<i>O</i> - <i>rha</i>	H	<i>O</i> - <i>rha</i>	OH
		182	<i>O</i> - <i>glc</i> -(2''- <i>O</i> - <i>p</i> - <i>coumaroyl</i>)	H	OH	OH
		183	OH	H	<i>O</i> - <i>rha</i> - <i>O</i> - <i>glc</i>	OH
		184	<i>O</i> - <i>rha</i>	H	<i>O</i> -glucuronic acid	OH
		185	<i>O</i> -β-D- <i>glc</i> -(1''→2'')-β-D- <i>glc</i>	OH	OH	OH
		186	OH	H	OH	OCH ₃
		187	<i>O</i> -[2 ^G -β-D- <i>glc</i>]-β- <i>rut</i>	H	OH	OCH ₃
		188	<i>O</i> -β-D- <i>glc</i> -[<i>glc</i> (1''→2'')- <i>O</i> -α-L- <i>rha</i> (1''→6'')]	H	OH	OCH ₃
		b		189	OH	OH
190	<i>O</i> -α-L- <i>rut</i>			OH	OH	
191	<i>O</i> -β-D- <i>gal</i>			OH	OH	
192	<i>O</i> -β-D- <i>glc</i>			OH	OH	
193	<i>O</i> - <i>glc</i>			OCH ₃	OH	
194	<i>O</i> -β-D- <i>glc</i> -(6''- <i>O</i> - <i>ara</i>)			OH	OH	
195	<i>O</i> -β-D- <i>glu</i> -(2''- <i>O</i> -α-L- <i>rha</i>)			OH	OH	
196	<i>O</i> -β-D- <i>glc</i> - <i>O</i> - <i>gal</i>			OH	OH	
197	<i>O</i> -β-D- <i>glc</i> - <i>O</i> -β-D- <i>glc</i>			OH	OH	
198	<i>O</i> -β-D- <i>glc</i> -[(1''→6'')-2''- <i>O</i> - <i>acetyl</i> - <i>O</i> -β-D- <i>xyf</i>]			OH	OH	
199	<i>O</i> -β-D- <i>gal</i> -[(1''→6'')-2''- <i>O</i> - <i>acetyl</i> - <i>O</i> -β-D- <i>glc</i>]			OH	OH	
200	<i>O</i> -β-D- <i>glc</i> -[(1''→2'')-D- <i>apio</i> - <i>O</i> -β-D- <i>furanosyl</i> - <i>O</i> -α-L- <i>rha</i> (1''→6'' ^v)]			OH	OH	
201	<i>O</i> - <i>glc</i>			<i>O</i> - <i>rha</i>	OH	
202	OH			OCH ₃	OH	
203	OH			<i>O</i> -β-D- <i>glc</i>	OH	
204	OH			OH	OCH ₃	
205	<i>O</i> -β-D- <i>glc</i>			OH	OCH ₃	
206	<i>O</i> - <i>rut</i>			OH	OCH ₃	
207	<i>O</i> - <i>rab</i>	OH	OCH ₃			
208	<i>O</i> -(3''- <i>O</i> - <i>feruloyl</i>)	OH	OCH ₃			
209	<i>O</i> - <i>glc</i>	OCH ₃	OCH ₃			
210	<i>O</i> - <i>rut</i>	OCH ₃	OCH ₃			
211	<i>O</i> - <i>hex</i> - <i>O</i> - <i>hex</i>	OH	OCH ₃			
212	<i>O</i> -β-D- <i>glc</i> -[(1''→6'')-2''- <i>O</i> - <i>acetyl</i> - <i>O</i> -β-D- <i>xyf</i> -(1''→6'' ^v)-β-D- <i>apiofuranosyl</i>]	OH	OCH ₃			
213	<i>O</i> -β-D- <i>glc</i> -[(1''→6'')-2''- <i>O</i> - <i>acetyl</i> - <i>O</i> -β-D- <i>xyf</i>]	OH	OCH ₃			
c		214	OH	OCH ₃	OH	
		215	<i>O</i> - <i>glc</i>	OH	OH	
		216	OH	OCH ₃	OCH ₃	

Fig. 13 The chemical structures of the flavonols identified in Caryophyllaceae. Kaempferol and its derivatives (a), quercetin and its derivatives (b), myricetin and its derivatives (c) identified in species of Caryophyllaceae family

this review, are commonly found in the Caryophyllaceae family. The genera *Silene* Mill., *Dianthus* L., *Stellaria* L., *Herniaria* L., *Spergularia* Presl., *Gypsophila* L. and *Cerastium* L. appear to contain high abundances of flavonoid compounds.

In summary, the structural diversity of flavonoids established in the Caryophyllaceae family makes them an interesting object of phytochemical and pharmacological investigations.



	217	R	R₁	R₂	R₃
	218	OH	H	OCH ₃	OH
	219	OH	H	OH	OCH ₃
	220	O-β-D-glc-(α-L-rha)	H	OH	OCH ₃
	221	OH	H	H	H
	222	O-β-D-glc-(2''-O-α-L-rha)	H	H	H
	223	O-glc	H	H	H
	223	O-β-D-glc	α-L-ara	H	H
	224	OH	H	H	OCH ₃
	225	O-glc	H	H	OCH ₃
	226	OCH ₃	H	H	OH
	227	OH	OCH ₃	H	OH
	228	O-glc	OCH ₃	OH	OH
	229	OH	H	OH	OH
	230	O-glc	H	OH	OH
	231	O-glc	H	H	OH

Fig. 14 The chemical structures of the flavonones (a) and isoflavones (b) identified in species of Caryophyllaceae family

Authors' contributions Conceptualization and Methodology, K.J., M.T.; Formal Analysis, K.J.; Investigation, K.J.; Writing – Original Draft Preparation, K.J.; Writing – Review and Editing, M.T., M.W.; Supervision, M.T.; Project Administration, M.T.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Adjadj M, Baghiani A, Boumerfeg S, Noureddine C, Khennouf S, Arrar L, Mubarak MS (2015) Protective effect of *Paronychia argentea* L. on acetic acid induced ulcerative colitis in mice by regulating antioxidant parameters and inflammatory markers. *Wulfenia J* 22:148–172
- Ahmad V, Ali Z, Ali M, Zahid M (1998) Chemical constituents of *Silene conoidea*. *Fitoterapia* 69:406–408
- Aldhebiani AY, Mufarah N (2017) Phytochemical screening of some wild plants from Wadi Yalmlam, Saudi Arabia. *IOSR J Pharm Biol Sci* 12:25–27
- Ali Z, Ahmad VU, Ali MS, Iqbal F, Zahid M, Alam N (1999) Two new C-glycosylflavones from *Silene conoidea*. *Nat Prod Lett* 13:121–129
- Allaoua Z, Benkhaled M, Dibi A, Long C, Aberkane MC, Bouzidi S, Haba H (2016) Chemical composition, antioxidant and antibacterial properties of *Pteranthus dichotomus* from Algerian Sahara. *Nat Prod Res* 30:700–704
- Al-Snafi AE (2017) Chemical contents and medical importance of *Dianthus caryophyllus*—a review. *ISOR J Pharm* 7:61–71
- Altay A (2018) HPLC Analysis of phenolic compounds from *Gypsophila aucheri* Boiss. and investigation of antioxidant and cytotoxic activity of *Gypsophila aucheri* Boiss. extracts. *J Sci Technol* 11:168–181
- Altay A, Degirmenci S, Korkmaz M, Cankaya M, Koksall E (2018) *In vitro* evaluation of antioxidant and anti-proliferative activities of *Gypsophila sphaerocephala* (Caryophyllaceae) extracts together with their phenolic profiles. *J Food Meas Charact* 12:2936–2945
- Altay A, Tohma H, Durmaz L, Taslimi P, Korkmaz M, Gulcin I, Koksall E (2019) Preliminary phytochemical analysis and evaluation of *in vitro* antioxidant, antiproliferative, antidiabetic, and anticholinergic effects of endemic *Gypsophila* taxa from Turkey. *J Food Biochem* 43:1–11
- Amosova EN, Zueva EP, Lopatina KA, Safonova EA, Razina TG, Rybalkina OY, Zibareva LN (2019) Influence of *Lychnis chalconica* L. flavonoids on transplanted tumor development and cytostatic therapy effectiveness in mice. *Pharm Chem J* 53:454–457
- Ancheeva E, Daletos G, Muharini R, Lin WH, Teslov L, Proksch P (2015) Flavonoids from *Stellaria nemorum* and *Stellaria holostea*. *Nat Prod Commun* 10:437–440
- Atkinson C, Compston JE, Day NE, Dowsett M, Bingham SA (2004) The effects of phytoestrogen isoflavones on bone density in women: a double-blind, randomized, placebo-controlled trial. *Am J Clin Nutr* 79:326–333
- Atta EM, Nassar AA, Hasan NM, Hasa AR (2013) New flavonoid glycoside and pharmacological activities of *Pteranthus dichotomus* forssk. *Rec Nat Prod* 7:69–79
- Avunduk S, Lacaille-Dubois MA, Miyamoto T, Bedir E, Şenol SG, Çalişkan ÖA (2007) Chionaeosides A-D, triterpene

- saponins from *Paronychia chionaea*. J Nat Prod 70:1830–1833
- Azadi B, Sohrabi Y (2014) Chemical composition of *Silene morganae* Freyn volatile oil. Nat Prod Res 29:791–794
- Baeva RT, Karryev MO, Litvinenko VI, Abubacirov NK (1975) Glycosides of *Vaccaria segetalis* V. Vaccarin. Chem Nat Compd 10:182–186
- Bahar A, Mubashir HM, Shamshir K (2008) *Lychnis coronaria* Linn. A review. Nat Prod Indian J 4:22–25
- Bakroglu A, Kökten K, Kavurmaci Z (2014) Tannin, protein contents and fatty acid compositions of *Silene compacta* Fische seeds from Bingöl, Turkey. Turk J Agric Nat Sci 1:441–444
- Balamurugan K, Sakthidevi G, Mohan VR (2013) Antiulcer activity of *Polycarpaea corymbosa* (L.) Lam. whole plant extracts (Caryophyllaceae). Int J Biol Med Res 4:3379–3382
- Balsevich JJ, Ramirez-Erosa I, Hickie RA, Dunlop DM, Bishop GG, Deibert LK (2011) Antiproliferative activity of *Saponaria vaccaria* constituents and related compounds. Fitoterapia 83:170–181
- Baruah CC, Pal SK, Baruah AG, Roy JD, Buragohain B, Bora RS, Lahon LC (2009) Analgesic activity of methanolic extract of *Drymaria cordata* Willd Caryophyllaceae. Pharmacologyonline 2:470–476
- Bathori M, Varga E, Szendrei K, Lafont R (1990) Isolation and identification of new edcysteroids from the Caryophyllaceae. J Nat Prod 5:279–293
- Báthori M, Lafont R, Girault JP, Máthé I (2001) Structural diversity of ecdysteroids of *Lychnis flos-cuculi*. Acta Pharm Hung 71:157–167
- Bechlem H, Mencherini T, Bouheroum M, Benayache S, Cotugno R, Braca A, De Tommasi N (2017) New constituents from *Gymnocarpus decander*. Planta Med 83:1200–1206
- Blaut M, Schoefer L, Braune A (2003) Transformation of flavonoids by intestinal microorganisms. Int J Vitam Nutr Res 73:79–87
- Boguslavskaya LI (1976) Phenolic compounds of *Dianthus platycodon*. Chem Nat Compd 12:485
- Boguslavskaya LI, Dem'yanenko SI, Salam DK (1983) Flavonoids of some species of the genus *Dianthus*. Khimiya Prirodnikh Soedineni 61:366
- Boguslavskaya LI, Tikhonov AI, Pashnev PD, Zhemal B, Sklyar VI (1985a) C-glycosides of *Stellaria holostea*. Khim Prir Soedin 21:385
- Boguslavskaya LI, Tikhonov AI, Pashnev PD, Jemal B, Sklyar VI (1985b) Flavonoid compounds of *Herniaria polygama*. Chem Nat Compd 21:386–411
- Böttger S, Melzig MF (2011) Triterpenoid saponins of the Caryophyllaceae and Illecebraceae family. Phytochem Lett 4:59–68
- Bouillant ML, de Arce FF, Favre-Bonvin J, Chopin J, Zoll A, Mathieu G (1979) Nouvelles C-glycosylflavones extraites de *Spergularia rubra*. Phytochemistry 18:1043–1047
- Braca A, Bader A, Siciliano T, De Tommasi N (2008) Secondary metabolites from *Paronychia argentea*. Magn Reson Chem 46:88–93
- Bracher F, Puzik A (2004) β -Carboline Alkaloids 9 [1]. Total synthesis of the β -carboline alkaloids arenarine A and (\pm)-arenarine B. J Heterocycl Chem 41:173–176
- Brahmachari G, Gorai D (2006) Progress in the research on naturally occurring flavones and flavonols: an overview. Curr Org Chem 10:873–898
- Brockington SF, Walker RH, Glover BJ, Soltis PS, Soltis DE (2011) Complex pigment evolution in the Caryophyllales. New Phytol 190:854–864
- Bustamante-Rangel M, Delgado-Zamarreño MM, Pérez-Martin L, Rodríguez-Gonzalo E, Dominguez-Alvarez J (2018) Analysis of isoflavones in foods. Compr Rev Food Sci Food Saf 17:391–411
- Cambie RC (1959) Identity of isovitexin (“homovitexin”) and saponaretin. Chem Ind 1959:87–88
- Cheikh-Ali S, Farman M, Lacaille-Dubois MA, Semmar N (2019) Structural organization of saponins in Caryophyllaceae. Phytochem Rev 18:405–441
- Chen Q, Luo JG, Kong LY (2010a) Triterpenoid saponins from *Gypsophila altissima* L. Chem Pharm Bull 58:412–414
- Chen YF, Kuo PC, Chan HH, Kuo IJ, Lin FW, Su CR, Wu TS (2010b) β -carboline alkaloids from *Stellaria dichotoma* var. *lanceolata* and their anti-inflammatory activity. J Nat Prod 73:1993–1998
- Chen R, Qi QL, Wang MT, Li QY (2016) Therapeutic potential of naringin: an overview. Pharm Biol 54:3203–3210
- Cheng YX, Zhou J, Tan NH, Teng RW, Lu Y, Wang C, Zheng QT (2002) Isolation and characterization of brachystemidines A-E, novel alkaloids from *Brachystemma calycinum*. J Nat Prod 65:750–752
- Cheriti A, Sekkoum K (1996) Flavonoids from *Herniaria mauritanica*. Indian J Pharm Sci 58:203–204
- Cho HJ, Park JHY (2013) Kaempferol induces cell cycle arrest in HT-29 human colon cancer cells. J Cancer Prev 18:257–263
- Cho JY, Kim MS, Lee YG, Jeong HY, Lee HJ, Ham KS, Moon JH (2016) A phenyl lipid alkaloid and flavone C-diglucosides from *Spergularia marina*. Food Sci Biotechnol 25:63–69
- Chopin MJ, Bouillant ML, Wagner H, Galle K (1974) Endgültige struktur von schaftosid aus *Silene schafta*. Phytochemistry 13:2583–2586
- Chou S, Everngam MC, Beck JJ (2008) Allelochemical phenolic acids from *Gypsophila paniculata*. J Undergraduate Chem Res 7:2–4
- Clarkson TB (2002) Soy, soy phytoestrogens and cardiovascular disease. J Nutr 132:566S–569S
- Cook NC, Samman S (1996) Flavonoids—chemistry, metabolism, cardioprotective effects, and dietary sources. Nutritional Biochemistry 7:66–76
- Cosme P, Rodríguez AB, Espino J, Garrido M (2020) Plant phenolics: bioavailability as a key determinant of their potential health-promoting applications. Antioxidants 9:1–20
- Crozier A, Del Rio D, Clifford MN (2010) Bioavailability of dietary flavonoids and phenolic compounds. Mol Aspects Med 31:446–467
- Cui YL, Shen N, Zhao JQ, Dang J (2017a) Phytochemical constituents of *Arenaria kansuensis*. Chem Nat Compd 53:1002–1004
- Cui Y, Shen N, Yuan X, Dang J, Shao Y, Mei L, Liu Z (2017b) Two-dimensional chromatography based on on-line HPLC-DPPH bioactivity-guided assay for the preparative

- isolation of analogue antioxidant compound from *Arenaria kansuensis*. J Chromatogr B 1046:81–86
- Cui Y, Tao Y, Wang S (2018) Antihypoxic activities of constituents from *Arenaria kansuensis*. Phytomedicine 38:175–182
- Cui Y, Shao Y, Wang Q, Mei L, Tao Y (2019) Purification of flavonolignan diastereoisomers from *Arenaria kansuensis* by two-dimensional liquid chromatography combined with solid-phase extraction. J Chromatogr Sci 57:1–8
- Curir P, Dolci M, Lanzotti V, Tagliatalata-Scafati O (2001) Kaempferide triglycoside: A possible factor of resistance of carnation (*Dianthus caryophyllus*) to *Fusarium oxysporum* f. sp. *dianthi*. Phytochemistry 56:717–721
- Curir P, Lanzotti V, Dolci M, Dolci P, Pasini C, Tollin G (2003) Purification and properties of a new *S*-adenosyl-L-methionine: flavonoid 4'-*O*-methyltransferase from carnation (*Dianthus caryophyllus* L.). Eur J Biochem 270:3422–3431
- Curir P, Dolci M, Galeotti F (2005) A phytoalexin-like flavonol involved in the carnation (*Dianthus caryophyllus*)—*Fusarium oxysporum* f. sp. *dianthi* pathosystem. J Phytopathol 153:65–67
- Dai J, Dan W, Schneider U, Wang J (2018) β -Carboline alkaloid monomers and dimers: occurrence, structural diversity, and biological activities. Eur J Med Chem 157:622–656
- Darmograi VN (1977) Flavonoids of plants of the genera *Silene* and *Otites adans*, family Caryophyllaceae. Chem Nat Compd 13:102–103
- Darmograi VN (1979) Flavonoids of some species of the genera *Arenaria* and *Cerastium*. Khim Prir Soedin 1:93255
- de Andrade Teles RB, Diniz TC, Pinto TCC, de Oliveira Júnior RG, Silva MG, de Lavour EM, Fernandes AWC, de Oliveira AP, de Almeida Ribeiro FPR, da Silva AAM, Cavalcante TCF, Quintans Júnior LJ, da Silva Almeida JRG (2018) Flavonoids as therapeutic agents in Alzheimer's and Parkinson's diseases: a systematic review of preclinical evidences. Oxid Med Cell Longev 21:1–21
- del Valle JC, Buide ML, Casimiro-Soriguer I, Whittall JB, Narbona E (2015) On flavonoid accumulation in different plant parts: Variation patterns among individuals and populations in the shore campion (*Silene littorea*). Front Plant Sci 6:1–13
- Devkota HP, Fukusako K, Ishiguro K, Yahara S (2013) Flavone C-glycosides from *Lychnis senno* and their antioxidative activity. Nat Prod Commun 8:1413–1414
- Dhanya R, Arun KB, Nisha VM, Syama HP, Nisha P, Santhosh Kumar TR, Jayamurthy P (2015) Preconditioning L6 muscle cells with naringin ameliorates oxidative stress and increases glucose uptake. PLoS ONE 10:e0132429
- Di Lorenzo Ch, Colombo F, Biella S, Stockley C, Restani P (2021) Polyphenols and human health: the role of bioavailability. Nutrients 13:1–30
- Dinan L, Balducci C, Guibout L, Lafont R (2020) Small-scale analysis of phytoecdysteroids in seeds by HPLC/DAD-MS for the identification and quantification of specific analogues, dereplication and chemotaxonomy. Phytochem Anal 31:1–19
- Ding Z, Zhou J, Tan N (1999) A novel flavonoid glycoside from *Drymaria diandra*. Planta Med 65:578–579
- Ding Z, Zhou J, Tan N, Teng R (2000) Two new cyclic peptides from *Drymaria diandra*. Planta Med 66:386–388
- Ding ZT, Yang XQ, Cao QE, Li F (2005) Three new flavone glycosides from *Drymaria diandra* Bl. J Integr Plant Biol 47:1140–1144
- Dong Q, Huang Y, Qiao S (2007) Studies on chemical constituents from *Stellaria media*. Chin Mater Med 32:1048–1051
- Dötterl S, Jürgens A (2005) Spatial fragrance patterns in flowers of *Silene latifolia*: Lilac compounds as olfactory nectar guides? Plant Syst Evol 255:99–109
- Dubois M, Zoll A, Bouillant M, Delaveau P (1982) Di-C-Glycosylflavones du *Cerastium arvense* ssp. *arvense* nouvelles pour les Caryophyllaceae. Planta Med 46:56–57
- Dubois MA, Zoll A, Chopin J (1983) 7,2''-di-*O*-Glycosyl-6-C-glycosylflavones from *Cerastium arvense*. Phytochemistry 22:2879–2880
- Dubois MA, Zoll A, Markham KR, Bouillant ML, Dellamonica G, Chopin J (1984) 6-C- β -D-glucopyranosyl-8-C- β -D-galactopyranosylapigenin from *Cerastium arvense*. Phytochemistry 23:706–707
- Dubois MA, Zoll A, Chopin J (1985) Isomollupentin-*O*-glucosides from *Cerastium arvense*. Phytochemistry 24:1077–1080
- Egert S, Bösny-Westphal A, Seiberl J, Kurbitz C, Settler U, Plachta-Danielzik S, Wagner AE, Frank J, Schrezenmeier J, Rimbach G, Wolfram S, Müller MJ (2009) Quercetin reduces systolic blood pressure and plasma oxidised low-density lipoprotein concentrations in overweight subjects with a high-cardiovascular disease risk phenotype: a double-blinded, placebo-controlled cross-over study. Br J Nutr 7:1065–1074
- El Mabruki K, Klemper AV, Kaukhova IE, Sorokin VV (2014) Establishment of rupturewort (*Herniaria glabra*) herb identity characteristics and quality indices. Pharmacia 6:21–24
- Elbandy M, Miyamoto T, Lacaille-Dubois MA (2007) Sulfated lupane triterpene derivatives and a flavone C-glycoside from *Gypsophila repens*. Chem Pharm Bull 55:808–811
- El-Dien OG, Shawky E, Aly AH, Abdallah RM, Abdel-Salam NA (2013) A validated high-performance thin-layer chromatography (HPTLC) method for the quantitative determination of tricrin in two *Spergularia* Species. Am J Anal Chem 4:668–673
- Elgamal MHA, Soliman HSM, Karawya MS, Mikhova B, Duddeck H (1995) Isolation of triterpene saponins from *Gypsophila capillaris*. Phytochemistry 38:1481–1485
- Elhagali G, Abozeed A, Abdelnaser K, Youssif Y (2019) Investigation of bioactive constituents and biological activities of different fractions from *Herniaria hemistemon*. J Gay Al-Azhar Bull Sci 30:67–80
- El-Hawary SS, Mubarek MM, Lotfy AR, Hassan AR, Sobeh M, Okba MM (2020) Validation of antidiabetic potential of *Gymnocarpus decandrus* Forssk. Nat Prod Res 13:1–6
- Ferreres F, Gil-Izquierdo A, Vinholes J, Grosso C, Valentão P, Andrade PB (2011) Approach to the study of C-glycosyl flavones acylated with aliphatic and aromatic acids from *Spergularia rubra* by high-performance liquid chromatography-photodiode array detection/electrospray ionization multi-stage mass spectrometry. Rapid Commun Mass Spectrom 25:700–712
- Fukui Y, Tanaka Y, Kusumi T, Iwashita T, Nomoto K (2003) A rationale for the shift in colour towards blue in transgenic

- carnation flowers expressing the flavonoid 3',5'-hydroxylase gene. *Phytochemistry* 63:15–23
- Galeotti F, Barile E, Curir P, Dolci M, Lanzotti V (2008a) Flavonoids from carnation (*Dianthus caryophyllus*) and their antifungal activity. *Phytochem Lett* 1:44–48
- Galeotti F, Barile E, Lanzotti V, Dolci M, Curir P (2008b) Quantification of major flavonoids in carnation tissues (*Dianthus caryophyllus*) as a tool for cultivar discrimination. *Z Naturforsch* 63C:161–168
- Ganesan K, Xu BJ (2017) Molecular targets of vitexin and isovitexin in cancer therapy: a critical review. *Ann N Y Acad Sci* 1401:102–113
- Ganeshpurkar A, Saluja AK (2017) The pharmacological potential of rutin. *Saudi Pharm J* 25:149–164
- Ganeshpurkar A, Saluja A (2019) The pharmacological potential of hesperidin. *Indian J Biochem Biophys* 56:287–300
- Gevrenova R, Bardarov K, Bouguet-Bonnet S, Voynikov Y, Balabanova V, Zheleva-Dimitrova D, Henry M (2018) A new liquid chromatography-high resolution orbitrap mass spectrometry-based strategy to characterize glucuronide oleanane-type triterpenoid carboxylic acid 3, 28-*O*-bidesmosides (GOTCAB) saponins. A case study of *Gypsophila glomerata* Pall ex M. B. (Caryophyllaceae). *J Pharm Biomed Anal* 159:567–581
- Griffiths LA (1959) On the distribution of gentisic acid in green plants. *J Exp Bot* 10:437–442
- Grundmann O, Wang J, McGregor GP, Butterweck V (2008) Anxiolytic activity of a phytochemically characterized *Passiflora incarnata* extract is mediated via the GABAergic system. *Planta Med* 74:1769–1773
- Gullón B, Lú-Chau TA, Moreira MT, Lema JM, Eibes G (2017) Rutin: A review on extraction, identification and purification methods, biological activities and approaches to enhance its bioavailability. *Trends Food Sci Technol* 67:220–235
- Hegnauer R (1964) *Chemotaxonomy of plants*, vol 18. Springer Basel AG, Basel, p 379
- Hegnauer R (1989) *Caryophyllaceae. Chemotaxonomy of Plants*, vol 30. Springer Basel AG, Basel, pp 215–220
- Heim KE, Tagliaferro AR, Bobilya DJ (2002) Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *J Nutr Biochem* 13:572–584
- Heinsbroek R, van Brederode J, van Nigtevecht G, Maas J, Kamsteeg J, Besson E, Chopin J (1980) The 2''-*O*-glucosylation of vitexin and isovitexin in petals of *Silene alba* is catalysed by two different enzymes. *Phytochemistry* 19:1935–1937
- Hollman PCH (2004) Absorption, bioavailability, and metabolism of flavonoids. *Pharm Biol* 42:74–83
- Hostetler GL, Ralston RA, Schwartz SJ (2017) Flavones: food sources, bioavailability, metabolism, and bioactivity. *Adv Nutr* 28:423–435
- Hsieh PW, Chang FR, Lee KH, Hwang TL, Chang SM, Wu YC (2004a) A new anti-HIV alkaloid, drymaritin, and a new C-glycoside flavonoid, diandraflavone, from *Drymaria diandra*. *J Nat Prod* 67:1175–1177
- Hsieh PW, Chang FR, Wu CC, Wu KY, Li CM, Wang WY, Wu YC (2004b) Selective inhibition of collagen-induced platelet aggregation by a cyclic peptide from *Drymaria diandra*. *Helv Chim Acta* 87:57–66
- Huang QF, Zhang SJ, Zheng L, Liao M, He M, Huang RB, Lin X (2012) Protective effect of isoorientin-2'-*O*- α -L-arabinopyranosyl isolated from *Gypsophila elegans* on alcohol induced hepatic fibrosis in rats. *Food Chem Toxicol* 50:1992–2001
- Hussein IA, Srivedavyasari R, El-Hela AA, Mohammad AI, Ross SA (2019) Antimicrobial secondary metabolites from *Silene rubella* growing in Egypt. *J Biomed Pharm Res* 8:81–84
- Hussein IA, Srivedavyasari R, El-Hela AA, Mohammad AI, Ross SA (2020) Chemical constituents from *Silene schimperiana* Boiss. belonging to Caryophyllaceae and their chemotaxonomic significance. *Biochem Syst Ecol* 92:1–4
- Iacopini P, Baldi M, Storchi P, Sebastiani L (2008) Catechin, epicatechin, quercetin, rutin and resveratrol in red grape: content, *in vitro* antioxidant activity and interactions. *J Food Compos Anal* 21:589–598
- Imran M, Rauf A, Shah ZA, Saeed F, Imran A, Arshad MU, Mubarak MS (2018) Chemo-preventive and therapeutic effect of the dietary flavonoid kaempferol: a comprehensive review. *Phytother Res* 2018:1–13
- Imran M, Rauf A, Abu-Izneid T, Nadeem M, Shariati MA, Khan IA, Mubarak MS (2019a) Luteolin, a flavonoid, as an anticancer agent: a review. *Biomed Pharmacother* 112:108612
- Imran M, Salehi B, Sharifi-rad J, Gondal TA, Saeed F, Imran A, Estevinho LM (2019b) Kaempferol: A key emphasis to its anticancer potential. *Molecules* 24:1–16
- Imran M, Aslam GT, Atif M, Shahbaz M, Batool QT, Hanif MM, Sharifi-Rad J (2020) Apigenin as an anticancer agent. *Phytother Res* 26:1–17
- Itokawa H, Yun Y, Morita H, Takeya K, Yamada K (1995) Estrogen-like activity of cyclic peptides from *Vaccaria segetalis* extracts. *Planta Med* 61:561–562
- Iwashina T, Yamaguchi MA, Nakayama M, Onozaki T, Yoshida H, Kawanobu S, Okamura M (2010) Kaempferol glycosides in the flowers of carnation and their contribution to the creamy white flower color. *Nat Prod Commun* 5:1903–1906
- Jakimiuk K, Strawa JW, Granica S, Tomczyk M (2020) Flavonoids from the aerial parts of *Scleranthus perennis*. T20 PSE Conference Liverpool 2020, "Contemporary Natural Products Discovery Research", 6.03.2020, Liverpool, United Kingdom, p. 57
- Jia AQ, Tan NH, Yang YP, Wu SG, Wang LQ, Zhou J (2004) Cyclopeptides from three arctic Caryophyllaceae plants, chemotaxonomy and distribution significance of Caryophyllaceae cyclopeptides. *Acta Bot Sin* 46:625–630
- Jovanović O, Radulović N, Palić R, Zlatković B (2009) Volatiles of *Minuartia recurva* (All.) Schinz et Thell. subsp. *recurva* (Caryophyllaceae) from Serbia. *J Essent Oil Res* 21:429–432
- Jung HJG, Batzli GO, Seigler DS (1979) Patterns in the phytochemistry of arctic plants. *Biochem Syst Ecol* 7:203–209
- Jürgens A (2004) Flower scent composition in diurnal *Silene* species (Caryophyllaceae): phylogenetic constraints or adaptation to flower visitors? *Biochem Syst Ecol* 32:841–859
- Jürgens A, Witt T, Gottsberger G (2002) Flower scent composition in night-flowering *Silene* species (Caryophyllaceae). *Biochem Syst Ecol* 30:383–397

- Jürgens A, Witt T, Gottsberger G (2003) Flower scent composition in *Dianthus* and *Saponaria* species (Caryophyllaceae) and its relevance for pollination biology and taxonomy. *Biochem Syst Ecol* 31:345–357
- Kamsreoo J, van Brederode J, van Nigtevecht G (1980) Genetical and biochemical evidence that the hydroxylation pattern of the anthocyanin B-ring *Silene dioica* is determined at the *p*-coumaroyl-coenzyme a stage. *Phytochemistry* 19:1459–1462
- Kamsteeg J, van Brederode J, van Nigtevecht G (1976) Pleiotropic effect of a pelargonidin-hydroxylation gene in *Silene dioica*. *Phytochemistry* 15:1917–1918
- Ke JY, Cole RM, Hamad EM, Hsiao YH, Cotten BM, Powell KA, Belury MA (2016) Citrus flavonoid, naringenin, increases locomotor activity and reduces diacylglycerol accumulation in skeletal muscle of obese ovariectomized mice. *Mol Nutr Food Res* 60:313–324
- Kew Science (2020) The Royal Botanic Gardens, Great Britain. <http://plantsoftheworldonline.org> Accessed 27 Nov 2020
- Kılınc H, Masullo M, Bottone A, Karayıldırım T, Alankuş Ö, Piacente S (2019) Chemical constituents of *Silene montbretiana*. *Nat Prod Res* 33:335–339
- Kim DW, Hwang IK, Lim SS, Yoo KY, Li H, Kim YS, Kwon DY, Moon WK, Kim DW, Won MH (2009) Germinated Buckwheat extract decreases blood pressure and nitrotyrosine immunoreactivity in aortic endothelial cells in spontaneously hypertensive rats. *Phytother Res* 23:993–998
- Kim YB, Reed DW, Covello PS (2015) Production of triterpenoid saponins in hairy root cultures of *Silene vulgaris*. *Nat Prod Commun* 10:1919–1922
- Kirilov V, Stikhareva T, Suleimen Y, Serafimovich M, Kabanova S, Mukanov B (2017) Chemical composition of the essential oil from carnation coniferous (*Dianthus acicularis* Fisch. ex Ledeb) growing wild in Northern Kazakhstan. *Nat Prod Res* 31:117–123
- Kitanov GM (1992) Phenolic acids and flavonoids from *Stellaria media* (L.) Vill. (Caryophyllaceae). *Pharmazie* 47:470–471
- Koike K, Jia Z, Nikaido T (1998) Triterpenoid saponins from *Vaccaria segetalis*. *Phytochemistry* 47:1343–1349
- Koike K, Jia Z, Nikaido T (1999) New triterpenoid saponins and saponinins from *Saponaria officinalis*. *J Nat Prod* 62:1655–1659
- Kozachok S, Pecio Ł, Kolodziejczyk-Czepas J, Marchyshyn S, Nowak P, Mołdoch J, Oleszek W (2018) γ -Pyrone compounds: flavonoids and maltol glucoside derivatives from *Herniaria glabra* L. collected in the Ternopil region of the Ukraine. *Phytochemistry* 152:213–222
- Kozłowska J, Potaniec B, Zarowska B, Aniol M (2017) Synthesis and biological activity of novel *o*-alkyl derivatives of naringenin and their oximes. *Molecules* 22:1–14
- Krasteva IN, Popov IS, Balabanova VI, Nikolov SD, Pencheva IP (2008) Phytochemical study of *Gypsophila trichotoma* Wend. (Caryophyllaceae). *Quim Nova* 31:1125–1126
- Kremer D, Košir IJ, Potočnik T, Rogulj N, Načinović K, Randić M, Srećec S, Jurišić-Grubešić R (2021) Phenolic compounds in two subspecies of *Drypis spinosa* L. (Caryophyllaceae) growing in Croatia. *Acta Bot Croatica*. <https://doi.org/10.37427/botcro-2020-015>
- Križová L, Dadáková K, Kašparovská J, Kašparovský T (2019) Isoflavones. *Molecules* 24:1–28
- Królikowska M, Szymańska M, Wolbiś M (1983) Rhamnazin 3-rutinoside from *Herniaria ciliolata* Meld. spp. *robusta* Chaudhri. *Acta Pol Pharm* 40:643–648
- Kubitzki K (1993) In: Kubitzki K, Rohwer JG, Bittrich V (eds) Flowering plants dicotyledons. Springer, Berlin
- Kukongviriyapan U, Sompamit K, Pannangpetch P, Kukongviriyapan V, Donpunha W (2012) Preventive and therapeutic effects of quercetin on lipopolysaccharide-induced oxidative stress and vascular dysfunction in mice. *Can J Physiol Pharmacol* 90:1345–1353
- Kulevanova S, Stefova M, Kadifkova Panovska T, Stafilov T (2003) HPLC identification and determination of myricetin, quercetin, kaempferol and total flavonoids in herbal drugs. *Maced Pharm Bull* 48:25–30
- Kumar P, Khanna P (1994) Flavonoids from *Saponaria vaccaria* Linn. *Indian J Plant Physiol* 37:76–78
- Kumar RK, Herbert C, Foster PS (2008) The “classical” ovalbumin challenge model of asthma in mice. *Curr Cancer Drug Targets* 9:485–494
- Kuwayama S, Nakata M, Godo T, Nakano M (2005) Analyses of anthocyanidins and anthocyanins in flower petals of *Lychnis senno* and its related species (Caryophyllaceae). *Bull Fac Agric Niigata Univ* 58:35–38
- Lam KY, Ling APK, Koh RY, Wong YP, Say YH (2016) A review on medicinal properties of orientin. *Adv Pharmacol Sci* 2016:4104595
- Larhsini M, Marston A, Hostettmann K (2003) Triterpenoid saponins from the roots of *Silene cucubalus*. *Fitoterapia* 74:237–241
- Liang X, Li Y, Fan H, Huang W, Zhang H, Cui Y, Song X (2019) Chemical constituents from the roots and rhizomes of *Silene tatarinowii* Regel. *Biochem Syst Ecol* 86:103932
- Lin X, Chen Y, Lv S, Tan S, Zhang S, Huang R, Huang Q (2015) *Gypsophila elegans* isoorientin attenuates CCl₄-induced hepatic fibrosis in rats via modulation of NF- κ B and TGF- β 1/Smad signaling pathways. *Int Immunopharmacol* 28:305–312
- Lin X, Wei J, Chen Y, He P, Lin J, Tan S, Huang Q (2016) Isoorientin from *Gypsophila elegans* induces apoptosis in liver cancer cells via mitochondrial-mediated pathway. *J Ethnopharmacol* 187:187–194
- Litvinenko VI, Amanmuradov K, Abubakirov NK (1967) Glycosides of *Vaccaria segetalis* IV Isosaponarin. *Chem Nat Compd* 3:131–134
- Liu XX, Wang L, Wang Q, Qiu B (2007) Chemical constituents from root of *Psammosilene tunicoides*. *China J Chin Mater Med* 32:921–923
- Liu Z, Lindemeyer AK, Liang J, Wallner M, Shao XM, Shao Y, Olsen RW (2018) Flavonoids isolated from Tibetan medicines, binding to GABA_A receptor and the anticonvulsant activity. *Phytomedicine* 50:1–7
- Liu Y, Song FM, Ma ST, Moro A, Feng WY, Liao SJ, Liu Q (2019) Vaccarin prevents titanium particle-induced osteolysis and inhibits RANKL-induced osteoclastogenesis by blocking NF- κ B and MAPK signaling pathways. *J Cell Physiol* 234:13832–13842
- Luo JG, Cao LH, Kong LY (2012) Two new β -carboline-type alkaloids from *Stellaria dichotoma* var. *lanceolata*. *Chin Chem Lett* 23:1385–1388

- Ma X, Wu C, Wang W, Li X (2006) Peptides from plants: a new source for antitumor drug research. *Asian J Tradit Med* 1:85–90
- Ma L, Gu YC, Luo JG, Wang JS, Huang XF, Kong LY (2009) Triterpenoid saponins from *Dianthus versicolor*. *J Nat Prod* 72:640–644
- Ma LY, Liu RH, Xu XD, Yu MQ, Zhang Q, Liu HL (2010) The pharmacokinetics of C-glycosyl flavones of Hawthorn leaf flavonoids in rat after single dose oral administration. *Phytomedicine* 17:640–645
- Maleš Ž, Crkvenčić M, Pilepić KH, Herenda F (2013) Investigation of flavonoids, phenolic acids and amino acids of smooth raptunewort—*Herniaria glabra* L. *Farm Glas* 69:673–684
- Maliński MP, Michalska AD, Tomczykowa M, Tomczyk M, Thiem B (2014) Ragged Robin (*Lychnis flos-cuculi*)—a plant with potential medicinal value. *Rev Bras* 24:722–730
- Maliński MP, Kikowska M, Kruska D, Napierała M, Florek E, Śliwińska E, Thiem B (2019) Various *in vitro* systems of Ragged Robin (*Lychnis flos-cuculi* L.): a new potential source of phytoecdysteroids? *Plant Cell Tissue Organ Cult* 139:39–52
- Mamadalieva NZ, Zibareva LN, Lafont R, Dainan L, Saatov Z (2004) Phytoecdysteroids from the *Silene* genus. *Chem Nat Compd* 40:574–578
- Mamadalieva NZ, Egamberdieva D, Lafont R, Girault JP (2008) Phytoecdysteroids and antibacterial activity of the plant *Coronaria flos-cuculi*. *Chem Nat Compd* 44:404–406
- Mamadalieva NZ, El-Readi MZ, Janibekov AA, Tahrani A, Wink M (2011) Phytoecdysteroids of *Silene guntensis* and their *in vitro* cytotoxic and antioxidant activity. *Z Naturforsch* 66C:215–224
- Mamadalieva NZ, Lafont R, Wink M (2014) Diversity of secondary metabolites in the genus *Silene* L. (Caryophyllaceae)—structures, distribution, and biological properties. *Diversity* 6:415–499
- Mandal P, Misra TK, Ghosal M (2009) Free-radical scavenging activity and phytochemical analysis in the leaf and stem of *Drymaria diandra* Blume. *Int J Integr Biol* 7:80–84
- Martineti V, Tognarini I, Azzari C, Sala SC, Clematis F, Dolci M, Curir P (2010) Inhibition of *in vitro* growth and arrest in the G0/G1 phase of HCT8 line human colon cancer cells by kaempferide triglycoside from *Dianthus caryophyllus*. *Phytother Res* 24:1302–1308
- Mbark AN, Charrouf Z, Guillaume D, Kol O (1999) New glycosides from *Herniaria fontanesii*. *Stud Plant Sci* 6:314–319
- Melnyk MV, Vodoslavskiy VM, Obodianskyi MA (2018) Research of phenolic compounds of *Ruta graveolens* L. and *Stellaria media* (L.) Vill. *Asian J Pharm Clin Res* 11:152–156
- Meng Y, Whiting P, Zibareva L, Bertho G, Girault JP, Lafont R, Dinan L (2001) Identification and quantitative analysis of the phytoecdysteroids in *Silene* species (Caryophyllaceae) by high-performance liquid chromatography: Novel ecdysteroids from *S. pseudotites*. *J Chromatogr A* 935:309–319
- Meselhy MR, Aboutabl ES (1997) Hopane-type saponins from *Polycarpon succulentum* growing in Egypt. *Phytochemistry* 44:925–929
- Mihaylova D, Vrancheva R, Desseva I, Ivanov I, Dincheva I, Popova M, Popova A (2018) Analysis of the GC-MS of volatile compounds and the phytochemical profile and antioxidant activities of some Bulgarian medicinal plants. *Z Naturforschung* 74C:45–54
- Mikšátková P, Ancheeva E, Hejtmánková K, Teslov L, Lapčík O (2014) Determination of flavonoids in *Stellaria* by high-performance liquid chromatography-tandem mass spectrometry. *Anal Lett* 47:2317–2331
- Morales P, Carvalho AM, Sánchez-Mata MC, Cámara M, Molina M, Ferreira ICFR (2012) Tocopherol composition and antioxidant activity of Spanish wild vegetables. *Genet Resour Crop Evol* 59:851–863
- Mubarek MM (2019) Pharmacognostical studies on *Gymnocarpus decandrus* Forrsk. growing at North Western Coast in Egypt, Doctoral dissertation, Cairo University
- Nakano T, Sugimoto S, Matsunami K, Otsuka H (2011) Dianthosaponins A-F, triterpene saponins, flavonoid glycoside, aromatic amide glucoside and γ -pyrone glucoside from *Dianthus japonicus*. *Chem Pharm Bull* 59:1141–1148
- Nakayama M, Koshioka M, Yoshida H, Kan Y, Fukui Y, Koike A, Yamaguchi MA (2000) Cyclic malyl anthocyanins in *Dianthus caryophyllus*. *Phytochemistry* 55:937–939
- Nerio LS, Olivero-Verbel J, Stashenko E (2010) Repellent activity of essential oils: A review. *Biores Technol* 101:372–378
- Niemann GJ (1984) Leaf flavonoid glycosylation and sprout morphogenesis in *Silene pratensis* influenced by the spectral composition of light. *J Plant Physiol* 115:311–318
- Niemann GJ (1993) The anthranilamide phytoalexins of the Caryophyllaceae and related compounds. *Phytochemistry* 34:319–328
- Nono RN, Nguerefack-Mbuyo EP, Nzowa LK, Ponou BK, Teponno RB, Nguerefack TB, Park HJ (2016) Antioxidant C-glycosylflavones of *Drymaria cordata* (Linn.) Willd. *Arch Pharmacol Res* 39:43–50
- Obmann A, Zehl M, Purevsuren S, Narantuya S, Reznicek G, Kletter C, Glasl S (2011b) Quantification of flavonoid glycosides in an aqueous extract from the traditional Mongolian medicinal plant *Dianthus versicolor* Fisch. *J Sep Sci* 34:292–298
- Obmann A, Werner I, Presser A, Zehl M, Swoboda Z, Purevsuren S, Glasl S (2011a) Flavonoid C- and O-glycosides from the Mongolian medicinal plant *Dianthus versicolor* Fisch. *Carbohydr Res* 346:1868–1875
- Obmann A, Purevsuren S, Zehl M, Kletter C, Reznicek G, Narantuya S, Glasl S (2012) HPLC determination of flavonoid glycosides in Mongolian *Dianthus versicolor* Fisch. (Caryophyllaceae) compared with quantification by UV spectrophotometry. *Phytochem Anal* 23:254–259
- Ogata J, Itoh Y, Ishida M, Yoshida H, Ozeki Y (2004) Cloning and heterologous expression of cDNAs encoding flavonoid glucosyltransferases from *Dianthus caryophyllus*. *Plant Biotechnol* 21:367–375
- Olennikov DN (2020) Silenerepin—a new C-glycosylflavone from *Silene repens*. *Chem Nat Compd* 56:423–426
- Pacifico S, Scognamiglio M, D'Abrosca B, Piccolella S, Tsafantakis N, Gallicchio M, Fiorentino A (2010) Spectroscopic characterization and antiproliferative activity on HepG2 human hepatoblastoma cells of flavonoid C-

- glycosides from *Petrorhagia velutina*. J Nat Prod 73:1973–1978
- Panche AN, Diwan AD, Chandra SR (2016) Flavonoids: an overview. J Nutr Sci 5:1–15
- Pei R, Liu X, Bolling B (2020) Flavonoids and gut health. Curr Opin Biotechnol 61:153–159
- Peng Y, Gan R, Li H, Yang M, McClements DJ (2020) Absorption, metabolism, and bioactivity of vitexin: recent advances in understanding the efficacy of an important nutraceutical. Critical Reviews in Food Science and Nutrition 1–16
- Plant Database (2020) United States Department of Agriculture, United States of America. <https://plants.sc.egov.usda.gov>. Accessed 27 Nov 2020
- Powers JM, Seco R, Faiola CL, Sakai AK, Weller SG, Campbell DR, Guenther A (2020) Floral scent composition and fine-scale timing in two moth-pollinated Hawaiian *Schiedea* (Caryophyllaceae). Front Plant Sci 11:1–16
- Praveena R, Sadasivam K, Deepha V, Sivakumar R (2014) Antioxidant potential of orientin: a combined experimental and DFT approach. J Mol Struct 1061:114–123
- Qi P, Li Z, Chen M, Sun Z, Huang C (2013) Metabolism and tissue distribution study of *Vaccaria* seeds (Wang-Bu-Liu-Xing) in benign prostatic hyperplasia model rat: toward an in-depth study for its bioactive components. J Pharm Biomed Anal 85:218–230
- Qi P, Zhang F, Xue R, Li Z, Chen M, Sun Z, Huang C (2014) Identification of multiple constituents from seed of *Vaccaria segetalis* with an adsorbent-separation strategy based on liquid chromatography coupled to quadrupole time-of-flight mass spectrometry. Rapid Commun Mass Spectrom 28:1243–1257
- Radulović NS, Ristić MN, Ristić NR, Dekić VS, Dekić BR, Mladenović MZ (2018) The floral scent of *Dianthus cruentus* Griseb (Caryophyllaceae). Fac Univ Ser Phys Chem Technol 16:161
- Richardson M (1978) Flavonols and C-glycosylflavonoids of the caryophyllales. Biochem Syst Ecol 6:283–286
- Rizk AM (1986) Phytochemistry of the flora of Qatar. Scientific and Applied Research Centre, University of Qatar, Qatar
- Rogowska M, Lenart M, Srećec S, Ziaja M, Parzonko A, Bazyłko A (2017) Chemical composition, antioxidative and enzyme inhibition activities of chickweed herb (*Stellaria media* L., Vill.) ethanolic and aqueous extracts. Ind Crops Prod 97:448–454
- Sachdeva AK, Kuhad A, Chopra K (2014) Naringin ameliorates memory deficits in experimental paradigm of Alzheimer's disease by attenuating mitochondrial dysfunction. Pharmacol Biochem Behav 127:101–110
- Said RB, Hamed AI, Masullo M, Al-Ayed AS, Moustafa MFM, Mahalel UA, Piacente S (2019) Flavone C-glycosides from *Vaccaria pyramidata*: structure elucidation by spectroscopy and theoretical calculations. Phytochem Lett 29:119–124
- Sait S, Hamri-Zeghichi S, Boulekbache-Makhlouf L, Madani K, Rigou P, Brighenti V, Pellati F (2015) HPLC-UV/DAD and ESI-MS_n analysis of flavonoids and antioxidant activity of an Algerian medicinal plant: *Paronychia argentea* Lam. J Pharm Biomed Anal 111:231–240
- Salehi B, Fokou PVT, Sharifi-Rad M, Zucca P, Pezzani R, Martins N, Sharifi-Rad J (2019) The therapeutic potential of naringenin: A review of clinical trials. Pharmaceuticals 12:1–18
- Salt TA, Adler JH (1986) Dominance of Δ^7 -sterols in the family Caryophyllaceae. Lipids 21:754–758
- Sang SM, Xia ZH, Mao SL, Lao A, Chen ZL (2000) Studies on the flavonol glycosides from the seeds of *Vaccaria segetalis*. China J Chin Mater Med 25:221–222
- Sang S, Xia Z, Lao A, Cao L, Chen Z, Uzawa J, Fujimoto Y (2003b) Studies on the constituents of the seeds of *Vaccaria segetalis*. Heterocycles 59:811–821
- Sang S, Lao A, Chen Z, Uzawa J, Fujimoto Y (2003) In: Ho CT (ed.) Oriental foods and herbs. Oxford University Press, Washington
- Schmidt J, Bohme F, Adam G (1996) 24-Epibrassinolide from *Gypsophila perfoliata*. Z Naturforsch 51C:897–899
- Schweingruber FH (2007) Stem anatomy of Caryophyllaceae. Flora - Morphology, Distribution, Functional Ecology of Plants 202:281–292
- Seo C, Shin HS, Lee JE, Jung YW, Kim JK, Kwon JG, Hong SS (2020) Isolation and structure elucidation of siliendines A–D, new β -carboline alkaloids from *Silene seoulensis*. Phytochem Lett 36:58–62
- Seraya L, Birke K, Khimenko SV, Boguslavskaya L (1978) Flavonoid compounds of *Dianthus superbus*. Khim Prir Soedin 6:802–803
- Serra A, Macià A, Romero MP, Reguant J, Ortega N, Motilva MJ (2012) Metabolic pathways of the colonic metabolism of flavonoids (flavonols, flavones and flavanones) and phenolic acids. Food Chem 130:383–393
- Shafaghat A, Shafaghatlonbar M (2011) Antimicrobial activity and chemical constituents of the essential oils from flower, leaf and stem of *Gypsophila bicolor* from Iran. Nat Prod Commun 6:275–276
- Sharma A, Arora D (2012) Phytochemical and pharmacological potential of genus *Stellaria*: A review. J Pharm Res 5:3591–3596
- Shinjiro O, Junko M, Godo T, Kato Y (2009) Possibility for selective accumulation of polyphenolics in tissue cultures of *Senno* (*Lychnis senno* Siebold et Zucc.). Nat Prod Commun 4:377–380
- Shukla S, Gupta S (2010) Apigenin: a promising molecule for cancer prevention. Pharm Res 27:962–978
- Simeonova R, Kondeva-Burdina M, Vitcheva V, Krasteva I, Manov V, Mitcheva M (2014) Protective effects of the apigenin-O/C-diglucoside saponarin from *Gypsophila trichotoma* on carbone tetrachloride-induced hepatotoxicity *in vitro/in vivo* in rats. Phytomedicine 21:148–154
- Singh M, Kaur M, Silakari O (2014) Flavones: an important scaffold for medicinal chemistry. Eur J Med Chem 84:206–239
- Slavokhotova AA, Odintsova TI, Rogozhin EA, Musolyamov AK, Andreev YA, Grishin EV, Egorov TA (2011) Isolation, molecular cloning and antimicrobial activity of novel defensins from common chickweed (*Stellaria media* L.) seeds. Biochimie 93:450–456
- Smolyakova IM, Avdeenko SN, Kalinkina GI, Yusubov MS, Zibareva LN (2010) Analysis of the chemical composition of *Lychnis chalconica* cultivated in Western Siberia. Chem Plant Mater 2010:95–102
- Stich K, Eidenberger T, Wurst F, Forkmann G (1992) Enzymatic conversion of dihydroflavonols to flavan-3,4-diols

- using flower extracts of *Dianthus caryophyllus* L. (carnation). *Planta* 187:103–108
- Sun J, Yu JH, Song JL, Jiang CS, Yuan T, Zhang H (2019) Two new quinolone alkaloids from *Dianthus superbis* var. *superbus*. *Tetrahedron Lett* 60:161–163
- Taskin T, Bitis L (2013) Antioxidant activity of *Silene alba* subsp. *divaricata* and *Stellaria media* subsp. *media* from Caryophyllaceae. *Spatula* DD 3:1–5
- Thiem B, Kikowska M, Maliński MP, Kruszka D, Napierała M, Florek E (2016) Ecdysteroids: production in plant *in vitro* cultures. *Phytochem Rev* 16:603–622
- Thilakarathna SH, Rupasinghe HPV (2013) Flavonoid bioavailability and attempts for bioavailability enhancement. *Nutrients* 5:3367–3387
- Tlili H, Hanen N, Arfa AB, Neffati M, Boubakri A, Buonocore D, Doria E (2019) Biochemical profile and *in vitro* biological activities of extracts from seven folk medicinal plants growing wild in southern Tunisia. *PLoS ONE* 14:1–18
- Tomczyk M (2008) Preliminary phytochemical investigation of *Lychnis flos-cuculi* herbs. *J Nat Med* 62:473–475
- Tong Y, Luo JG, Wang R, Wang XB, Kong LY (2012) New cyclic peptides with osteoblastic proliferative activity from *Dianthus superbis*. *Bioorg Med Chem Lett* 22:1908–1911
- Tong H, Sun BG, ChangTao W, Sun XT, Xue Z (2014) Study on surfactant-assisted extraction process and preliminary structural analysis of total flavonoids from *Arenaria kansuensis* Maxim. *Food Res Dev* 35:14–18
- Tu Y, Zhu S, Wang J, Burstein E, Jia D (2019) Natural compounds in the chemoprevention of alcoholic liver disease. *Phytother Res* 33:2192–2212
- Ullah F, Ayaz A, Saqib S, Zaman W, Butt MA, Ullah A (2019) *Silene conoidea* L.: A review on its systematic, ethnobotany and phytochemical profile. *Plant Sci Today* 6:373–382
- Uma Devi P, Ganasoundari A, Vrinda B, Srinivasan KK, Unnikrishnan MK (2000) Radiation protection by the *Ocimum* flavonoids orientin and vicenin: mechanisms of action. *Radiat Res* 154:455–460
- Van Wyk BE, Wink M (2017) *Medicinal plants of the World*. Briza Publications, Pretoria
- Van Brederode J, van Genderen HH, Berendsen W (1982) Morphological effects of the flavone isovitexin in a non-glycosylating genotype of *Silene pratensis* (Caryophyllaceae). *Experientia* 38:929–931
- Van Dooren I, Foubert K, Bijttebier S, Theunis M, Velichkova S, Claeys M, Apers S (2016) Saponins and flavonoids from an infusion of *Herniaria hirsuta*. *Planta Med* 82:1576–1583
- Vardavas CI, Majchrzak D, Wagner KH, Elmadfa I, Kafatos A (2006) The antioxidant and phyloquinone content of wildy grown greens in Crete. *Food Chem* 99:813–821
- Vincken JP, Heng L, de Groot A, Gruppen H (2007) Saponins, classification and occurrence in the plant kingdom. *Phytochemistry* 68:275–297
- Vinholes J, Grosso C, Andrade PB, Gil-Izquierdo A, Valentão P, Pinho PGD, Ferreres F (2011) *In vitro* studies to assess the antidiabetic, anti-cholinesterase and antioxidant potential of *Spergularia rubra*. *Food Chem* 129:454–462
- Viskupičová J, Ondrejovič M, Šturdík E (2008) Bioavailability and metabolism of flavonoids. *J Food Nutr Res* 47:151–162
- Vitale DC, Piazza C, Melilli B, Drago F, Salomone S (2013) Isoflavones: estrogenic activity, biological effect and bioavailability. *Eur J Drug Metab Pharmacokinet* 38:15–25
- Vitcheva V, Simeonova R, Krasteva I, Yotova M, Nikolov S, Mitcheva M (2011) Hepatoprotective effects of saponarin, isolated from *Gypsophila trichotoma* Wend. on cocaine-induced oxidative stress in rats. *Redox Rep* 16:56–61
- Volodin VV, Volodina SO (2015) Floristic and molecular phylogenetic analysis of the distribution of phytoecdysteroids among plants of North-East Russia (Asteraceae and Caryophyllaceae). *Biol Med* 7:1
- Wang X, Dong H, Liu Y, Yang B, Wang X, Huang L (2011) Application of high-speed counter-current chromatography for preparative separation of cyclic peptides from *Vaccaria segetalis*. *J Chromatogr B* 879:811–814
- Wang G, Luo JG, Yang MH, Wang XB, Kong LY (2013) Six new cyclic peptides from the roots of *Gypsophila oldhamiana*. *Chem Pharm Bull* 61:489–495
- Wang W, Sun C, Mao L, Ma P, Liu F, Yang J, Gao Y (2016) The biological activities, chemical stability, metabolism and delivery systems of quercetin: a review. *Trends Food Sci Technol* 56:21–38
- Wink M (2011) *Biochemistry of Plant Secondary Metabolism*, 2nd edn. Wiley-Blackwell, Chichester
- Wink M (2015) Modes of action of herbal medicines and plant secondary metabolites. *Medicines* 2:251–286
- Wolf SJ, Denford KE, Packer JG (1979) A study of the flavonoids of the *Minuartia rossii* complex. *Can J Bot* 57:2374–2377
- Wu FE, Koike K, Nikaido T, Sakamoto Y, Ohmoto T, Ikeda K (1989) New β -Carboline alkaloids from a Chinese medicinal plant, *Arenaria kansuensis*. Structures of arenarines A, B, C, D. *Chem Pharm Bull* 37:1808–1809
- Wu FE, Koike K, Nikaido T, Ishii K, Ohmoto T, Ikeda K (1990) Terpenoids and flavonoids from *Arenaria kansuensis*. *Chem Pharm Bull* 38:2281–2282
- Yasukawa K, Yamanouchi S, Takido M (1981) Studies on the constituents in the water extracts of crude drugs. III. On the roots of *Stellaria dichotoma* L. var. *lanceolata* BGE. *Yakugaku Zasshi* 101:64–66
- Yayli N, Seymen H, Baltacı C (2001) Flavone C-glycosides from *Scleranthus uncinatus*. *Phytochemistry* 58:607–610
- Yayli N, Baltacı C, Genç H, Terzioğlu S (2002) Phenolic and flavone C-glycosides from *Scleranthus uncinatus*. *Pharm Biol* 40:369–373
- Yoshida H, Itoh Y, Ozeki Y, Iwashina T, Yamaguchi MA (2004) Variation in chalcononaringenin 2'-O-glucoside content in the petals of carnations (*Dianthus caryophyllus*) bearing yellow flowers. *Sci Hortic* 99:175–186
- Yücel TB, Yayli N (2018) GC/MS analysis and antimicrobial activity of the volatile compounds from *Dianthus carmelitarum* Reut. ex Boiss and *Dianthus calocephalus* Boiss. grown in Turkey. *J Agric Fac Ege Univ* 55:89–94
- Zanotti SD, de Abreu Ribeiro GK, Zeppone LC, Borges CT (2013) Orange juice and hesperidin promote differential innate immune response in macrophages *ex vivo*. *Int J Vitam Nutr Res* 83:162–167
- Zaychenko SG, Zernov AS (2017) Structural features of the seed coat in Caucasian representatives of *Minuartia* (Caryophyllaceae). *Wulfenia J* 24:205–220

- Zdraveva P, Gevrenova R, Dimitrova B (2004) Phenolic compounds of *Scleranthus annuus* L. (Caryophyllaceae). 3rd Conference on Medicinal and Aromatic Plants of Southeast European Countries. Nitra, Slovakia Republic 2004:57
- Zdraveva P, Pencheva I, Popova P, Ionkova I, Krasteva I (2015) Production of saponarin in *in vitro* cultures of *Gypsophila species*. J Chem Pharm Res 7:829–832
- Zemtsova GN, Glyzin VY, Dzhumyrko SF (1976) Flavones and their C-glycosides from *Silene saxatilis*. Chem Nat Compd 11:538
- Zhang H (2012) Profiling analysis of the seeds of *Vaccaria segetalis* (Necr.) Gracke by HPLC-ESI-MS. Adv Mater Res 396–398:96–98
- Zhang FM, Tai ZG, Cai L, Yang YB, Li F, Ding ZT (2011) Flavonoids from *Gypsophila elegans* and their antioxidant activities. J Yunnan Univ (Nat Sci Ed) 33:93–95
- Zhang H, Wang K, Wu J, Chen Y, He P (2011) A new flavonoid glycoside from *Vaccaria hispanica*. Nat Prod Commun 6:1599–1602
- Zhang Y, Wang G, Lv H, Luo J, Kong L (2015) Two new β -carboline alkaloids from the roots of *Gypsophila oldhamiana*. Nat Prod Res 29:1207–1211
- Zheleva-Dimitrova D, Zengin G, Balabanova V, Voynikov Y, Lozanov V, Lazarova I, Gevrenova R (2018) Chemical characterization with *in vitro* biological activities of *Gypsophila* species. J Pharm Biomed Anal 155:56–69
- Zhou X, Wang L, Tian Y, Gong X, Zhao C, Yang S (2013) Chemical constituents from roots of *Psammossilene tunioides*. Zhongguo Zhong Yao Za Zhi 38:3507–3509
- Zhou G, Tang L, Wang T, Zhou X, Kou Z, Wu J, Wang Z (2016) Phytochemistry and pharmacological activities of *Vaccaria hispanica* (Miller) Rauschert: a review. Phytochem Rev 15:813–827
- Zhou G, Wu H, Wang T, Guo R, Xu J, Zhang Q, Wang Z (2017) C-glycosylflavone with rotational isomers from *Vaccaria hispanica* (Miller) Rauschert seeds. Phytochem Lett 19:241–247
- Zhu G, Liu X, Li H, Yan Y, Hong X, Lin Z (2018) Kaempferol inhibits proliferation, migration, and invasion of liver cancer HepG2 cells by down-regulation of microRNA-21. Int J Immunopathol Pharmacol 32:1–12
- Zhuang L (1983) C-Glycosylflavones from Qi Gu Cao (*Sagina japonica*). Zhongcaoyao 14:295–297
- Zitouni M (2017) Profil polyphénolique et activité antioxydante de deux plantes médicinales *Pistacia lentiscus*. L et *Gymnocarpos decander* Forsk. Université Abou Bekr Belkaid, Tlemcen
- Zoll A, Nouvel G (1974) Comparative study of C-glycosyl flavones of two Caryophyllaceae. *Spergularia rubra* and *Stellaria holostea*. Phytochemistry 8:134–140

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.