



# Correction: Ectopic overexpression of *TaHsfA5* promotes thermomorphogenesis in *Arabidopsis thaliana* and thermotolerance in *Oryza sativa*

Harsha Samtani<sup>1</sup> · Aishwarye Sharma<sup>1</sup> · Paramjit Khurana<sup>1</sup>

Published online: 27 June 2023

© The Author(s), under exclusive licence to Springer Nature B.V. 2023

## Correction to: Plant Molecular Biology

<https://doi.org/10.1007/s11103-023-01355-3>

In the original publication, Fig. 5 was a duplicate of Fig. 4. Both Figs. 4 and 5 should have appeared as shown below and the original article has been corrected.


---

The original article can be found online at <https://doi.org/10.1007/s11103-023-01355-3>.

---

✉ Paramjit Khurana  
[param@genomeindia.org](mailto:param@genomeindia.org)

<sup>1</sup> Department of Plant Molecular Biology, University of Delhi South Campus, Benito Juarez Road, New Delhi 110021, India



**Fig. 4** Interaction of TaHsfA5 with TaHsfA4, TaHsfA3, TaHsfA2 and TaHSBP2. The Yeast-2-hybrid assay was performed to check the interaction of TaHsfA5 with TaHsfA4, TaHsfA3, TaHsfA2, TaHSBP1, and TaHSBP2 proteins. Growth of co-transformed yeast cells was analyzed on SD/-Leu/-Trp (-LW) medium and SD/-Leu/-

Trp/-His (-HLW) medium. Drop assay was performed on media lacking Histidine, Leucine and Tryptophan (-HLW) along with 0.5 mM 3-aminotriazole (3AT). Pair of plasmids pGBKT7-53 and pGADT7-T were used as positive control, and pGBKT7-Lam and pGADT7-T were used as the negative control

**Fig. 5** BiFC assay showing the interaction of TaHsfA5 with TaHsfA4, TaHsfA3 and TaHSBP2 in the nucleus and cytoplasm in onion epidermal cells. Interaction between TaHsfA5 and TaHsfA2 was used as a negative control

