CORRECTION

Correction to: Quantification of the contrasting root systems of *Pinus thunbergii* in soils with different groundwater levels in a coastal forest in Japan

Yasuhiro Hirano · Chikage Todo · Keitaro Yamase ·
Toko Tanikawa · Masako Dannoura · Mizue Ohashi ·
Ryuusei Doi · Ryusei Wada · Hidetoshi Ikeno

Published online: 2 May 2018
© Springer International Publishing AG, part of Springer Nature 2018

Correction to: Plant Soil

<https://doi.org/10.1007/s11104-018-3630-9>

The original version of this article unfortunately contained a mistake. Table 3 was published erroneously. This Table has now been corrected.

The online version of the original article can be found at
<https://doi.org/10.1007/s11104-018-3630-9>

Y. Hirano (✉) · R. Doi · R. Wada
Graduate School of Environmental Studies, Nagoya University,
Nagoya 464-8601, Japan
e-mail: yhirano@nagoya-u.jp

C. Todo · K. Yamase
Hyogo Prefectural Technology Center for Agriculture, Forestry
and Fisheries, Shiso 671-2515, Japan

T. Tanikawa
Kansai Research Center, Forestry and Forest Products Research
Institute, Kyoto 612-0855, Japan

M. Dannoura
Graduate School of Global Environmental Studies, Kyoto
University, Kyoto 606-8501, Japan

M. Dannoura
Graduate School of Agriculture, Kyoto University,
Kyoto 606-8502, Japan

M. Ohashi · H. Ikeno
School of Human Science and Environment, University of Hyogo,
Himeji, Hyogo 670-0092, Japan

Table 3 Belowground traits of the harvested *Pinus thunbergii* trees in the sea- and land-side plots (Mean $\pm$ SE, $n = 3$)

	Sea side	Land side
Maximum depth of roots at 2 cm in diameter (cm)	56 $\pm$ 12	212 $\pm$ 19 ^{**}
Maximum length of horizontal roots at 3 cm in diameter (cm)	655 $\pm$ 13	450 $\pm$ 95 ^{ns}
Maximum length of horizontal roots to maximum root depth (cm cm ⁻¹)	13.0 $\pm$ 3.2	2.1 $\pm$ 0.3 [*]
H to maximum root depth (m m ⁻¹)	12.0 $\pm$ 0.9	5.4 $\pm$ 0.2 ^{**}
H to maximum length of horizontal roots (m m ⁻¹)	1.0 $\pm$ 0.2	2.6 $\pm$ 0.3 ^{**}
Number of horizontal roots at 100 cm distance from stem (no.)	29 $\pm$ 4	23 $\pm$ 8 ^{ns}
Number of horizontal roots at 150 cm distance from stem (no.)	37 $\pm$ 12	16 $\pm$ 3 ^{ns}
Sum of cross sectional area of horizontal roots at 100 cm distance from stem (cm ²)	535 $\pm$ 155	264 $\pm$ 106 ^{ns}
Sum of cross sectional area of horizontal roots at 150 cm distance from stem (cm ²)	435 $\pm$ 156	129 $\pm$ 39 ^{ns}
Number of vertical roots at 50 cm depth (no.)	14 $\pm$ 9	8 $\pm$ 2 ^{ns}
Sum of cross sectional area of vertical roots at 50 cm depth (cm ²)	66 $\pm$ 60	848 $\pm$ 497 ^{ns}
Dry weight of roots at less than 50 cm depth (kg dw)	102 $\pm$ 29	49 $\pm$ 9 ^{ns}
Dry weight of roots at deeper than 50 cm depth (kg dw)	0.8 $\pm$ 0.7	25 $\pm$ 9 ^{ns}
Dry weight of smaller roots <1 cm in diameter (kg dw)	1.0 $\pm$ 0.3	1.2 $\pm$ 0.6 ^{ns}
Dry weight of the whole roots (kg dw)	104 $\pm$ 28	75 $\pm$ 14 ^{ns}
Root to shoot ratio (kg dw kg dw ⁻¹)	0.81 $\pm$ 0.07	0.57 $\pm$ 0.02 ^{**}

H: stem height,

Significant differences between the sea- and land-side plots at ^{**} $p < 0.01$, ^{*} $p < 0.05$,

^{ns}: not significant