



Correction to: Below-ground nitrogen transfer from oak seedlings facilitates *Molinia* growth: ^{15}N pulse-chase labelling

Marine Fernandez · Philippe Malagoli · Antoine Vernay · Thierry Améglio · Philippe Balandier

Published online: 30 April 2020
© Springer Nature Switzerland AG 2020

Correction to: Plant and Soil.

<https://doi.org/10.1007/s11104-020-04473-9>

In the published version of this paper, the titles for figs. 3 and 4 should be presented as below:

Fig. 3 ^{15}N allocation (expressed as % of total ^{15}N measured in all compartments) among shoots, roots and soil in donor oak and receiver *Molinia*. Species

were sole-grown (a, b, c and d) or mixed-grown (e, f, g and h), for each season: summer (a and e), autumn (b and f), winter (c and g) and spring (d and h). Values are reported as means \pm SE ($n = 4$)

Fig. 4 ^{15}N allocation (expressed as % of total ^{15}N measured in all compartments) among shoots, roots and soil in donor oak and receiver oak. Species were sole-grown (a, b, c and d) or mixed-grown (e, f, g and h), for each season: summer (a and e), autumn (b and f), winter (c and g) and spring (d and h). Values are reported as means \pm SE ($n = 4$)

The online version of the original article can be found at
<https://doi.org/10.1007/s11104-020-04473-9>

M. Fernandez · P. Malagoli (✉) · T. Améglio
Université Clermont Auvergne, INRAE, PIAF,
F-63000 Clermont-Ferrand, France
e-mail: philippe.malagoli@uca.fr

M. Fernandez
e-mail: marine.fernandez@uca.fr

T. Améglio
e-mail: thierry.ameglio@inrae.fr

A. Vernay
Department of Forest Ecology and Management, Swedish,
University of Agricultural Sciences (SLU), SE-901 83 Umeå,
Sweden
e-mail: antoine.vernay@slu.se

P. Balandier
INRAE, UR EFNO, Domaine des Barres,
F-45290 Nogent-sur-Vernisson, France
e-mail: philippe.balandier@inrae.fr

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.