



# Influence of small-scale spatial variability of soil properties on yield formation of winter wheat

Jonas Groß · Norman Gentsch · Jens Boy · Diana Heuermann ·  
Dörte Schweneker · Ulf Feuerstein · Johannes Brunner ·  
Nicolaus von Wirén · Georg Guggenberger · Bernhard Bauer

Received: 5 April 2023 / Accepted: 2 August 2023 / Published online: 20 September 2023  
© The Author(s) 2023

## Abstract

**Background** With the increasing development of sophisticated precision farming techniques, high-resolution application maps are frequently discussed as a key factor in increasing yield potential. However, yield potential maps based on multiple soil properties measurements are rarely part of current farming practices. Furthermore, small-scale differences in soil properties have not been taken into account.

**Methods** To investigate the impact of soil property changes at high resolution on yield, a field trial has been divided into a sampling grid of 42 plots. The soil properties in each plot were determined at three soil depths. Grain yield and yield formation of winter wheat were analyzed at two sites.

**Results** Multiple regression analyses of soil properties with yield measures showed that the soil contents of organic carbon, silt, and clay in the top and subsoil explained 45–46% of the variability in grain yield. However, an increasing clay content in the topsoil correlated positively with grain yield and tiller density. In contrast, a higher clay content in the subsoil led to a decrease in grain yield. A cluster analysis of soil texture was deployed to evaluate whether the soil's small-scale differences caused crucial differences in yield formation. Significant differences in soil organic carbon, yield, and yield formation were observed among clusters in each soil depth.

**Conclusion** These results show that small-scale lateral and vertical differences in soil properties can strongly impact crop yields and should be considered to improve site-specific cropping techniques further.

---

Responsible Editor: Rémi Cardinael.

---

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1007/s11104-023-06212-2>.

---

J. Groß · J. Brunner · B. Bauer (✉)  
Crop Production and Crop Protection, Institute of Biomass Research, University of Applied Sciences Weihenstephan-Triesdorf, Markgrafenstraße 16, 91746 Weidenbach, Germany  
e-mail: [bernhard.bauer@hswt.de](mailto:bernhard.bauer@hswt.de)

J. Groß  
Kuratorium für Technik und Bauwesen in der Landwirtschaft e.V. (KTBL), Bartningstraße 48, 24689 Darmstadt, Germany

**Keywords** Soil texture · Soil organic carbon · Lateral and vertical small-scale variability · *Triticum aestivum* L. · Yield formation · Precision farming

N. Gentsch · J. Boy · G. Guggenberger  
Institute of Soil Science, Leibniz Universität Hannover, Herrenhäuser Straße 2, 30419 Hannover, Germany

D. Heuermann · N. von Wirén  
Molecular Plant Nutrition, Leibniz Institute of Plant Genetics and Crop Plant Research Gatersleben, Corrensstraße 3, 06466 Stadt Seeland, Germany

D. Schweneker · U. Feuerstein  
Deutsche Saatveredelung AG, Steimker Weg 7, 27330 Asendorf, Germany

## Introduction

The impact of global warming, environmental constraints, and increasing demand for food due to a growing world population requires an urgent improvement of crop management, including efficient use of farming resources such as fertilizers and pesticides, while increasing crop yield at the same time (Cassmann 1999; Foley et al. 2011; Gebbers and Adamchuk 2010; Katalin et al. 2014; Kühn et al. 2009; Machado et al. 2000; Mulla 2013; Schimmelpfennig 2018). Besides crop management, the yield formation of crops is mainly driven by abiotic factors such as climate and soil properties. Especially at the field level, the impact of climatic factors on crop growth can be seen as homogeneous. At the same time, small-scale differences in soil properties may provoke heterogeneity in yield formation within a uniformly managed crop stand.

Soil texture, as a fundamental master variable in soils (Ad hoc AG Boden 2005), mainly influences the available water capacity in the effective root space and the air capacity. These properties are directly linked to the crop's nutrient and water availability (Ad hoc AG Boden 2005; Amelung et al. 2018; Maidl et al. 1999). At most locations, soil texture changes in the landscape and with soil depth, influencing root development, water availability, and soil aeration depending on the geomorphology and the parent material for soil formation (Barraclough and Leigh 1984; Breuning Madsen 1985; Poehlau and Kätterer 2017; Schulte-Eickholt 2009; White and Kirkegaard 2010). Compared to the topsoil, the subsoil's nutrient content is mostly substantially lower in arable fields if there are no significant changes in soil texture (Crist and Weaver 1924; Kautz et al. 2013). Nevertheless, the subsoil is an essential resource for macro- and micronutrient mining by plant roots in nutrient depletion in the topsoil or topsoil drought (Kautz et al. 2013). A gradient in the nutrient content from the top to the subsoil is mainly driven by plant litter decomposition, surface fertilizer application, topsoil cultivation, and nutrient leakage or dislocation.

In contrast to texture, the soil organic carbon (SOC) content can be influenced by management practices from a medium to long-term perspective (Freibauer et al. 2004; Poehlau et al. 2020, 2021). In the topsoil, SOC can be increased by crop management, for example, through the use of organic fertilizers and the incorporation of plant residues (Guo

et al. 2019; Wuest and Gollany 2013) by including soil organic matter (SOM) preserving crops into crop rotations (Börjesson et al. 2018; Osanai et al. 2021; Šeremešić et al. 2020), or the cultivation of cover crops (Liang et al. 2022; Poehlau et al. 2021). SOC is related to higher water- and nutrient- availability for plants (Ghaley et al. 2018) and is known to have beneficial effects on soil properties, like soil aggregate stability, bulk density, aeration, and soil biological processes, which all affect plant growth (Ghaley et al. 2018; Palmer et al. 2017; Šeremešić et al. 2020; Stein et al. 1997; Yost and Hartemink 2019). However, SOC is not distributed homogeneously either within soil profiles or across fields (Hbirkou et al. 2012; Rentschler et al. 2020), and crop production has the potential even to increase site-specific variations (Conant et al. 2011; Virto et al. 2012).

Due to its high economic value, winter wheat is one of the most important cereals cultivated in temperate regions of Europe (Eurostat 2021). Nevertheless, multiple environmental factors influence individual yield components during different stages of their development. This high responsiveness makes it suitable as a model crop for determining small-scale soil differences. The yield components, plant density, tiller density, spike density, number of grains per spike, and thousand-grain weight, form the winter wheat grain yield. However, the yield components are strongly interrelated and plastic (Slafer and Rawson 1994; Slafer 2003; Roth et al. 1984) so that they can compensate for each other (Bastos et al. 2020).

Under heterogeneous soil conditions in a field, differences in grain yield can be large (Camarano et al. 2019; Buttafuoco et al. 2017; Taylor et al. 2003). Within a small distance of < 10 m, the variation in winter wheat grain yield can be significant (Crain et al. 2013). Data from several studies suggest that the variability in grain yield is mainly linked to the variability of soil properties (Bölenius et al. 2017; Cox et al. 2003; Gozdowski et al. 2017; Maidl et al. 1999; Rodriguez-Moreno et al. 2014; Wang and Shen 2015). When subdividing a field into management zones, precision agriculture improves the efficient use of pesticides and fertilizers and can help to homogenize the nutrient availability and growth conditions in a field (Argento et al. 2021; Hørfarter et al. 2019; Katalin et al. 2014; Maidl et al. 1999; Nordmeyer 2006; Patzold et al. 2008; Stewart et al. 2002; Wagner and Marz 2017). Site-specific application techniques for sowing, crop protection, N fertilization (Barnes et al.

2019), and irrigation have already been implemented in practical crop production to a certain extent. Especially under highly heterogeneous soil conditions, high-resolution mapping of soil properties is likely to increase the profitability of, for example, site-specific N management compared to coarser mapping approaches (Denora et al. 2022; Späti et al. 2021). However, most field experiments investigating the impact of spatial variability of soil properties on winter wheat yield were performed on larger grid sizes > 0.5 ha (LaRuffa et al. 2001; Mallarino et al. 1999).

Site-specific management at higher resolution can improve resource efficiency and become increasingly crucial for future farming practices (Boenecke et al. 2018; Brogi et al. 2021; LaRuffa et al. 2001; Skakun et al. 2021; Tubaña et al. 2008; Zebarth et al. 2021). In this context, spatial variation in soil properties in combination with soil depth is often not considered (Boenecke et al. 2018; Cammarano et al. 2019). Below 30 cm, variation in soil properties becomes crucial at later development stages of cereals (Kirkegaard et al. 2007). For example, Cammarano et al. (2019) showed that higher clay content in the subsoil led to low and unstable spring barley yields. Steadily evolving high-precision application techniques enhance the resolution of site-specific management. These more precise application techniques will only benefit site-specific management if the field's lateral and vertical resolution of specific management zones increases. However, site-specific management often considers only two-dimensional lateral differences in soil properties.

This study aims to evaluate the impact of small-scale vertical and lateral soil heterogeneity in two typical locations in Germany on the yield formation of winter wheat. Therefore, we (i) assessed the lateral and vertical small-scale variability of soil properties, (ii) analyzed the influence of soil properties in different soil depths on winter wheat grain yield and its components at a small-scale level, and (iii) identified clusters of soil texture and their influence on yield components and grain yield.

## Materials and methods

### Locations

The field experiment was conducted from autumn 2015 to summer 2016 at two locations in Germany:

Triesdorf is located in Northern Bavaria (450 m a.s.l., 49°12'36.5"N 10°38'33.9"E) on a Stagnic Cambisol (IUSS Working Group WRB 2015). The area with a mean annual air temperature of 8.7 °C and annual precipitation of 674 mm (2005–2015) is characterized by a warm, fully humid climate with warm summer (Kottek et al. 2006). The second location is Asendorf in Ems-Hunte-Geest (Lower Saxony, 49 m a.s.l., 52°45'48.4"N 9°01'24.3"E). The climatic conditions are similarly warm and humid, with a mean annual air temperature of 8.9 °C and annual precipitation of 705 mm (2005–2015) (Kottek et al. 2006). The soil is also described as a Stagnic Cambisol (IUSS Working Group WRB 2015).

### Experimental design

At both locations, a subarea of a field was selected for the field trial with a size of 2 areas of 63.0 m x 34.5 m separated by a tramline and divided into a sampling grid of 42 squares to assess the effects of spatial variability of soil properties on crop yield and the structure of its components. The plot size was 10.5 m x 9.0 m each.

### Plant material and growth conditions

On both sites, the field experiment started with the harvest of the previous crop, spring wheat, in 2015. Tillage and seedbed preparation was performed with conservation tillage. On 13th October 2015 in Triesdorf and on 21st October 2015 in Asendorf, winter wheat (cultivar "Patras") was sown with a sowing rate of 345 grains per m<sup>2</sup>. During the vegetation period, crop management followed common local practices adapted to the two regions. The crop protection strategy was based on a homogeneous growth regulator application at the beginning of stem elongation and a fungicide application at the flag leaf stage. In total, 220 kg ha<sup>-1</sup> in Triesdorf and 160 kg ha<sup>-1</sup> in Asendorf of nitrogen were applied according to the different yield potentials of the locations. The fertilizer was applied in three partial doses during winter wheat's tillering, stem elongation, and grain filling phase.

## Analysis of soil measures

Soil samples for bulk density were taken manually from an open pit in depths of 0–10, 10–30, and 30–60 cm before sowing in autumn 2015 at both locations. The soil texture was analyzed according to DIN ISO 11,277. The pH value was quantified in soil suspension with water at a solid-to-solution ratio of 1:2.5, with 10 g of dry soil in 25 ml H<sub>2</sub>O. A stainless-steel cutter (100 cm<sup>3</sup>) was used for sampling bulk density. Samples were dried at 105 °C, and bulk density was determined gravimetrically (Rawls 1983). SOC content was measured with an EA system (Elementar Analysensysteme GmbH, Hanau, Germany). Traces of carbonates (<0.5%) were removed by acid fumigation, according to Harris et al. (2001), and subsequent neutralization was carried out over NaOH pellets, 48 h for each sample.

## Analysis of plant measures

The plant development of winter wheat was monitored in each plot over the growing period following the BBCH code (Hess et al. 1997). Stand density was assessed by counting the number of tillers per m<sup>2</sup> at the beginning of stem elongation (BBCH 31). Spike density was determined before harvest (BBCH 79) based on one square meter per plot. Winter wheat was harvested on 4th August 2016 in Triesdorf and 27th September 2016 in Asendorf. Plots were harvested with the plot combine Hege 140 (Hege Saatzuchtmaschinen, Hohebuch, Germany) for the exact quantification of grain yield. Only the score area of the plots was harvested to minimize border effects. The total plot area was divided into three repetitions of 15.75 m<sup>2</sup> each, so yield components could be determined in three replicates per plot. Grain samples were dried for 24 h at 65 °C to determine grain dry matter yield gravimetrically. Postharvest, the thousand-grain weight was determined by counting 500 grains per sample and quantified weight gravimetrically based on 14.0% moisture (Pfeuffer Contador, Germany). Additionally, grain protein content was determined with near-infrared spectroscopy using the DA 7250 At-line NIR Instrument (PerkinElmer, Waltham, United States).


## Statistical analysis

In the first step, data were analyzed with descriptive statistics like minimum and maximum values, mean, standard deviation (SD), and coefficient of variation (CV), and the dataset was analyzed for distribution. Statistical analysis was performed using the R program (R Core Team 2021). Variations of soil texture of all soil layers were visualized in the soil triangle according to the German soil classification system (Ad hoc AG Boden 2005) using the R package “soil texture” (Moey 2018). Further, as a measure of linear correlation, Pearson’s correlation coefficients ( $r$ ) between soil measures and plant measures were calculated, which is recommended for sampling numbers <50 points (Leroux and Tisseyre 2019). To investigate soil properties’ influence on winter wheat yield and its yield components, we performed a stepwise multiple regression analysis using the Akaike information criterion (AIC) as a selection criterion for removing or adding variables of soil properties against the plant properties (Bozdogan 1987). Distributions of the experimental errors were tested for normality by executing Shapiro-Wilk’s test for each regression.

Variance inflation factor (VIF) was used to test multiple regression models on collinearity (Manly and Navarro Alberto 2004; Miles 2014). To visualize the influence of the soil texture (especially for silt and clay) in different soil depths on the development of yield and yield components, the R-package “Akima” (Version 0.6–2.1) was used to create linear interpolated contour plots (Akima et al. 2020).

An additional approach for data analysis was clustering the dataset. Clustering by soil properties is common practice in site-specific farming approaches and is done by zoning (Taylor et al. 2003). Soil texture is a constant physical soil measure that barely changes over time. To distinguish subsets in the data set characterized by minimized intra-cluster variation, a k-means cluster analysis was performed (Hartigan and Wong 1979). Based on this analysis, clusters have been identified, mainly based on the measures of sand and clay (Fig. 1). For comparison of the mean values of these clusters, a two-way analysis of variance (ANOVA) was conducted. To check the basic assumptions for an ANOVA, the homogeneity of variances was tested using the Levene test (Schultz 1985).

**Fig. 1** Schema of clustering the dataset by measures of the soil texture in Triesdorf



The normal distribution of the residuals was verified using the Shapiro-Wilk test. When needed, data were transformed (log8 or exponential), and a one-way analysis of means (not assuming equal variances) was executed (Welch 1951), followed by a multi-comparison of means with Tukey’s honestly significant difference (HSD) test with  $p < 0.05$ . A multiple comparison of means calculating the least significant difference (LSD) on all measures with  $p < 0.05$  was calculated using the R-package “Agricolae” (Mendiburu 2021).

## Results

### Lateral and vertical variability of soil properties


Small-scale lateral variability of soil properties was analyzed using a grid of 10.5 m x 9.0 m plots at two locations. Samples were also taken in three soil depths to define the vertical variability. Firstly a plot-wise classification of the prevailing soil type according to Ad hoc AG Boden (2005) was performed to get an idea of soil variability at both locations (Fig. 2a). The observed soil types in Triesdorf ranged from medium loamy sand (SI3), strong loamy sand (SI4) over medium sandy loam (Ls3) to strong sandy loam (Ls4) in the topsoil containing two upper analyzed soil layers (0–10 cm and 10–30 cm). In the subsoil (30–60 cm), the soil type SI3 was present in 19% of the plots, while SI4 characterized 33%, Ls 24%, Ls 19%, partly silty loamy sand (Slu) only 2%, an sandy clay loam (Lts) 2% of the plots in Triesdorf (Fig. 2b). So, clay content showed a higher coefficient of variation (CV) in each soil layer compared to the fractions of silt and sand. The values range from 12.5 to 23.9%, with the highest CV in the subsoil layer (30–60 cm) (Table S1).

At the second location Asendorf, the soil was generally characterized on average by 12% points smaller clay and 43% points higher silt contents than at Triesdorf. So, the observed soil types are sandy silt (Us), pure silt (Uu), sandy clayey silt (Uls), and slight clayey silt (Ut2) (Fig. 2a). Also, the soil texture’s variability was low in the subsoil, with 83.3% of the soil samples being classified as sandy silt (Us) (Fig. 2c). Hence, the two locations showing different soil types, and Asendorf delivered more homogenous conditions in small-scale variability for the field trials compared to Triesdorf.

As expected, the SOC content decreased from the topsoil to the subsoil. In Triesdorf, on average, 14.5 mg g<sup>-1</sup> SOC was observed in the first soil layer of 0–10 cm, 12.4 mg g<sup>-1</sup> SOC in 10–30 cm soil depth, and 6.5 mg g<sup>-1</sup> SOC in the subsoil (30–60 cm) (Table 1). A similar gradient in SOC over soil depth was observed in Asendorf too. Nevertheless, in contrast to bulk density, high lateral variability of SOC content was observed in all three soil layers, indicated by CV values ranging from 16.1 to 32.9% at both sites. Bulk density increased with soil depth with a mean of 1.3 g cm<sup>-3</sup> in the topsoil, 1.5 g cm<sup>-3</sup> in 10–30 cm, and 1.7 g cm<sup>-3</sup> in the subsoil, and minor variations of the pH value were observed in Triesdorf. Similar vertical variations of the measured soil properties could be observed in the soil profile of the second location Asendorf (Table 1). Nevertheless, compared to soil texture, the lateral variability in SOC and bulk density were nearly comparable in Triesdorf and Asendorf (Table 1).

### Variability of grain yield and yield components

In order to characterize the impact of the presented small-scale lateral and vertical soil variability on yield formation, winter wheat was cultivated, and



**Fig. 2** Soil texture in the soil layers of 0–10 cm, 10–30 cm, and 30–60 cm (a) and the number of soil samples that could be assigned to individual soil texture classes [n] to soil depths

in Triesdorf (b) and Asendorf (c), following the German soil classification system (Ad hoc AG Boden 2005)

yield components were analyzed in each plot. A sowing rate of 345 grains  $m^{-2}$  was applied homogeneously over all plots, and in the field emergence rate (72.8%), no statistically significant differences were observed. Wheat plants tillered differently in the plots during the vegetative development (until BBCH 31), so a variation of 367 to 591 tillers per  $m^2$  with a CV of 10.8% was observed in Triesdorf (Table 2). The variation of tiller density between plots decreased even during the tiller reduction phase, resulting in 388 and 538 spikes  $m^{-2}$  and a CV of 7.5% at BBCH 79. Lateral differences between plots were not only observed in the spike density. Also, the number of grains per spike and the single spike weight showed a CV of 10.7% and 9.5%, respectively. While the measures representing organ setting were strongly influenced by soil's inhomogeneity, like spike density and numbers of grains per spike, resulting in a high grain density range from 17,313 to 22,107 grains  $m^{-2}$ , grain weight was not strongly influenced. In Triesdorf, the thousand-grain weight ranged from 48.7 to 57.3 g, and the soil's small-scale differences caused a

CV of 3.5%. Likewise, the CV of 4.1% was relatively low for the grain yield. Nevertheless, the mean grain yield was 10.4  $t ha^{-1}$  ranging from 9.6 to 11.2  $t ha^{-1}$ , and soil's small-scale effects on grain protein content with values between 12.9 and 13.8% and a CV of only 1.6% were low.

In Asendorf, vegetative development was much more substantial and led to more tillers. On average, 816 tillers per  $m^2$  were observed at the end of tillering period (BBCH 31), and the CV of 3.7% was low (Table 2). However, tiller reduction during booting led to a mean of 543 spikes per  $m^2$ , and variability between plots increased to a CV of 7.7% before harvest. Due to elevated spike density, the crop stand was characterized by a mean number of 31.1 grains per spike and 16,820 grains per  $m^2$ . Hence, the grain setting was much lower in comparison to Triesdorf. Even though fewer grains were created per  $m^{-2}$ , grain weight was relatively low in Asendorf compared to Triesdorf. Hence, thousand-grain weight ranged from 28.1 to 44.2 g with a CV of 7.2%, indicating that small-scale differences in soil properties influenced

**Table 1** Descriptive statistics of soil properties SOC, bulk density, and pH-value according to different soil depths at the field sites in Triesdorf and Asendorf in 2016, including minimum, maximum, mean, standard deviation, and the coefficient of variation in % (CV);  $n=42$  for each location

Soil depth [cm]	Soil properties	Minimum	Maximum	Mean	Standard deviation	CV [%]
Triesdorf						
0–10	Soil organic carbon [mg g <sup>-1</sup> ]	4.9	19.2	14.5	3.5	23.7
	Bulk density [g cm <sup>-3</sup> ]	1.0	1.6	1.3	0.1	10.3
	pH value	6.8	7.9	7.5	0.4	
10–30	Soil organic carbon [mg g <sup>-1</sup> ]	7.8	17.1	12.4	2.3	18.2
	Bulk density [g cm <sup>-3</sup> ]	1.3	1.8	1.5	0.1	7.0
	pH value	6.8	7.9	7.5	0.3	
30–60	Soil organic carbon [mg g <sup>-1</sup> ]	3.8	9.1	6.5	1.3	20.2
	Bulk density [g cm <sup>-3</sup> ]	1.5	1.9	1.7	0.1	4.3
	pH value	7.0	7.9	7.5	0.2	
Asendorf						
0–10	Soil organic carbon [mg g <sup>-1</sup> ]	8.8	23.5	18.0	3.0	16.4
	Bulk density [g cm <sup>-3</sup> ]	1.1	1.3	1.2	0.1	4.4
	pH value	5.3	6.6	6.1	0.3	
10–30	Soil organic carbon [mg g <sup>-1</sup> ]	13.8	24.9	17.9	2.9	16.1
	Bulk density [g cm <sup>-3</sup> ]	1.2	1.4	1.3	0.1	4.3
	pH value	5.5	6.8	6.2	0.3	
30–60	Soil organic carbon [mg g <sup>-1</sup> ]	4.4	20.8	9.8	3.2	32.9
	Bulk density [g cm <sup>-3</sup> ]	1.4	1.8	1.7	0.1	6.6
	pH value	5.9	6.9	6.4	0.2	

**Table 2** Descriptive statistics of the yield components and measures tiller density, spike density, grain per spike, single spike weight, grain density, thousand-grain weight, grain yield, and grain protein content in Triesdorf and Asendorf in 2016, including minimum, maximum, mean, standard deviation, and the coefficient of variation [%],  $n=42$  for each location

Plant properties	Minimum	Maximum	Mean	SD	CV [%]
Triesdorf					
Tillers density [tillers m <sup>-2</sup> ]	367	591	485	52.84	10.8
Spikes density [spike m <sup>-2</sup> ]	388	538	468	35.38	7.5
Grains per spike [n]	34	53	42	4.6	10.7
Single spike weight [g]	1.87	2.80	2.24	0.22	9.5
Grain density [grains m <sup>-2</sup> ]	17,313	22,107	19,608	1,130	5.7
Thousand-grain weight [1000 grains g <sup>-1</sup> ]	48.7	57.3	53.3	1.9	3.5
Grain yield [t ha <sup>-1</sup> ]	9.6	11.2	10.4	0.43	4.1
Protein content [%]	12.9	13.8	13.4	0.2	1.6
Asendorf					
Tillers density [tillers m <sup>-2</sup> ]	725	885	816	31	3.7
Spikes density [spike m <sup>-2</sup> ]	461	676	543	41	7.7
Grains per spike	25	43	31	2.9	9.4
Single spike weight [g]	1.00	1.50	1.23	0.11	9.1
Grain density [grains m <sup>-2</sup> ]	14,894	22,753	16,820	1,228	7.3
Thousand-grain weight [1000 grains g <sup>-1</sup> ]	28.1	44.2	39.6	2.8	7.2
Grain yield [t ha <sup>-1</sup> ]	5.8	7.2	6.6	0.31	4.8
Protein content [%]	10.9	12.3	11.6	0.3	2.4

yield formation much stronger in the later stages compared to the vegetative growth in this location. As a result, grain yield ranged from 5.8 to 7.2 t ha<sup>-1</sup> with a mean yield of 6.6 t ha<sup>-1</sup>. Even though the lower yield level in Asendorf, the grain protein content was only 11.6%, with a small CV of 2.4%.

In summary, the two sites' different growing conditions and soil properties had contrasting impacts on plant development. While vegetative development was retarded and showed high variations due to soil properties in Triesdorf, the later-formed yield components compensate for the limited vegetative development. Thus, the influence of soil differences became less important at later developmental stages. In contrast, the solid vegetative development was hardly influenced by soil differences in Asendorf. However, the impact of soil properties increased during the generative developmental stages but could not be associated with plant properties like in Triesdorf, showing higher soil heterogeneity (Fig. S2).

#### Influence of soil properties on yield formation

To further assess the relationship between plant measures and soil properties, Pearson's correlation coefficients ( $r$ ) were calculated. In Triesdorf, the soil's small-scale variability was much more pronounced than in Asendorf. Grain yield correlated positively with clay and silt in the topsoil (0–30 cm). The correlation coefficient ranged from 0.32 to 0.39,  $p < 0.05$  (Table 3). In contrast, the correlation with

sand showed negative values in all three tested soil layers. Interestingly, clay content was negatively associated with grain yield in the subsoil ( $r = -0.31$ ,  $p < 0.05$ ). While thousand-grain weight showed no significant correlation with any measure of the soil texture, tillering showed a clear positive association with clay content ( $r = 0.44$ ,  $p < 0.01$ ) and a negative with sand ( $r = -0.36$ ,  $p < 0.05$  and  $-0.44$ ,  $p < 0.01$ ) in the topsoil down to 30 cm. Nevertheless, spike density strongly based on tillering showed no significant association with any measure of the soil texture in the first tested layer (0–10 cm). In the second layer of the topsoil (10–30 cm), the correlation for clay ( $r = 0.34$ ,  $p < 0.05$ ) was significantly positive, while sand ( $r = -0.31$ ,  $p < 0.05$ ) was negative. No significant correlations were observed in the subsoil. Grain protein content was only significantly correlated with silt content ( $r = 0.32$ ,  $p < 0.05$ ) and clay content ( $r = -0.42$ ,  $p < 0.01$ ) in the subsoil, indicating the importance of the subsoil during grain filling, even if this was not observable for the thousand-grain weight.

In summary, wheat yield formation in Triesdorf profited from higher clay contents in the topsoil, while larger sand contents were counterproductive to form yield components. In contrast, the subsoil silt content was the most decisive for yield formation.

A stepwise multiple regression analysis was performed to understand the functional relationship between grain yield and the yield components as the dependent and the soil properties as the independent variables. Additionally, soil measures were expanded to include SOC and bulk density.

**Table 3** Pearson's correlation coefficient ( $r$ ) and  $p$ -value ( $p$ ) of soil texture divided into clay (0–2  $\mu\text{m}$ ), silt (2–63  $\mu\text{m}$ ), and sand (63–200  $\mu\text{m}$ ), as well as the plant measures tiller density, spike density, grain yield, thousand-grain weight, and grain protein content in Triesdorf, 2016

Soil depth [cm]	Soil texture	tiller density [tillers m <sup>-2</sup> ]		spike density [spike m <sup>-2</sup> ]		grain yield [t ha <sup>-1</sup> ]		thousand-grain weight [1000 grains g <sup>-1</sup> ]		protein content [%]	
		r <sup>2</sup>	$p$	r <sup>2</sup>	$p$	r <sup>2</sup>	$p$	r <sup>2</sup>	$p$	r <sup>2</sup>	$p$
0–10	Clay	0.44	***	0.27	n.s.	0.36	*	0.14	n.s.	0.05	n.s.
	Silt	0.23	n.s.	0.14	n.s.	0.39	*	0.04	n.s.	-0.01	n.s.
	Sand	-0.44	**	-0.27	n.s.	-0.41	**	-0.12	n.s.	-0.02	n.s.
10–30	Clay	0.44	**	0.34	*	0.32	*	0.11	n.s.	0.01	n.s.
	Silt	0.27	n.s.	0.26	n.s.	0.32	*	0.1	n.s.	0.02	n.s.
	Sand	-0.36	*	-0.31	*	-0.34	*	-0.11	n.s.	-0.01	n.s.
30–60	Clay	-0.28	n.s.	0.27	n.s.	-0.31	*	0.03	n.s.	-0.42	**
	Silt	0.34	*	0.13	n.s.	0.49	***	-0.08	n.s.	0.32	*
	Sand	-0.11	n.s.	-0.27	n.s.	-0.22	n.s.	0.05	n.s.	-0.01	n.s.

Significance at different probability levels ( $p$ ) is indicated as “\*\*\*”  $\leq 0.001$ , “\*\*”  $\leq 0.01$ , “\*”  $\leq 0.05$ , n.s. = non-significant,  $n = 42$



**Table 4** Stepwise multiple regression relationships relating tiller density [tillers m<sup>-2</sup>], spike density [spikes m<sup>-2</sup>], grain yield [t ha<sup>-1</sup>], and grain protein content [%] to soil properties in different soil depths in Triesdorf 2016

Soil depth [cm]	Dependent variable	Regression relationship	R <sup>2</sup>	p
0–10	Tiller density	51.66 ± 29.43 (SOC) + 8.68 ± 4.75 (Clay)	0.43	<0.001
	Spike density			n.s.
	Grain yield	10.94 ± 2.48 (SOC) + 0.52 ± 0.18 (Silt) – 0.99 ± 0.44 (Clay)	0.46	<0.001
	Protein content			n.s.
10–30	Tiller density	127.36 ± 30.70 (SOC)	0.30	<0.001
	Spike density	9.89 ± 3.26 (Clay) – 95.34 ± 46.63 (BD) – 55.61 ± 29.43 (SOC)	0.24	<0.01
	Grain yield	11.30 ± 2.42 (SOC)	0.35	<0.001
	Protein content			n.s.
30–60	Tiller density	224.28 ± 52.34 (SOC) – 5.97 ± 1.77 (Clay)	0.37	<0.001
	Spike density			n.s.
	Grain yield	0.26 ± 0.11 (Silt) – 0.46 ± 0.14 (Clay) + 13.09 ± 4.40 (SOC)	0.45	<0.001
	Protein content	-0.02 ± 0.01 (Clay) + 0.01 ± 0.01 (Silt)	0.27	<0.01

The regression relationship shows coefficients with standard error of the factors included in the multiple regression  
 SOC = Soil organic carbon (mg g<sup>-1</sup>); Silt = Silt content (%); Clay = Clay content (%); BD = Bulk density (g cm<sup>-3</sup>)


Together the soil measures SOC and clay content in the topsoil (0–10 cm) explained 43.0% ( $p < 0.001$ ) of the variability in tiller density (Table 4) in Triesdorf. Plotting the influence of clay and silt content on tiller formation until BBCH 31 in a contour plot clarified that increasing clay content promotes tiller density (Fig. 3a), especially if clay content reaches levels above 16%. A clear impact of silt content on tillering could not be observed. Notably, data for these graphics have not been corrected for outliers and smoothed by moving averages. While it is common practice in precision farming to generate maps from higher numbers of data points showing smooth gradients (Joernsgaard and Halmoe 2003; Karampoiki et al. 2021), this study deliberately avoids doing so. Therefore, single values not fitting into the gradients became visible and impeded uniform slopes. In contrast to tillering, soil properties did not significantly affect spike density and protein content. In 0–10 cm soil depth, SOC, silt plus clay content explained 46.0% of the variance in grain yield of winter wheat ( $p < 0.001$ ) (Table 4). However, the contour plot conveys that silt content influenced grain yield much more than the soil's clay content (Fig. 3c).

In contrast, in the subsoil (30–60 cm), higher clay content was associated with a reduced tiller density, especially under silt concentrations <30% (Fig. 3b). Areas with higher silt content showed no adverse effects

of increasing clay contents on tillering. Despite this, the variance in tiller density was explained to 37.0% by SOC plus clay ( $p < 0.001$ ) (Table 4). Unexpectedly, the variation in spike density could not be significantly related to any subsoil's measures, even so rooting depth usually reached its maximum during this developmental period (Rasmussen and Thorup-Kristensen 2016). In contrast, subsoil's SOC, silt plus clay explained 45.0% of the variability in grain yield ( $<0.001$ ) (Table 4). A reduced grain yield was observed whenever clay content was higher than 12% paired with a silt content below 25% (Fig. 3d). The variables clay and silt could explain the variance in grain protein content to 27.0% ( $p < 0.01$ ).

In Asendorf, tiller and spike density and grain protein content were explained by the same variables in a multiple regression analysis, but the effects were far less pronounced (Table S3). No regression relationships existed between soil properties and grain yield and its components, tiller density, spike density, or grain protein content in the topsoil.

Summing up, the less pronounced impact of soil measures on wheat yield and its yield components can be attributed to the lower variability of the soil measures at Asendorf (Fig. S2). These results indicate that small-scale differences in a subsite of a field of 4200 m<sup>2</sup> caused a difference in wheat grain yield of 1.6 t ha<sup>-1</sup> in Triesdorf and 1.4 t ha<sup>-1</sup> in Asendorf (Table 2). Interestingly, when the soil variability



**Fig. 3** Influence of silt and clay content in the soil depth of 0–10 cm (**a** & **c**) and 30–60 cm (**b** & **d**) on tiller density [tillers  $m^{-2}$ ] in BBCH 31 (**a** & **b**) and on grain yield [ $t\ ha^{-1}$ ] of winter wheat (**c** & **d**) in Triesdorf, 2016

was high, as in Triesdorf, 35 to 46% of the variance in grain yield could be explained by only a few soil measures depending on soil depth (Table 4). However, no interaction was observed under more homogeneous conditions, as in Asendorf (Table S3).

#### Impact of small-scale differences in soil properties on yield components

To analyze if our approach of evaluating yield formation in response to vertical and lateral small-scale variations in soil properties is relevant for the practical application of site-specific crop production, we first clustered the dataset by soil texture. Farm practice typically uses this method to develop application maps based on large-scale soil differences, such as for fertilization approaches. Therefore, the Triesdorf

dataset was clustered by soil properties in different depths. Because clusters were mainly described by sand and clay content, the 3 clusters were named low sand (LS) with a sand content  $\leq 50\%$ , high sand, and low clay (HS-LC) characterized by a sand content  $> 50\%$  in combination with clay content  $\leq 15\%$ , and high sand and high clay (HS-HC) with sand content  $> 50\%$  and clay content  $> 15\%$  (Fig. 1).

All 3 clusters showed significant differences in sand content in every tested soil depth, with consistently smallest sand contents in LS and highest in HS-LC (Table 5). The highest clay content of the topsoil layers (0–30 cm), with around 18%, was observed in the LS cluster. In comparison, clay content decreased over HS-HC to around 14% in HS-LC. In contrast, the highest clay contents in the subsoil were recorded in the HS-HC cluster.

**Table 5** Soil texture of the 3 clusters characterized by low sand (LS), high sand & low clay (HS-LC), and high sand & high clay (HS-HC) (clay, silt, sand [%], and soil organic carbon (SOC) content [ $\text{mg g}^{-1}$ ]) and their impact on the plant measures grain yield [ $\text{t ha}^{-1}$ ], tiller density [ $\text{tillers m}^{-2}$ ], spike density [ $\text{spike m}^{-2}$ ], thousand-grain weight [ $1000 \text{ grains g}^{-1}$ ], rain per spike [ $\text{m}$ ], grain density [ $\text{grains m}^{-2}$ ], single spike weight [ $\text{g}$ ], grain density [ $\text{grains m}^{-2}$ ], thousand-grain weight [ $1000 \text{ grains g}^{-1}$ ], and grain protein content [%] at Triesdorf in 2016

Soil measures	0–10 cm				10–30 cm				30–60 cm			
	LS		HS		LS		HS		LS		HS	
	LC	HC	LC	HC	LC	HC	LC	HC	LC	HC	LC	HC
	(n=11)	(n=15)	(n=15)	(n=15)	(n=11)	(n=16)	(n=15)	(n=16)	(n=11)	(n=16)	(n=16)	(n=15)
Sand	47.8 c	57.9 a	53.7 b	47.0 c	47.0 c	57.2 a	53.4 b	45.7 c	45.7 c	59.0 a	52.5 b	52.5 b
Silt	34.3 a	27.9 b	28.7 b	34.9 a	34.9 a	27.3 c	29.5 b	36.7 a	36.7 a	28.6 b	28.0 b	28.0 b
Clay	18.0* a	14.1* b	17.6* a	18.1 a	18.1 a	13.9 c	16.6 b	17.5 b	17.5 b	12.3 c	19.5 a	19.5 a
SOC	16.4* a	10.9* b	16.4* a	14.0 a	14.0 a	10.9 b	12.8 a	7.6 a	7.6 a	5.9 b	6.5 b	6.5 b
Grain yield	10.8 a	10.1 c	10.4 b	10.8 a	10.8 a	10.2 b	10.4 b	10.7 a	10.7 a	10.5 a	10.2 b	10.2 b
Tillers density	508 a	444 b	508 a	516 a	516 a	461 b	511 ab	513 a	513 a	494 a	454 b	454 b
Spikes density	475 a	465 a	469 a	482 a	482 a	459 b	477 ab	483 a	483 a	456 a	471 a	471 a
Thousand-grain weight	53.5 a	53.6 a	53.0 a	53.2 a	53.2 a	53.1 a	53.7 a	53.1 a	53.1 a	53.3 a	53.4 a	53.4 a
Grains per spike	42.7 a	41.1 a	42.3 a	42.1 a	42.1 a	42.7 a	40.5 a	42.0 a	42.0 a	43.5 a	40.8 a	40.8 a
Grain density	20,253 a	18,909 b	19,730 a	20,229 a	20,229 a	19,416 b	19,303 ab	20,213 a	20,213 a	19,716 ab	19,048 b	19,048 b
Single spike weight	2.3 a	2.2 a	2.2 a	2.2 a	2.2 a	2.3 a	2.2 a	2.2 a	2.2 a	2.3 a	2.2 a	2.2 a
Protein content	13.4 a	13.3 a	13.4 a	13.4 a	13.4 a	13.3 a	13.3 a	13.4 ab	13.4 ab	13.4 a	13.3 b	13.3 b

Values representing means and different letters indicate significant differences among clusters at  $p \leq 0.05$  by Fisher-LSD test or (\*) Tukey HSD test

Irrespective of the soil depth, the highest SOC content was observed in the LS cluster with 16.4 mg g<sup>-1</sup>, 14.0 mg g<sup>-1</sup>, and 7.6 mg g<sup>-1</sup> from the topsoil to the subsoil and lowest in HS-LC with 10.9 mg g<sup>-1</sup>, 10.9 mg g<sup>-1</sup>, 5.9 mg g<sup>-1</sup>.

The grain yield of wheat responded significantly to differences in the soil texture of the 3 clusters. With 10.7 to 10.8 t ha<sup>-1</sup>, grain yield was highest in the LS cluster in all soil layers (Table 5). Only HS-LC in the subsoil reached nearly the same yield level with 10.5 t ha<sup>-1</sup>. At the high grain yield level of average 10.4 t ha<sup>-1</sup> in our study, the soil's small-scale differences caused only a maximum span of 10.1 to 10.8 t ha<sup>-1</sup> between different clusters but still caused a maximum variation in grain yield of 6.9%.

Interestingly, variations in yield between clusters in all tested soil layers were neither related to the thousand-grain weight nor the number of grains per spike or single-spike weight. Nevertheless, grain density showed significant differences between clusters. Similar to the findings in grain yield, the highest grain density was observed in the LS cluster for all 3 soil layers with a range of 20,213 to 20,253 grains m<sup>-2</sup>. A significantly smaller grain number was observed in the topsoil's HS-LC cluster, with 18,909 (0–10 cm) and 19,416 grains m<sup>-2</sup> (10–30 cm). Only in the subsoil, the HS-HC cluster with 19,048 grains m<sup>-2</sup> led to the lowest grain setting. Interestingly, yield components formed during vegetative plant development responded stronger to differences in soil texture than those formed during the generative phase. In fact, tillering in the clusters followed the same pattern as grain density. Even the levels of significance were the same. Moreover, the tiller number was relatively low, with values between 444 and 516 tillers m<sup>-2</sup>, compared to other experiments reaching a high grain yield (Sieling et al. 2005; Zhang et al. 2020). The only drawback was that the tiller density was not transferred to spike density in the 0–10 cm and the subsoil layer. Therefore, only the clusters of the second topsoil layer (10–30 cm) showed the same patterns as grain density and grain yield. Grains' protein content was the only measure during the grain-filling phase, which was influenced by the soil differences represented by the clusters. Nevertheless, only the clustering in the subsoil led to significant effects in which HS-LC, with 13.4%, reached the highest protein content. At the same time, HS-HC resulted in only an infinitesimal reduction of 0.1% points to the smallest value in this analysis.

Summing up, the impact of soils' small-scale differences at Triesdorf was prominent on tillering, spike density, grain density, grain yield, and even grain protein content employing the zoning technique as common practice in site-specific farming. At Asendorf, the cluster approach did not lead to statistically significant differences in plant measures due to a less pronounced small-scale variability of soil parameters (Table 1).

## Discussion

Defining the appropriate resolution of sensor data, application maps, and the section size of agricultural machinery is an important step toward improving crop production systems through site-specific farming approaches. Especially when the economic value added is limited or even absent in practice, a higher resolution is often cited to improve the benefit of site-specific application strategies (Katalin et al. 2014; Späti et al. 2021). Indeed, this approach works well, especially in image-based decision systems, such as herbicide applications with single nozzle switching (Hørfarter et al. 2019; Kämpfer and Nordmeyer 2021; Oerke et al. 2010) or in the evaluation of canopy shapes and above-ground biomass, especially in multi-species cover crops (Kümmerer et al. 2023). In practice, however, high resolution is still limited, especially for soil data, like for variable-rate application in seeding (Šarauskiš et al. 2022) and in nitrogen (Argento et al. 2021) or basic fertilization (Wagner and Marz 2017).

To tackle whether an increase in resolution can improve yield or resource efficiency, we did not conduct seeding or fertilization experiments to evaluate the performance of treatments under different soil conditions. Instead, we chose an inverse approach and evaluated the plant development under a regime of homogeneous crop production by different soil conditions. For this purpose, the small-scale variability of soil in the lateral and vertical dimensions was detected, and grain yield, its yield components, and quality measures were analyzed at two locations.

Spatial variability of small-scale differences in soil properties

In common practice, the site-specific variability of soil is detected laterally and vertically by electromagnetic resistivity (André et al. 2012; Basso et al.

2010; Boenecke et al. 2018) or soil conductivity (Cossel et al. 2019; Kühn et al. 2009; Stadler et al. 2015). The resolution of these methods is limited more by sampling grid and application conditions than by the limitations of the technology itself (André et al. 2012). Nevertheless, the exact composition of soil properties that trigger the signal, and thus the exact impact on plant growth, remains to be determined.

Soil texture variability increased from the topsoil with four to subsoil with six texture classes according to German soil classification in the field trials (Fig. 2), representing one factor for inhomogeneous growing conditions in Triesdorf. In Asendorf, soil conditions were more homogeneous. Here, the topsoil was represented by only two different soil texture classes, with the majority of 74–76% classified as Us. The homogeneity was even more pronounced in the subsoil, where 35 of the 42 plots were classified as Us (Fig. 2c).

At the Triesdorf field site, geological clay lenses resulted in a CV of 23.9% of the clay content, indicating a high variability of 9.6 to 25.6% clay in the subsoil (Table S1). Studies reported similar CV in subsoil's soil texture with larger grid sizes on experimental sites of 11–30 ha with 20 soil sampling points (Boenecke et al. 2018; Cammarano et al. 2019). Thus, there is clear evidence that small-scale variability can typically be as large as observed across the entire field site. This is also supported by other studies that report similar small-scale differences in soil textures as identified in our experiment (Grote et al. 2010; Martinez-Turanzas et al. 1997), and recent studies have shown that heterogeneous soil texture is common in arable fields and that higher resolution is beneficial (Boenecke et al. 2018; Bölenius et al. 2017; Mallarino et al. 1999; Stadler et al. 2015; Usovich and Lipiec 2017; van Meirvenne 2003). To our knowledge, no studies have combined high-resolution sampling (9.0 m x 10.5 m grid size) of soil properties at lateral and vertical scales with yield measures in European environments.

**Benefit of considering small-scale variability on plant properties**

At the two field sites, fundamental differences in yield formation were observed due to different soil

properties and climatic differences (Fig. S3), such as a milder winter and a more extended period for tillering in Asendorf. Although leaf appearance and tillering are strongly influenced by temperature and photoperiod (Ochagavía et al. 2017), this could not be the only explanation for the enormous tiller formation in Asendorf compared to Triesdorf (Table 2). Of course, the seeding time differs in agricultural practice between different sites, but increased vegetative development cannot explain differences in tillering between the northern and southern locations (Budzyński et al. 2018). Therefore, the main driver for the increased tillering in Asendorf must be the different soil textures (Fig. 2). Several studies have shown the impact of soil clay content, in particular, on the formation of yield components such as tillering in cereals (Bölenius et al. 2017; Dou et al. 2016; Gozdowski et al. 2017; Yunusa and Sedgley 1992). Nevertheless, the effect of soil texture on tiller number could only be observed in tendency in the stepwise multiple regression analysis for Asendorf (Table S3), and the tiller number only ranged from 725 to 855 tillers  $m^{-2}$  (Table 2). However, other studies have observed the impact of soil texture on wheat development and grain yield under similar climatic conditions (Johnen et al. 2014) or in general (Carvalho Mendes et al. 2021; Pan et al. 2020). On the other hand, the influence of soil texture on tillering was present in Triesdorf and was mainly determined by clay and SOC content in the top and subsoil (Table 4).

Interestingly, although the site-specific differences by soil types are at first sight more heterogeneous in Triesdorf (Fig. 2), the CV for clay, silt, and sand were at a similar level as in Asendorf (Tables S1, S2). Nevertheless, the variation in Asendorf caused a variation in tillering and resulted in a CV for tiller density of 3.7% compared to 10.8% for Triesdorf. As an explanation for this phenomenon, high silt concentrations (Watson et al. 2002), as observed in Asendorf, are often mentioned in the literature.

However, the high tiller density in Asendorf, where site-specific variation was low, did not result in similarly high yields as in Triesdorf (Table 2), even though the site is generally considered to have higher yield potential due to favorable climatic conditions for winter wheat (Lüttger and Feike 2018). Nevertheless, the tiller density in Asendorf was too

high compared to Triesdorf, which is indicated by an average of 9 fewer grains per spike, resulting in a decrease in grain density by 2788 grains  $m^{-2}$ . In addition, the grain filling was also negatively influenced by the elevated tiller density, so the single spike weight was reduced by approximately 1.0 g compared to Triesdorf. The thousand-grain weight reached only 39.6 g compared to 53.3 g in Triesdorf (Table 2). The observation of decreased grain filling under a high organ setting, e.g., tillering under the condition of a specific location, is consistent with the findings of other studies (Slafer et al. 2014; Berry et al. 2003; Langridge 2017). Besides the general differences in yield formation between the two contrasting sites, the influence of small-scale variability in soil properties on yield was evident on both sites. It strongly depended on lateral soil texture differences across the field site and on vertical differences along the soil depth profile, especially under the condition of the soil properties in Triesdorf (Fig. 2). Subsoil conditions, which are important for water and nutrient availability (Kautz et al. 2013; Kuhlmann and Baumgärtel 1991; Kuhlmann 1990; Kuhlmann et al. 1989; Zheng et al. 2023), explain the differences in yield formation, especially during generative development. In another study, small-scale variability in subsoil texture was observed to influence water and nutrient fluxes (Leinemann et al. 2016).

However, the influence of small-scale soil texture differences in different soil layers on tiller density was rarely investigated. Interestingly, the soil measures SOC and clay in the first soil layer of 0–10 cm explained 43.0% of the variability in tiller density (Table 4) and decreased to 30% in the subsoil. At Asendorf, soil properties had little or no significant impact on tillering, regardless of which soil layer was analyzed. Previous studies have shown that winter wheat roots reach the subsoil before tillering (Hodgkinson et al. 2017; Zhou et al. 2022). Increasing clay content was associated with increased tiller density in the topsoil layer, while opposite effects occurred in the subsoil (Fig. 3; Table 4). An explanation for this observation is provided by Coelho Filho et al. (2013), who suggested that increasing soil impedance due to changes in soil texture negatively affects tiller density. At Triesdorf, 46.0% of the yield variability was explained by soil properties of the first layer (0–10 cm), 35.0% of the 10–30 cm layer, and 45.0% of the subsoil (Table 4). Similar observations of small-scale lateral differences

in soil texture and chemical soil properties linked to grain yield were reported in former studies (Hausherr Lüder et al. 2018; Maidl et al. 1999). Nevertheless, under the conditions of Asendorf, no significant impact of soil properties on yield variability has been observed (Table S3).

Taken together, we suggest a smaller grid size for site-specific crop management beyond the current standard grid size of approx. 20 m to 20 m, derived from satellite-based remote sensing, can increase grain yield, especially on sites with highly inhomogeneous soil properties. Based on our results, we recommend using a higher resolution of at least 10.5 m to 9.0 m as the grid size.

SOC is relevant at all soil depths

In addition to soil texture, the wheat yield was strongly determined by SOC, which explained 45–46% of the yield variability in Triesdorf, irrespective of the soil depth studied (Table 4). This is consistent with previous findings of increased grain yield at higher subsoil SOC (Usowicz and Lipiec 2017). In line with other studies, SOC content decreased with soil depth and was associated with the soil type (Börjesson et al. 2018; Poeplau et al. 2020). In Asendorf, no relationships between soil SOC and grain yield were observed (Table S3). This could be explained by a combination of non-limited precipitation (Fig. S3), less pronounced subsoil heterogeneity (Fig. 2c), and a generally lower yield level (Table 2), which did not fully exploit the yield potential. In previous studies, SOC increased nitrogen use efficiency (Oelofse et al. 2015) and had a positive influence on cereal grain yield (Šeremešić et al. 2020; Stein et al. 1997), even under soil conditions similar to the present study, as reported in Poland on sandy soils in a coarse sampling grid (Usowicz and Lipiec 2017). In addition, a previous study by Cammarano et al. (2019) on winter barley highlighted the importance of soil texture and bulk density, closely related to SOC, on the vertical distribution of nitrogen below 30 cm soil depth and showed an impact on crop yield stability. In general, the need for nutrient and water uptake from the subsoil is closely related to the topsoil conditions (Engels et al. 1994; Jin et al. 2015), such that nutrient uptake from the subsoil does not necessarily result in higher grain yield (Hodgkinson et al. 2017). Thus, in Asendorf, non-limiting precipitation may have reduced the impact of the soil properties on grain yield.

Our data support the view that SOC provides a valuable indication of growing conditions and can benefit site-specific management decisions. Because site-specific yield potential and mineralization patterns are closely related to soil properties, variable-rate mineral fertilizer application will likely improve nitrogen use efficiency. High-resolution information on soil properties is essential to achieve this efficiency (Argento et al. 2021; Tubaña et al. 2008). However, despite the environmental benefits, the economic value of the technology for smaller farms is still questionable (Späti et al. 2021).

## Conclusion

The present study shows lateral and vertical soil property variations significantly influence winter wheat development and grain yield, even at small field scales. The approach to defining clusters of physical soil properties within different soil depths is suitable for quantifying the effect of soil heterogeneity on yield patterns and assessing the impact and reliability for practical use. The described approach is particularly relevant whenever soils show small-scale soil texture or SOC differences. Future studies should define the optimal grid size for site-specific crop production strategies by considering soil heterogeneity to improve grain yield, yield stability, and input efficiency for sustainable crop production.

**Acknowledgements** This work was supported by the German Federal Ministry of Education and Research [grant number 031A559 - CATCHY]. We are grateful for the excellent technical support by Silke Bokeloh, Gerald Fiedler, Steffen Schierding, Farruh Ulmasov, and Stefan Uhl during the field trial, soil sampling, and lab work.

**Funding** Open Access funding enabled and organized by Projekt DEAL.

**Data availability** The Datasets from the current study are available in the zenodo database, <https://doi.org/10.5281/zenodo.8245951>.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not

included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Ad hoc AG Boden (2005) *Bodenkundliche Kartieranleitung*, 5th edn. Schweizerbart i. Komm; Bundesanst. für Geowiss. und Rohstoffe, Stuttgart
- Akima H, Gebhardt A, Petzold T, Maechler M (2020) Package 'akima'. Interpolation of Irregularly and Regularly Spaced Data. R. <https://cran.r-project.org/web/packages/akima/index.html>
- Amelung W, Blume H-P, Fleige H, Horn R, Kandeler E, Kögel-Knabner I, Kretschmar R, Stahr K, Wilke B-M (2018) *Scheffer/Schachtschabel Lehrbuch der Bodenkunde*, 17th edn. Springer, Berlin
- André F, van Leeuwen C, Saussez S, van Durmen R, Bogaert P, Moghadas D, de Ressaéguier L, Delvaux B, Vereecken H, Lambot S (2012) High-resolution imaging of a vineyard in south of France using ground-penetrating radar, electromagnetic induction and electrical resistivity tomography. *J Appl Geophys* 78:113–122. <https://doi.org/10.1016/j.jappgeo.2011.08.002>
- Argento F, Anken T, Abt F, Vogelsanger E, Walter A, Liebisch F (2021) Site-specific nitrogen management in winter wheat supported by low-altitude remote sensing and soil data. *Precision Agric* 22:364–386. <https://doi.org/10.1007/s11119-020-09733-3>
- Barnes AP, Soto I, Eory V, Beck B, Balafoutis A, Sánchez B, Vangeyer J, Fountas S, van der Wal T, Gómez-Barbero M (2019) Exploring the adoption of precision agricultural technologies: a cross regional study of EU farmers. *Land Use Policy* 80:163–174. <https://doi.org/10.1016/j.landusepol.2018.10.004>
- Barraclough PB, Leigh RA (1984) The growth and activity of winter wheat roots in the field: the effect of sowing date and soil type on root growth of high-yielding crops. *J Agric Sci* 103:59–74. <https://doi.org/10.1017/S002185960004332X>
- Basso B, Amato M, Bitella G, Rossi R, Kravchenko A, Sartori L, Carvahlo LM, Gomes J (2010) Two-dimensional spatial and temporal variation of soil physical properties in tillage systems using electrical resistivity tomography. *Agron J* 102:440–449. <https://doi.org/10.2134/agronj2009.0298>
- Bastos LM, Carciochi W, Lollato RP, Jaenisch BR, Rezende CR, Schwalbert R, Vara Prasad PV, Zhang G, Fritz AK, Foster C, Wright Y, Young S, Bradley P, Ciampitti IA (2020) Winter wheat yield response to plant density as a function of yield environment and tillering potential: a review and field studies. *Front Plant Sci* 11:54. <https://doi.org/10.3389/fpls.2020.00054>
- Berry P, Spink J, Foulkes M, Wade A (2003) Quantifying the contributions and losses of dry matter from non-surviving shoots in four cultivars of winter wheat. *Field Crop Res* 80:111–121. [https://doi.org/10.1016/S0378-4290\(02\)00174-0](https://doi.org/10.1016/S0378-4290(02)00174-0)

- Boenecke E, Lueck E, Ruehlmann J, Gruendling R, Franko U (2018) Determining the within-field yield variability from seasonally changing soil conditions. *Precis Agric* 19:750–769. <https://doi.org/10.1007/s11119-017-9556-z>
- Bölenius E, Stenberg B, Arvidsson J (2017) Within field cereal yield variability as affected by soil physical properties and weather variations – a case study in east central Sweden. *Geoderma Reg* 11:96–103. <https://doi.org/10.1016/j.geodrs.2017.11.001>
- Börjesson G, Bolinder MA, Kirchmann H, Kätterer T (2018) Organic carbon stocks in topsoil and subsoil in long-term ley and cereal monoculture rotations. *Biol Fertil Soils* 54:549–558. <https://doi.org/10.1007/s00374-018-1281-x>
- Bozdogan H (1987) Model selection and Akaike's Information Criterion (AIC): the general theory and its analytical extensions. *Psychometrika* 52:345–370. <https://doi.org/10.1007/BF02294361>
- Breuning Madsen H (1985) Distribution of spring barley roots in danish soils, of different texture and under different climatic conditions. *Plant Soil* 88:31–43. <https://doi.org/10.1007/BF02140664>
- Brogi C, Huisman JA, Weihermüller L, Herbst M, Vereecken H (2021) Added value of geophysics-based soil mapping in agro-ecosystem simulations. *Soil* 7:125–143. <https://doi.org/10.5194/soil-7-125-2021>
- Budzyński WS, Bepirszcz K, Jankowski KJ, Dubis B, Hłasko-Nasalska A, Sokólski MM, Olszewski J, Załuski D (2018) The responses of winter cultivars of common wheat, durum wheat and spelt to agronomic factors. *J Agric Sci* 156:1163–1174. <https://doi.org/10.1017/S0021859619000054>
- Buttafuoco G, Castrignanò A, Cucci G, Lacolla G, Lucà F (2017) Geostatistical modelling of within-field soil and yield variability for management zones delineation: a case study in a durum wheat field. *Precis Agric* 18:37–58. <https://doi.org/10.1007/s11119-016-9462-9>
- Cammarano D, Holland J, Basso B, Fontana F, Murgia T, Lange C, Taylor J, Ronga D (2019) Integrating geospatial tools and a crop simulation model to understand spatial and temporal variability of cereals in Scotland. In: Stafford JV (ed) *Precision agriculture '19*. Wageningen Academic Publishers, Wageningen, pp 29–35
- Carvalho Mendes I, Martinhão Gomes Sousa D, Dario Dantas O, Alves Castro Lopes A, Bueno Reis Junior F, Ines Oliveira M, Montandon Chaer G (2021) Soil quality and grain yield: a win-win combination in clayey tropical oxisols. *Geoderma* 388:114880. <https://doi.org/10.1016/j.geoderma.2020.114880>
- Cassmann KG (1999) Ecological intensification of cereal production systems: yield potential, soil quality. *Precis Agric* 69:5952–5959
- Coelho Filho MA, Colebrook EH, Lloyd DPA, Webster CP, Mooney SJ, Phillips AL, Hedden P, Whalley WR (2013) The involvement of gibberellin signalling in the effect of soil resistance to root penetration on leaf elongation and tiller number in wheat. *Plant Soil* 371:81–94. <https://doi.org/10.1007/s11104-013-1662-8>
- Conant RT, Ogle SM, Paul EA, Paustian K (2011) Measuring and monitoring soil organic carbon stocks in agricultural lands for climate mitigation. *Front Ecol Environ* 9:169–173. <https://doi.org/10.1890/090153>
- Cox MS, Gerard PD, Wardlaw MC, Abshire MJ (2003) Variability of selected soil properties and their relationships with soybean yield. *Soil Sci Soc Am J* 67:1296–1302
- Crain JL, Waldschmidt KM, Raun WR (2013) Small-scale spatial variability in winter wheat production. *Commun Soil Sci Plant Anal* 44:2830–2838. <https://doi.org/10.1080/00103624.2013.812735>
- Crist JW, Weaver JE (1924) Absorption of nutrients from subsoil in relation to crop yield. *Bot Gaz* 77:121–148. <https://doi.org/10.1086/333295>
- Denora M, Amato M, Brunetti G, de Mastro F, Perniola M (2022) Geophysical field zoning for nitrogen fertilization in durum wheat (*Triticum durum* Desf). *PLoS ONE* 17:e0267219. <https://doi.org/10.1371/journal.pone.0267219>
- Dou F, Soriano J, Tabien RE, Chen K (2016) Soil texture and cultivar effects on rice (*Oryza sativa*, L.) grain yield, yield components and water productivity in three water regimes. *PLoS ONE* 11:e0150549. <https://doi.org/10.1371/journal.pone.0150549>
- Engels C, Mollenkopf M, Marschner H (1994) Effect of drying and rewetting the topsoil on root growth of maize and rape in different soil depths. *Z für Pflanzenernährung und Bodenkunde* 157:139–144. <https://doi.org/10.1002/jpln.19941570213>
- Eurostat (2021) Key figures on the european food chain: 2021 edition, 2021 edition. Publications Office of the European Union, Luxembourg
- Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M, Mueller ND, O'Connell C, Ray DK, West PC, Balzer C, Bennett EM, Carpenter SR, Hill J, Monfreda C, Polasky S, Rockström J, Sheehan J, Siebert S, Tilman D, Zaks DPM (2011) Solutions for a cultivated planet. *Nature* 478:337–342. <https://doi.org/10.1038/nature10452>
- Freibauer A, Rounsevell MD, Smith P, Verhagen J (2004) Carbon sequestration in the agricultural soils of Europe. *Geoderma* 122:1–23. <https://doi.org/10.1016/j.geoderma.2004.01.021>
- Gebbers R, Adamchuk VI (2010) Precision agriculture and food security. *Sci (New York N Y)* 327:828–831. <https://doi.org/10.1126/science.1183899>
- Ghaley BB, Wösten H, Olesen JE, Schelde K, Baby S, Karki YK, Børgesen CD, Smith P, Yeluripati J, Ferrise R, Bindi M, Kuikman P, Lesschen J-P, Porter JR (2018) Simulation of soil organic carbon effects on long-term winter wheat (*Triticum aestivum*) production under varying fertilizer inputs. *Front Plant Sci* 9:1158. <https://doi.org/10.3389/fpls.2018.01158>
- Gozdowski D, Leszczyńska E, Stepień M, Rozbicki J, Samborski S (2017) Within-field variability of winter wheat yield and grain quality versus soil properties. *Commun Soil Sci Plant Anal* 48:1029–1041. <https://doi.org/10.1080/00103624.2017.1323091>
- Grote K, Anger C, Kelly B, Hubbard S, Rubin Y (2010) Characterization of soil water content variability and soil texture using GPR Groundwave techniques. *J Environ Eng Geophys* 15:93–110. <https://doi.org/10.2113/JEEG15.3.93>
- Guo Z, Zhang Z, Zhou H, Wang D, Peng X (2019) The effect of 34-year continuous fertilization on the SOC



- physical fractions and its chemical composition in a vertisol. *Sci Rep* 9:2505. <https://doi.org/10.1038/s41598-019-38952-6>
- Harris D, Horwath WR, van Kessel C (2001) Acid fumigation of soils to remove carbonates prior to total organic carbon or CARBON-13 isotopic analysis. *Soil Sci Soc Am J* 65:1853–1856. <https://doi.org/10.2136/sssaj2001.1853>
- Hartigan JA, Wong MA (1979) A k-means clustering algorithm. *J Roy Stat Soc* 28:100–108. <https://doi.org/10.2307/2346830>
- Hausherr Lüder R-M, Qin R, Richner W, Stamp P, Noulas C (2018) Spatial variability of selected soil properties and its impact on the grain yield of oats (*Avena sativa* L.) in small fields. *J Plant Nutr* 41:2446–2469. <https://doi.org/10.1080/01904167.2018.1527935>
- Hbirkou C, Pätzold S, Mahlein A-K, Welp G (2012) Airborne hyperspectral imaging of spatial soil organic carbon heterogeneity at the field-scale. *Geoderma* 175–176:21–28. <https://doi.org/10.1016/j.geoderma.2012.01.017>
- Hess M, Barralis G, Bleiholder H, Buhner L, Eggers T, Hack H, Stauss R (1997) Use of the extended BBCH scale - general for the descriptions of the growth stages of mono- and dicotyledonous weed species. *Weed Res* 37:433–441. <https://doi.org/10.1046/j.1365-3180.1997.d01-70.x>
- Hodgkinson L, Dodd IC, Binley A, Ashton RW, White RP, Watts CW, Whalley WR (2017) Root growth in field-grown winter wheat: some effects of soil conditions, season and genotype. *Eur J Agron* 91:74–83. <https://doi.org/10.1016/j.eja.2017.09.014>
- Hørfarter R, Thorsted MD, Stougård K, Poulsen HV (2019) Precision spraying by combining a variable rate application map with an on/off map. 2019. In: Stafford JV (ed), Precision agriculture '19. Papers presented at the 12th European Conference on Precision Agriculture. Montpellier. Wageningen Academic Publishers, pp 53–59
- IUSS Working Group WRB (2015) World reference base for soil resources 2014. International soil classification system for naming soils and creating legends for soil maps. Update 2015, vol 106. FAO, Rome
- Jin K, Shen J, Ashton RW, White RP, Dodd IC, Parry MAJ, Whalley WR (2015) Wheat root growth responses to horizontal stratification of fertiliser in a water-limited environment. *Plant Soil* 386:77–88. <https://doi.org/10.1007/s11104-014-2249-8>
- Joernsgaard B, Halmoe S (2003) Intra-field yield variation over crops and years. *Eur J Agron* 19:23–33. [https://doi.org/10.1016/S1161-0301\(02\)00016-3](https://doi.org/10.1016/S1161-0301(02)00016-3)
- Johnen T, Boettcher U, Kage H (2014) An analysis of factors determining spatial variable grain yield of winter wheat. *Eur J Agron* 52:297–306. <https://doi.org/10.1016/j.eja.2013.08.005>
- Kämpfer C, Nordmeyer H (2021) Teilflächenspezifische Unkrautbekämpfung als ein Baustein von Smart Farming. 284–291 Seiten / Journal für Kulturpflanzen, Bd.73 Nr. 7–8 (2021). <https://doi.org/10.5073/JFK.2021.07-08.13>
- Karampoiki M, Heiß A, Sharipov GM, Mahmood S, Todman LC, Murdoch AJ, Griepentrog HW, Paraforos DS (2021) 77. Producing grain yield maps by merging combine harvester and remote sensing data. In: Stafford J (ed), Precision agriculture '21. Papers presented at the 13th European Conference on Precision Agriculture. Wageningen Academic Publishers, Wageningen, pp 645–651
- Katalin T-G, Rahoveanu T, Magdalena M, István T (2014) Sustainable new agricultural technology – Economic aspects of precision crop protection. *Procedia Econ Financ* 8:729–736. [https://doi.org/10.1016/S2212-5671\(14\)00151-8](https://doi.org/10.1016/S2212-5671(14)00151-8)
- Kautz T, Amelung W, Ewert F, Gaiser T, Horn R, Jahn R, Javaux M, Kemna A, Kuzyakov Y, Munch J-C, Pätzold S, Peth S, Scherer HW, Schloter M, Schneider H, Vunderborght J, Vetterlein D, Walter A, Wiesenberger GL, Köpke U (2013) Nutrient acquisition from arable subsoils in temperate climates: a review. *Rev Soil Biol Biochem* 57:1003–1022. <https://doi.org/10.1016/j.soilbio.2012.09.014>
- Kirkegaard JA, Lilley JM, Howe GN, Graham JM (2007) Impact of subsoil water use on wheat yield. *Aust J Agric Res* 58:303. <https://doi.org/10.1071/AR06285>
- Kottek M, Grieser J, Beck C, Rudolf B, Rubel F (2006) World Map of the Köppen-Geiger climate classification updated. *Meteorol Z* 15:259–263. <https://doi.org/10.1127/0941-2948/2006/0130>
- Kuhlmann H (1990) Importance of the subsoil for the K nutrition of crops. *Plant Soil* 127:129–136. <https://doi.org/10.1007/BF00010845>
- Kuhlmann H, Baumgärtel G (1991) Potential importance of the subsoil for the P and mg nutrition of wheat. *Plant Soil* 137:259–266. <https://doi.org/10.1007/BF00011204>
- Kuhlmann H, Barraclough PB, Weir AH (1989) Utilization of mineral nitrogen in the subsoil by winter wheat. *J Plant Nutr Soil Sci* 152:291–295. <https://doi.org/10.1002/jpln.19891520305>
- Kühn J, Brenning A, Wehrhan M, Koszinski S, Sommer M (2009) Interpretation of electrical conductivity patterns by soil properties and geological maps for precision agriculture. *Precision Agric* 10:490–507. <https://doi.org/10.1007/s11119-008-9103-z>
- Kümmerer R, Noack PO, Bauer B (2023) Using high-resolution UAV Imaging to measure canopy height of diverse cover crops and predict Biomass. *Remote Sens* 15:1520. <https://doi.org/10.3390/rs15061520>
- Langridge P (2017) Achieving sustainable cultivation of wheat volume 2. Cultivation techniques, 1st edn. Burleigh Dodds Science Publishing Limited, Cambridge
- LaRuffa JM, Raun WR, Phillips SB, Solie JB, Stone ML, Johnson GV (2001) Optimum field element size for maximum yield in winter wheat, using variable nitrogen rates. *J Plant Nutr* 24:313–325. <https://doi.org/10.1081/PLN-100001390>
- Leinemann T, Mikutta R, Kalbitz K, Schaarschmidt F, Guggenberger G (2016) Small scale variability of vertical water and dissolved organic matter fluxes in sandy Cambisol subsoils as revealed by segmented suction plates. *Biogeochemistry* 131:1–15. <https://doi.org/10.1007/s10533-016-0259-8>
- Leroux C, Tisseyre B (2019) How to measure and report within-field variability: a review of common indicators and their sensitivity. *Precision Agric* 20:562–590. <https://doi.org/10.1007/s11119-018-9598-x>
- Liang Z, Mortensen E, de Notaris C, Elsgaard L, Rasmussen J (2022) Subsoil carbon input by cover crops depends on management history. *Agric Ecosyst Environ* 326:107800. <https://doi.org/10.1016/j.agee.2021.107800>
- Lüttger AB, Feike T (2018) Development of heat and drought related extreme weather events and their effect on

- winter wheat yields in Germany. *Theoret Appl Climatol* 132:15–29. <https://doi.org/10.1007/s00704-017-2076-y>
- Machado S, Bynum JED, Archer TL, Lascano RJ, Wilson LT, Bordovsky J, Segarra E, Bronson K, Nesmith DM, Xu W (2000) Spatial and temporal variability of corn grain yield: Site-specific relationships of biotic and abiotic factors. *Precis Agric* 2:359–376. <https://doi.org/10.1023/A:1012352032031>
- Maidl F-X, Brunner R, Sticksel E, Fischbeck G (1999) Ursachen kleinräumiger Ertragsschwankungen im bayerischen Tertiärhügelland und Folgerungen für eine teilschlagbezogene Düngung. *J Plant Nutr Soil Sci* 162:337–342. [https://doi.org/10.1002/\(SICI\)1522-2624\(199906\)162:3%3C337::AID-JPLN337%3E3.0.CO;2-2](https://doi.org/10.1002/(SICI)1522-2624(199906)162:3%3C337::AID-JPLN337%3E3.0.CO;2-2)
- Mallarino AP, Oyarzabal ES, Hinz PN (1999) Interpreting within-field relationships between crop yields and soil and plant variables using factor analysis. *Precision Agric* 1:15–25. <https://doi.org/10.1023/A:1009940700478>
- Manly BFJ, Navarro Alberto JA (2004) *Multivariate statistical methods. A primer*. Chapman and Hall/CRC, an imprint of Taylor and Francis, Boca Raton
- Martinez-Turanzas GA, Coffin DP, Burke IC (1997) Development of microtopography in a semi-arid grassland: Effects of disturbance size and soil texture. *Plant Soil* 191:163–171. <https://doi.org/10.1023/A:1004286605052>
- Mendiburu F (2021) Package ‘agricolae’. *Statistical Procedures for Agricultural Research*. R. <https://cran.r-project.org/web/packages/agricolae/index.html>
- Miles J (2014) In: Balakrishnan N, Colton T, Everitt B, Piegorsch W, Ruggeri F, Teugels JL (eds) *Tolerance and variance inflation factor*. Statistics Reference Online. Wiley, Wiley StatsRef
- Moey J (2018) Package ‘soiltexture’. *Functions for Soil texture plot, classification and Transformation*. R. <https://cran.r-project.org/web/packages/soiltexture/index.html>
- Mulla DJ (2013) Twenty five years of remote sensing in precision agriculture: key advances and remaining knowledge gaps. *Biosyst Eng* 114:358–371. <https://doi.org/10.1016/j.biosystemseng.2012.08.009>
- Nordmeyer H (2006) Reduktionsprogramm chemischer Pflanzenschutz – Beitrag der teilflächenspezifischen Unkrautbekämpfung. *Nachrichtenbl Deut Pflanzenschutz* 58:317–322
- Ochagavía H, Prieto P, Savin R, Griffiths S, Slafer GA (2017) Duration of developmental phases, and dynamics of leaf appearance and tillering, as affected by source and doses of photoperiod insensitivity alleles in wheat under field conditions. *Field Crop Res* 214:45–55. <https://doi.org/10.1016/j.fcr.2017.08.015>
- Oelofse M, Markussen B, Knudsen L, Schelde K, Olesen JE, Jensen LS, Bruun S (2015) Do soil organic carbon levels affect potential yields and nitrogen use efficiency? An analysis of winter wheat and spring barley field trials. *Eur J Agron* 66:62–73. <https://doi.org/10.1016/j.eja.2015.02.009>
- Oerke E-C, Gerhards R, Menz G, Sikora RA (2010) *Precision crop protection - the challenge and use of heterogeneity*. Springer Netherlands, Dordrecht
- Osanaí Y, Knox O, Nachimuthu G, Wilson B (2021) Contrasting agricultural management effects on soil organic carbon dynamics between topsoil and subsoil. *Soil Res* 59:24. <https://doi.org/10.1071/SR19379>
- Palmer J, Thorburn PJ, Biggs JS, Dominati EJ, Probert ME, Meier EA, Huth NI, Dodd M, Snow V, Larsen JR, Parton WJ (2017) Nitrogen cycling from increased soil organic carbon contributes both positively and negatively to ecosystem services in wheat agro-ecosystems. *Front Plant Sci* 8:731. <https://doi.org/10.3389/fpls.2017.00731>
- Pan J, Shang Y, Zhang WJ, Chen X, Cui Z (2020) Improving soil quality for higher grain yields in chinese wheat and maize production. *Land Degrad Dev* 31:1125–1137. <https://doi.org/10.1002/ldr.3515>
- Patzold S, Mertens FM, Bornemann L, Koleczek B, Franke J, Feilhauer H, Welp G (2008) Soil heterogeneity at the field scale: a challenge for precision crop protection. *Precis Agric* 9:367–390. <https://doi.org/10.1007/s11119-008-9077-x>
- Poepflau C, Kätterer T (2017) Is soil texture a major controlling factor of root:shoot ratio in cereals? *Eur J Soil Sci* 68:964–970. <https://doi.org/10.1111/ejss.12466>
- Poepflau C, Jacobs A, Don A, Vos C, Schneider F, Wittnebel M, Tiemeyer B, Heidkamp A, Prietz R, Flessa H (2020) Stocks of organic carbon in german agricultural soils—key results of the first comprehensive inventory. *J Plant Nutr Soil Sci* 183:665–681. <https://doi.org/10.1002/jpln.202000113>
- Poepflau C, Don A, Schneider F (2021) Roots are key to increasing the mean residence time of organic carbon entering temperate agricultural soils. *Glob Change Biol* 27:4921–4934. <https://doi.org/10.1111/gcb.15787>
- R Core Team (2021) R: a language and environment for statistical computing. Foundation for Statistical Computing, Vienna
- Rasmussen IS, Thorup-Kristensen K (2016) Does earlier sowing of winter wheat improve root growth and N uptake? *Field Crop Res* 196:10–21. <https://doi.org/10.1016/j.fcr.2016.05.009>
- Rawls WJ (1983) Estimating soil bulk density from particle size analysis and organic matter content. *Soil Sci* 1983:123–125
- Rentschler T, Werban U, Ahner M, Behrens T, Gries P, Scholten T, Teuber S, Schmidt K (2020) 3D mapping of soil organic carbon content and soil moisture with multiple geophysical sensors and machine learning. *Vadose Zone J* 19:e20062. <https://doi.org/10.1002/vzj2.20062>
- Rodriguez-Moreno F, Lukas V, Neudert L, Dryšlová T (2014) Spatial interpretation of plant parameters in winter wheat. *Precis Agric* 15:447–465. <https://doi.org/10.1007/s11119-013-9340-7>
- Roth GW, Marshall HG, Hatley OE, Hill RR (1984) Effect of management practices on grain yield, test weight, and lodging of soft red winter wheat 1. *Agron J* 76:379–383. <https://doi.org/10.2134/agronj1984.00021962007600030007x>
- Šarauskius E, Kazlauskas M, Naujokienė V, Bručienė I, Steponavičius D, Romaneckas K, Jasinskas A (2022) Variable rate seeding in precision agriculture: recent advances and future perspectives. *Agriculture* 12:305. <https://doi.org/10.3390/agriculture12020305>
- Schimmelpfennig D (2018) *Crop production costs, profits, and ecosystem stewardship with precision agriculture*. *J Agric Appl Econ* 50:81–103. <https://doi.org/10.1017/aae.2017.23>
- Schulte-Eickholt A (2009) *Erfassung, Analyse und Modellierung des Wurzelwachstums von Weizen (Triticum aestivum L.) unter Berücksichtigung der räumlichen Heterogenität der Pedosphäre*. Dissertation. Humboldt-Universität zu Berlin, Berlin
- Schultz BB (1985) Levene’s test for relative variation. *Syst Biol* 34:449–456. <https://doi.org/10.1093/sysbio/34.4.449>
- Šeremešić S, Čirić V, Djalović I, Vasin J, Zeremski T, Siddique KH, Farooq M (2020) Long-term winter wheat cropping

- influenced soil organic carbon pools in different aggregate fractions of Chernozem soil. *Arch Agron Soil Sci* 66:2055–2066. <https://doi.org/10.1080/03650340.2019.1711065>
- Sieling K, Stahl C, Winkelmann C, Christen O (2005) Growth and yield of winter wheat in the first 3 years of a monoculture under varying N fertilization in NW Germany. *Eur J Agron* 22:71–84. <https://doi.org/10.1016/j.eja.2003.12.004>
- Skakun S, Kalecinski NI, Brown MGL, Johnson DM, Vermote EF, Roger J-C, Franch B (2021) Assessing within-field corn and soybean yield variability from WorldView-3, Planet, Sentinel-2, and Landsat 8 Satellite Imagery. *Remote Sens* 13:872. <https://doi.org/10.3390/rs13050872>
- Slafer GA (2003) Genetic basis of yield as viewed from a crop physiologist's perspective. *Ann Appl Biol* 142:117–128. <https://doi.org/10.1111/j.1744-7348.2003.tb00237.x>
- Slafer GA, Rawson HM (1994) Sensitivity of wheat phasic development to major environmental factors: a re-examination of some assumptions made by physiologists and modellers. *Funct Plant Biol* 21:393. <https://doi.org/10.1071/pp9940393>
- Slafer GA, Savin R, Sadras VO (2014) Coarse and fine regulation of wheat yield components in response to genotype and environment. *Field Crop Res* 157:71–83. <https://doi.org/10.1016/j.fcr.2013.12.004>
- Späti K, Huber R, Finger R (2021) Benefits of increasing information accuracy in variable rate technologies. *Ecol Econ* 185:107047. <https://doi.org/10.1016/j.ecolecon.2021.107047>
- Stadler A, Rudolph S, Kupisch M, Langensiepen M, van der Kruk J, Ewert F (2015) Quantifying the effects of soil variability on crop growth using apparent soil electrical conductivity measurements. *Eur J Agron* 64:8–20. <https://doi.org/10.1016/j.eja.2014.12.004>
- Stein A, Brouwer J, Bouma J (1997) Methods for comparing spatial variability patterns of millet yield and soil data. *Soil Sci Soc Am J* 61:861. <https://doi.org/10.2136/sssaj1997.03615995006100030021x>
- Stewart CM, McBratney AB, Skerritt JH (2002) Site-specific durum wheat quality and its relationship to soil properties in a single field in Northern New South Wales. *Precision Agric* 3:155–168. <https://doi.org/10.1023/A:1013871519665>
- Taylor JC, Wood GA, Earl R, Godwin RJ (2003) Soil factors and their influence on within-field crop variability, part II: spatial analysis and determination of management zones. *Biosyst Eng* 84:441–453. [https://doi.org/10.1016/S1537-5110\(03\)00005-9](https://doi.org/10.1016/S1537-5110(03)00005-9)
- Tubaña BS, Arnall DB, Holtz SL, Solie JB, Girma K, Raun WR (2008) Effect of treating field spatial variability in winter wheat at different resolutions. *J Plant Nutr* 31:1975–1998. <https://doi.org/10.1080/01904160802403144>
- Usovicz B, Lipiec J (2017) Spatial variability of soil properties and cereal yield in a cultivated field on sandy soil. *Soil Tillage Res* 174:241–250. <https://doi.org/10.1016/j.still.2017.07.015>
- van Meirvenne M (2003) Is the soil variability within the small fields of Flanders structured enough to allow precision agriculture? *Precision Agric* 4:193–201. <https://doi.org/10.1023/A:1024561406780>
- Virto I, Barré P, Burlot A, Chenu C (2012) Carbon input differences as the main factor explaining the variability in soil organic C storage in no-tilled compared to inversion tilled agrosystems. *Biogeochemistry* 108:17–26. <https://doi.org/10.1007/s10533-011-9600-4>
- von Cossel M, Druecker H, Hartung E (2019) Low-input estimation of site-specific Lime demand based on apparent soil electrical conductivity and in situ determined topsoil pH. *Sensors* 19:5280. <https://doi.org/10.3390/s19235280>
- Wagner P, Marz M (2017) Precision Farming– Langzeitversuche mit Grunddüngungsstrategien. *Digitale Transformation–Wege in eine zukunftsfähige Landwirtschaft. LectureNotes in Informatics (LNI), Gesellschaft für Informatik, Bonn 2017, S.157–160*
- Wang Y-P, Shen Y (2015) Identifying and characterizing yield limiting soil factors with the aid of remote sensing and data mining techniques. *Precision Agric* 16:99–118. <https://doi.org/10.1007/s11119-014-9365-6>
- Watson CA, Atkinson D, Gosling P, Jackson LR, Rayns FW (2002) Managing soil fertility in organic farming systems. *Soil Use Manag* 18:239–247. <https://doi.org/10.1111/j.1475-2743.2002.tb00265.x>
- Welch BL (1951) On the comparison of several mean values: an alternative approach. *Biometrika* 38:330–336. <https://doi.org/10.1093/biomet/38.3-4.330>
- White RG, Kirkegaard JA (2010) The distribution and abundance of wheat roots in a dense, structured subsoil—implications for water uptake. *Plant Cell Environ* 33:133–148. <https://doi.org/10.1111/j.1365-3040.2009.02059.x>
- Wuest SB, Gollany HT (2013) Soil organic carbon and nitrogen after application of nine organic amendments. *Soil Sci Soc Am J* 77:237–245. <https://doi.org/10.2136/sssaj2012.0184>
- Yost JL, Hartemink AE (2019) Soil organic carbon in sandy soils: a review. Elsevier, Amsterdam, pp 217–310
- Yunusa IAM, Sedgley RH (1992) Reduced Tillering Spring Wheats for Heavy Textured Soils in a semi-arid Mediterranean Environment. *J Agron Crop Sci* 168:159–168. <https://doi.org/10.1111/j.1439-037X.1992.tb00994.x>
- Zebarth BJ, Fillmore S, Watts S, Barrett R, Comeau L-P (2021) Soil factors related to within-field yield variation in commercial potato fields in Prince Edward Island Canada. *Am J Potato Res* 98:139–148. <https://doi.org/10.1007/s12230-021-09825-4>
- Zhang L, He X, Liang Z, Zhang W, Zou C, Chen X (2020) Tiller development affected by nitrogen fertilization in a high-yielding wheat production system. *Crop Sci* 60:1034–1047. <https://doi.org/10.1002/csc2.20140>
- Zheng F, Qin J, Hua Y, Chu J, Dai X, He M (2023) Nitrogen uptake of winter wheat from different soil depths under a modified sowing pattern. *Plant Soil* 1–14. <https://doi.org/10.1007/s11104-023-05952-5>
- Zhou J, Zhang Z, Xin Y, Chen G, Wu Q, Liang X, Zhai Y (2022) Effects of planting density on root spatial and temporal distribution and yield of winter wheat. *Agronomy* 12:3014. <https://doi.org/10.3390/agronomy12123014>