Geometric speed limit of neutrino oscillation

Fazeel Khan¹ · Jerzy Dajka^{1,2,3}

Received: 8 December 2020 / Accepted: 10 May 2021 / Published online: 26 May 2021
© The Author(s) 2021

Abstract

We investigate geometric quantum speed limit of neutrino oscillations in a presence of matter and CP -violation. We show that periodicity in the speed limit present in an unperturbed system becomes damped by interaction with a normal matter and decoherence. We also show that (hypothetical) CP -violation causes enhancement of periodicity and increases amplitude of an oscillating quantum speed limit and can quantify CP -violation.

Keywords Neutrino oscillation · Quantum speed limit · Decoherence

1 Introduction

Time evolving neutrinos [35] and their oscillation focus and stimulate research from various perspectives. Dynamic properties of neutrino oscillations [10,11,44] initially studied in the most natural particle physics [35] context become equipped and extended by multitude of various and seemingly far investigations concerning decoherence [8,9,11,24,31,41,47] or various aspects of quantum information [5,6,10,18,33,49] to mention but a few. Such a broad interest seems to be at least partially motivated not only by natural applicability in the domain of particle physics methods and computational techniques borrowed from quantum information processing but also by recent attempts of information transfer as a resource utilizing neutrinos [52] or gravitational waves [1] reflecting an everlasting human dream of interstellar communication [25]. One can divide the quantum-information-based neutrino research into two overlap-

✉ Jerzy Dajka
jerzy.dajka@us.edu.pl

Fazeel Khan
ahmed.khan@smcebi.edu.pl

¹ Institute of Physics, University of Silesia in Katowice, Katowice, Poland

² Institute of Computer Science, University of Silesia in Katowice, Katowice, Poland

³ Silesian Center for Education and Interdisciplinary Research, University of Silesia in Katowice, Chorzow, Poland

ping classes: the first, focused on quantum properties of evolving neutrino—with an entanglement or contextuality as examples—and relating them to fundamental properties of these fascinating particles; the second, seemingly more exotic, motivated by neutrino-based quantum information processing. However, history of science shows that an effective feedback coupling between fundamentals and applications is a *sine qua non* condition of progress [22].

In this paper, we consider one of fundamental features of neutrino oscillation analysed as a quantum dynamical process and, after short review of neutrino time-evolution in a qutrit approximation, we discuss *geometric* quantum speed limit [29] of neutrino oscillation. Quantum speed limits (QSL) or quantum limit of evolution time reviewed in Ref. [28] can be studied from various context-dependent perspectives [7,21,27,30,34,55]: quantum control, quantum information or metrology. There were various motivations driving these investigations. The first approaches [42,43] were unified [40] and further developed in very various, sometimes divergent directions [28]. Some of them focus practical role of speed limit as a natural bound for control strategies, the other emphasize fundamental relation between various speed limits and intrinsic time scales of quantum systems [28]. In particular, contrary to recent operational approaches [21,53], geometric QSL proposed in Ref. [29] allows to identify a class of fundamental statistical and informational properties of quantum dynamics since it constitutes an upper bound for time-dependent changes of a shortest (geodesic) path qualified by the Fisher–Rao statistical distance [13] between states. With this interpretation geometric speed limit of Deffner and Lutz [29] allows to identify how a rate of information, given by a change of statistical distance, relates to dynamical properties of quantum dynamical systems governed by a very general class of equations of motion. Following [10,12], we neglect non-Markovian effects in neutrino time evolution and describe it in a qutrit approximation (i.e. as an effective qutrit using three dimensional flavour F space) in terms of Gorini–Kossakowski–Sudarshan–Lindblad Markovian master equation ($\hbar = 1$) [4,20]:

$$\frac{d}{dt}\rho_F(t) = \mathbb{L}[\rho_F(t)], \quad (1)$$

where $\rho_F(t) \in \mathcal{B}(\mathbb{C}^3)$ is a neutrino state. The generator $\mathbb{L}[\cdot]$ consisting of Hamiltonian and Lindbladian (decohering) parts is specified in the following section. For a time-evolving quantum system Eq. (1), its geometric QSL reads [29] as follows:

$$v_{\text{QSL}}(t) = \frac{\|\mathbb{L}[\rho_F(t)]\|}{2 \cos(l(t)) \sin(l(t))} \quad (2)$$

where the operator norm $\|\cdot\|$ is given by $\|\mathbb{L}\| = \sigma_{\max}$ with $\sigma_{\max}$ the largest *singular* value of $\mathbb{L}[\rho_F(t)]$ and the Bures distance

$$l(t) = \arccos(\sqrt{F(\rho_F(0), \rho_F(t))}) \quad (3)$$

related to fidelity F [13]. The notion of QSL is based on suitably defined bound of time derivative of the Fisher–Rao metric [29]. For pure initial states $\rho_F(0)$ Eq. (2)

is derived [28,29] using $2 \cos(l) \sin(l) \dot{l} \leq |\text{Tr}[\dot{\rho}_F(t) \rho_F(0)]| \leq \|\mathbb{L}[\rho_F(t)]\| = \sigma_{\max}$ where the operator norm in a last inequality provides sharpest bound in comparison with trace norm or Hilbert–Schmidt [28]. The equality which relates QSL to a largest singular value $\sigma_{\max}$ of $\mathbb{L}[\rho_F(t)]$ is for pure initial states $\rho_F(0)$ granted. Let us emphasize that QSL, which is a *time-dependent* quantity, characterizes quantum dynamics at a given instant contrary to other often very different operational approaches to quantum speed limits such as e.g. [53].

In this work, we investigate relations between properties of the neutrino QSL and conditions affecting neutrino oscillation: (i) interaction with normal matter' (electrons or neutrons) induced by forward elastic scattering (in Sect. 2), (ii) an effect of decoherence resulting in nonunitary damping of neutrino oscillation (in Sect. 3). Motivated by recent experiments [54] indicating possible violation of CP (charge conjugation parity) symmetry for neutrinos we investigate (iii) the effect of CP violation relating this fundamental feature to the geometric QSL in Sect. 4. We show that the CP -violation is a highly non-trivial feature affecting QSL of neutrino oscillation being a candidate for a hallmark of CP -violation in neutrino systems. Finally, we discuss and summarize our results.

2 Neutrino as a qutrit

As the discovery of the neutrino oscillations, originating from mixing of *three* neutrino fields, was a first evidence in favor of a beyond the standard model physics attracting attention of particle community [14]. Following it, neutrino studies from a perspective of quantum information and communication are naturally supported by an effective description of neutrino's flavour in terms of a qutrit approximation. It is known [35] that a neutrino of a given flavour $\alpha \in \{e, \mu, \tau\}$ evolves in time and can be measured in a different flavour state. This phenomenon is an essence of neutrino oscillation and originates from a fundamental non-correspondence between neutrino's flavour $\{\nu_e, \nu_\mu, \nu_\tau\}$ and massive states $\{\nu_1, \nu_2, \nu_3\}$. These two sets of orthogonal states are unitarily related by Pontecorvo–Maki–Nakagawa–Sakata mixing matrix U_{PMNS} :

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U_{\text{PMNS}} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}, \quad (4)$$

parameterized by three mixing angles θ_{12} , θ_{23} , θ_{13} and one CP -violating phase δ [35]:

$$U_{\text{PMNS}} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \quad (5)$$

with $c_{ij} = \cos(\theta_{ij})$ and $s_{ij} = \sin(\theta_{ij})$ (following Ref. [23] we set $s_{12}^2 = 0.307$, $s_{13}^2 = 0.021$ and $s_{23}^2 = 0.5$) and δ being the CP -violating phase. The phase δ according to the most recent experiments [54] can be non-vanishing indicating necessity

of reformulation of most fundamental aspects of the universe at (almost) all scales. The symmetry between matter and antimatter, charge-conjugation parity-reversal CP symmetry, seemed to be a perfect symmetry of nature. However, it was proposed by Sakharov in Ref. [51] that CP violation is one of the necessary conditions for known and observed imbalance of matter and anti-matter abundance in the universe. As the CP violation known for quarks is too small to be sufficient the CP symmetry and its potential violation remains a central topic of many investigations in particle physics. In particular, results of the above mentioned experiment [54] motivate searching consequences of the neutrinos' CP violation in various areas of natural science.

Neutrino oscillations are customarily approximated in a framework of unitary evolving closed quantum systems. However, as every physical system coupled to a matter, neutrino is also subjected to decoherence effects [10,12] originating from interaction with the environment and resulting in *non-unitary* corrections to its time-evolution [20]. Decoherence effects not only serve as a possible explanation of certain experimental data [8,9,31,32,41,47]—cf. Ref. [24] for a summary of a recent progress on that topic—but also significantly modify dynamic properties of the oscillation. It is since in a presence of decoherence any system becomes *open* and its evolution is determined by a reduced (with respect to its environment) density matrix [4]. Description of a time evolution of an open system cannot be arbitrary—complete positivity [4,12,20] together with a semi-group property are two of the best known requirements.

Neutrino oscillation in a qutrit approximation Eq. (1) is given by a solution of a master equation

$$\frac{d}{dt}\rho_F = -i[\mathcal{H}_F, \rho_F] + L[\rho_F], \quad (6)$$

where initially we assume neutrino in a particular flavour state $\rho_F(0) = |\Psi_F\rangle\langle\Psi_F|$ which in our case chosen to be electron, i.e. $|\Psi_F\rangle = |\nu_e\rangle$. Neutrino oscillation can be quantified in a most natural way by considering an overlap between a flavour state solving Eq. (6) and a state of well-defined flavour $\rho(0) = |\nu_e\rangle\langle\nu_e|$ initializing oscillation Eq. (6). The corresponding probability

$$P_0 = \text{Tr}[\rho_F(0)\rho_F(t)] \quad (7)$$

serves in the following discussion as natural reference (benchmark) for QSL dynamics which properties are compared. Let us notice that for a neutrino initially in a pure state $\rho_F(0)$, the overlap P_0 in Eq. (7) is related to the fidelity of states [13]. The terms $[\mathcal{H}_F, \cdot]$ and $L[\cdot]$ are responsible for unitary (Hamiltonian generated) and dissipative (Lindbladian) part of time evolution Eq. (6), respectively. The Hamiltonian $\mathcal{H}_F$ [19, 26,35,50]:

$$\mathcal{H}_F = H_{\text{kin}} + H_{\text{pot}}, \quad (8)$$

includes the kinetic H_{kin} and the potential H_{pot} part, the latter resulting from neutrino interaction with normal matter. The Lindbladian L in Eq. (6) describes decoherence and a discussion of its influence on the QSL is postponed to the next Sections. In the absence of decoherence the time evolution of the neutrino, Eq. (6) reduces to the Schrödinger equation. Operating at an ultrarelativistic limit for the neutrinos (with neutrino's eigenenergies $E_k = E + \frac{m_k^2}{2E}$ and momentum $p \sim E$)—the kinetic part of

the Hamiltonian $\mathcal{H}_{\mathcal{F}}$ in the flavour basis reads:

$$H_{\text{kin}} = \frac{1}{2E} U_{\text{PMNS}} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \Delta m_{21}^2 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} U_{\text{PMNS}}^\dagger, \quad (9)$$

where $\Delta m_{ij}^2 = m_i^2 - m_j^2$ is the difference between the masses of two oscillating neutrinos. In the following numerical calculations, we set $\Delta m_{12}^2 = 7.37 \cdot 10^{-5} \text{ eV}^2$ and $\Delta m_{31}^2 = 2.52 \cdot 10^{-3} \text{ eV}^2$ [23]. The main contribution to the potential part H_{pot} originates from neutrino's interaction due to coherent forward elastic scattering (and for negligible non-coherent effects) with the matter's electrons or neutrons generating *charged-current potential* V_{CC} . The potential part of the Hamiltonian takes then the following form:

$$H_{\text{pot}} = \begin{pmatrix} V_C & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (10)$$

The charged-current potential V_C is related to the electron's density in the matter $V_C = \sqrt{2} G_F n_e$ with the Fermi coupling constant G_F and the electron number density n_e .

3 Results

Quantum speed limit of neutrino oscillation is analysed by means of numerical solution of Eq. (6) obtained using QuTip [38,39], a Python-based toolbox dedicated to dynamics and control of open quantum systems. We present results in three steps: first including normal matter, the second including decoherence effects and finally presenting our central results concerning QSL for CP -violating system. We investigate numerical solution of Eq. (6) with Hamiltonian and Lindbladian parts given in Eq. (8) and (introduced below) Eq. (11), respectively. We utilize QuTiP master equation solver `qutip.mesolve` (both for unitary and dissipative evolution) for a neutrino initially in the electron flavour. In calculation of quantum speed limit Eq. (2), we utilize `qutip.bures_angle` for Bures distance and `scipy.linalg.svdvals` to compute singular values of $\mathbb{L}[\rho_F(t)]$ in Eq. (2).

3.1 Normal matter

The charged-current potential V_C is the first quantity which significantly affects neutrino oscillation modifying its period as presented in an upper panel of Fig. 1. The modification is present also in the geometric QSL as presented in lower panel of Fig. 1. Let us notice that in an absence of normal matter i.e. for $V_C = 0$ maximal values $P_0 = 1$ correspond to *discontinuities* of $\nu_{\text{QSL}} \rightarrow \infty$ what is in agreement with interpretation of geometric QSL as a speed of changes of statistical distance between states [28,29]. Due to the discontinuities (vertical asymptotes) a range of ν_{QSL} for $V_C = 0$ presented in Fig. 1 is truncated. Increase in V_C results in changes of periodicity of oscillation and,

Fig. 1 Upper panel: decoherence-free neutrino oscillation qualified by P_0 Eq. (7) as a function of time for various values of the charge-current potential V_C . Lower panel: geometric quantum speed limit of decoherence-free neutrino oscillation ν_{QSL} Eq. (2) (for clarity of presentation truncated to $\nu_{QSL} < 4$) as a function of time for various values of the charge-current potential V_C . The CP violating phase $\delta = 0$

as a consequence of interaction with normal matter, discontinuities in ν_{QSL} disappear cf. Fig. 1 replaced by sharp but continuous peaks. As in the absence of decoherence quantum finite-dimensional systems are periodic or quasi-periodic, one obtains large values of ν_{QSL} provided that $P_0 \approx 1$, i.e. the time-evolving systems returns in a neighbourhood of its starting point. Inclusion of interaction with matter allows to abandon an artefact of discontinuous QSL time characteristics clearly absent in real systems.

3.2 Decoherence

One of peculiarities of neutrino as a fundamental particle is its extremely weak interaction with almost everything existing in the Universe. However, omnipresent decoherence is unavoidable and can serve as one of possible mechanisms [8,9,11,24,31,41,47] affecting unusual properties of a neutrino. We consider the simplest Markovian model

of decoherence based on natural requirement of complete positivity of time evolution [4] leading to a Master equation Eq. (6) with a (non-Hamiltonian) Lindbladian part

$$L[\rho_F] = \sum_{i,j=0}^{N^2-1} c_{ij} \left(F_i \rho_F F_j^\dagger - \frac{1}{2} \{ F_j^\dagger F_i, \rho_F \} \right) \quad (11)$$

being responsible for the non-standard effects connected with dissipation and decoherence. Here, $N = 3$ denotes the dimension of the system, $\{\cdot, \cdot\}$ is an anti-commutator, the matrices F_n stand for the generators of $SU(3)$. In the standard representation, they are given by the celebrated Gell-Mann matrices $F_n = \frac{\lambda_n}{2}$, where λ_n [37]:

$$\begin{aligned} \lambda_0 &= \mathbb{I}_3, \quad \lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ \lambda_3 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\ \lambda_5 &= \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \quad \lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \\ \lambda_7 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}. \end{aligned} \quad (12)$$

The coefficients c_{ij} in Eq. (11) satisfy the set of inequalities granting complete positivity [11,12]):

$$|c_{ij}| \leq \frac{1}{2}(c_{ii} + c_{jj}). \quad (13)$$

In our analysis, we make one more simplifying step and we limit our consideration to diagonal decoherence with $c_{ii} = \kappa$ and $c_{i \neq j} = 0$. In Fig. 2, we present simultaneous effect of decoherence and very weak interaction with normal matter with $V_C = 0.03$ in order to avoid artificial discontinuities in ν_{QSL} present in ideal noise-less systems. Let us notice that a range of ν_{QSL} presented in Fig. 2 for a noise-less case, $\kappa = 0$ is (as irrelevant for further discussion) truncated. Let us also notice initially asymptotic value of ν_{QSL} as $t \rightarrow 0^+$ due to a definition (and its geometric interpretation) of the geometric speed limit Eq. (2). Markovian decoherence, contrary to previously discussed interaction with normal matter, does not influence periodicity of neutrino oscillation but rather effects in amplitude damping as it can be inferred from an upper panel of Fig. 2. Non-vanishing decoherence removes discontinuities present in a noise-less time evolution of ν_{QSL} but does not affect position of peaks since decoherence of any amplitude κ does not alter positions of extreme values in QSL compared to the noise-less oscillations. For sufficiently large values of κ , there is a qualitative change in the QSL characteristics which, as it is presented in the lower panel of Fig. 2, starts to be monotonically decreasing function of time.

Fig. 2 Neutrino oscillation quantified by P_0 Eq. (7) (upper panel) and neutrino QSL quantified by ν_{QSL} Eq. (2) for clarity of presentation truncated to $\nu_{\text{QSL}} < 0.5$ (lower panel) for different amplitudes of decoherence $c_{ii} = \kappa$ in a presence of normal matter $V_C = 0.03$ and no CP -violation $\delta = 0$

3.3 CP -violation

An influence of the CP symmetry and its potential violation is visible both in decoherence-affected neutrino oscillation and geometric QSL as presented in upper and (respectively) lower panel of Fig. 3. For a better visualization of results, a range of ν_{QSL} is in the lower panel of Fig. 3 truncated. Let us emphasize the presence of initially asymptotic value of ν_{QSL} as $t \rightarrow 0^+$. Contrary to previously discussed cases of modification via decoherence only affecting height of P_0 graph, non-vanishing CP phase δ results in increase in P_0 depth which becomes maximal for $\delta = \pi$. In other words, neutrino oscillates with the same (quasi)period but in a presence of CP -violation it wanders off further (in a sense of decreasing fidelity) than it does for $\delta = 0$. Resulting behaviour of ν_{QSL} suggests possibility of using geometric QSL to quantify the CP -violation if one utilizes relation between amplitude of ν_{QSL} and a value of δ . However, let us emphasize an existence of time windows where both P_0 and ν_{QSL} become indistinguishable with respect to different δ limiting potential

Fig. 3 Neutrino oscillation quantified by P_0 Eq. (7) (upper panel) and neutrino QSL quantified by ν_{QSL} Eq. (2) for clarity of presentation truncated to $\nu_{\text{QSL}} < 0.4$ (lower panel) for different values of CP -violating phase δ in the presence of decoherence with $\kappa = 0.03$ in Eq. (11) and in the presence of normal matter $V_C = 0.03$

applicability of QSL as a hallmark of violation of the CP symmetry. Let us notice that the effect of CP -symmetry is even stronger if one considers ν_{QSL} cf. lower panel of Fig. 3. First, the non-vanishing δ enhances peaks of which otherwise are damped by decoherence. As a result, geometric QSL quantifies a potential CP violation provided that we remember the above mentioned time windows where both P_0 and ν_{QSL} for all values of δ are identical. It occurs close to local maxima of P_0 , i.e. at time instance when the neutrino returns close to its initial state.

4 Discussion

Neutrinos belong to a class of objects allowing for 'complementary' studies. The most common approaches of particle physics community become often success-

fully augmented by results concerning quantum informational properties studied from seemingly different perspective.

Our work belongs to a part of investigations indicating possibility of utilizing quantum informational perspective for better understanding the nature of neutrino. We considered geometric quantum speed limit and we showed its potential applicability as a hallmark for the CP -violation. First, we presented how QSL as a function of time becomes modified by interaction of neutrino with normal matter formed by matter's electrons or neutrons and generating the charge-current correction V_C in the Hamiltonian Eq. (8). Removal of artificial discontinuities in the QSL-time characteristics is the main effect of this interaction. Infinite peaks in ν_{QSL} typical for an ideal decoherence-free system become shifted and lowered by V_C and, for sufficiently large values of V_C lose their direct connection to the position of extremal values of P_0 Eq. (7). These results after inclusion of decoherence effect become supplemented by damping of oscillations of ν_{QSL} which, for sufficiently high decoherence (given by an amplitude κ of Lindblad dissipators Eq. (11)). Although we used purely phenomenological model of decoherence guided solely by the requirement of complete positivity, the results are generic for Markovian open quantum systems.

The most significant results allow to relate geometric QSL to a very hot topic of hypothetical violation of the CP symmetry for neutrinos [54]. We infer that such a violation indicated by non-vanishing δ in the Pontecorvo–Maki–Nakagawa–Sakata mixing matrix Eq. (5) results in an enhancement of ν_{QSL} amplitude and effective separation of time characteristics of geometric QSL of systems with different δ . We also identified time windows of geometric QSL where this effect becomes either particularly visible or vanishes making time characteristics almost δ -independent.

Geometric quantum speed limit Eq. (2) quantifies informational (statistical) content of a time-local (calculated at a given time instant t) rate of change of neutrino state. There are three factors affecting this rate which are studied in this paper: interaction between neutrino and normal matter (for $V_C \neq 0$), Markovian decoherence (for $\kappa \neq 0$ in Eq. (11) and potential violation of the CP symmetry (for $\delta \neq 0$ in Eq. (5)). From a mathematical perspective, the changes in the geometric speed limit Eq. (2) reported here are due to a collective effect of collaboration of these three factors. In particular, the effect of δ in ν_{QSL} starts to be visible provided that one allows for the presence of decoherence in a dynamical model of neutrino oscillation. It is due to an effective cancellation of the CP -violating phase both for the operator norm and the Bures distance $l(t)$ (related to fidelity and for pure states given by their overlap) in the numerator and denominator of Eq. (2), respectively, for conservative model of neutrino oscillation with $\kappa = 0$. The numerator of QSL Eq. (2) is given by maximal singular value which, to be calculated, demands multiplication of an operator and its conjugate [28] resulting in the above-mentioned cancellation of phase. In the presence of decoherence resulting in an information loss due to non-unitary dynamics Eq. (11), the effect of δ is present indicating limiting applicability of purely Hamiltonian models of a description of at least certain properties of neutrino oscillation in time. The QSL quantifying instantaneous rate of changes of neutrino state under non-unitary evolution is a quantity which allows to exhibit otherwise 'invisible' effects related to characteristic time scale of neutrino dynamics. The phenomenon of neutrino oscillations in time and for decoherence-free Hamiltonian description is related

Fig. 4 Quantum speed limit time τ_{QSL} Eq. (14) as a function of time: (i) upper panel: for various values of the charge-current potential V_C in the absence of decoherence $\kappa = 0$ and with CP violating phase $\delta = 0$. (ii) Central panel: for different amplitudes of decoherence $c_{ij} = \kappa$ in the presence of normal matter $V_C = 0.03$ and no CP -violation $\delta = 0$. (iii) Bottom panel: for different values of CP -violating phase δ in the presence of decoherence with $\kappa = 0.03$ in Eq. (11) and in a presence of normal matter $V_C = 0.03$

to the Mandelstam–Tamm time-energy uncertainty relation [16]. Despite of the presence of certain controversial issues [3, 15], it allows to set a characteristic time interval required for a significant change of the flavour neutrino state, i.e. an intrinsic time scale of the inter-flavour passage [16]. Let us emphasize that in particular the Mössbauer neutrinos produced in two-body decays of nuclei embedded in a crystal lattice [2] can be utilized for justification and verification of this property [17]. Working beyond conservative dynamics approximation, as we do here, one can utilize QSL as a natural generalization of a (local in time) bound for a rate of neutrino oscillation. Moreover, the geometric quantum speed limit Eq. (2) is directly associated with a quantum speed limit time [28]

$$\tau_{\text{QSL}}^{-1}(t) = \frac{1}{t} \int_0^t v_{\text{QSL}}(t') dt' \quad (14)$$

which thereafter can serve as a generalization of the characteristic time interval for a significant change of the flavour neutrino state in a very general, possibly non-Hamiltonian, systems. The quantum speed limit time Eq. (14) is directly related to geometric quantum speed limit QSL Eq. (2) and, by extension, it reflects the results reported in this work. The quantum speed limit time τ_{QSL} is presented in Fig. 4 for the same set of parameters as previously v_{QSL} . Let notice that although the results are qualitative only, one recognizes characteristic features of geometric speed limit Eq. (2) also present in the speed limit time Eq. (14). Following the interpretation given in Refs. [16, 17] of the time-energy uncertainty and a related time scale as being characteristic for duration of a neutrino inter-flavour passage, one observes both an interaction with normal matter of an amplitude V_C and Markovian decoherence quantified by κ in the model Eq. (11) significantly modifying the quantum speed limit time Eq. (14). In particular, the effect of V_C (as presented in the upper panel of Fig. 4 influences both a magnitude of τ_{QSL} and its periodicity as it was for geometric speed limit Eq. (2) in Fig. 1. Similar modification yet applied to an amplitude only is shared by geometric speed limit and its corresponding speed limit time under Markovian decoherence cf. Fig. 2 and the central panel of Fig. 4, respectively. The effect of CP violation, seemingly less 'spectacular', is also present (bottom panel of Fig. 4 indicating an enhancement of the quantum speed limit time due to an increasing value of the CP -violating phase δ). We strongly emphasize that all the neutrino experiments such as the recent one reported in Ref. [54]) belong to the most sophisticated experiments performed so far and our predictions concerning QSL and the corresponding time are nothing but qualitative signatures of certain properties of neutrino oscillation.

5 Conclusions

Quantum information processing models [46] such as circuit model [46], topological models, Zidan's model of quantum computing proposed in Refs. [56, 57] or quantum communication protocols [36, 45, 48] can—in principle—be realized using neutrinos [52]. However, despite recent experiments [52], such implementations, together with models using gravitational waves [1] as resource, still remain in a domain of science

fiction [25] rather than reality. Nevertheless, investigations of fundamental properties of neutrino with an emphasis given on CP -violation [51] inspire researchers of various branches of science. In our work, we provided one more quantifier, the geometric QSL, of dynamic properties of neutrino oscillations. Formal similarity of dynamical model governing neutrino oscillation and qutrit dynamics, which we utilize, is a starting point of many investigations concerning properties of neutrino conducted from a perspective of quantum information. However, expecting all the predictions of quantum information to be directly applicable or translatable to a language of neutrino physics can be due to their redundancy unjustifiable. Dynamics of neutrino oscillation even simplified to a qutrit model remains confined and limited by our (current) knowledge of neutrino's physics. As we currently know, neutrinos are massive and are affected—via charged-current potential V_{CC} , cf. Eq. (8)—by normal matter. We also expect that neutrinos undergo decoherence Eq. (11) which (for hardly interacting neutrinos) can safely be assumed Markovian. Moreover, most recent experiments [54] remove at least a part of these limitations due to a possible non-preservation of the CP symmetry. Extension of the domain of studies to geometric quantum speed limit Eq. (2) of CP -violating oscillation models in a presence of decoherence and normal matter, allowed to identify regimes where the QSL, being essentially dynamical property, becomes modified by changes of equations of motion Eq. (6) induced by all the three above mentioned factors, i.e. normal matter, decoherence and CP -violation. We recognized tailored range of time where the effect of modifications becomes most significant not only indicating an opportunity of using QSL as a diagnostic tool for neutrino properties, e.g. as a hallmark of the CP violation, but also expressing quantum informational content of neutrino oscillation in terms of its geometric quantum speed limit.

Declaration

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Abramowicz, M., Bejger, M.,ourgoulhon, E., Straub, O.: A galactic centre gravitational-wave messenger. *Sci. Rep.* **10**, 7054 (2020). <https://doi.org/10.1038/s41598-020-63206-1>
2. Akhmedov, E.K., Kopp, J., Lindner, M.: Oscillations of Mössbauer neutrinos. *J. High Energy Phys.* **2008**(05), 005 (2008). <https://doi.org/10.1088/1126-6708/2008/05/005>

3. Akhmedov, E.K., Kopp, J., Lindner, M.: Comment on ‘Time-energy uncertainty relations for neutrino oscillations and the Mössbauer neutrino experiment’. *J. Phys. G Nucl. Part. Phys.* **36**, 078001 (2009). <https://doi.org/10.1088/0954-3899/36/7/078001>
4. Alicki, R., Lendi, K.: *Quantum Dynamical Semigroups and Applications*. Springer, Berlin (2007)
5. Alok, A.K., Banerjee, S., Sankar, S.U.: Quantum correlations in terms of neutrino oscillation probabilities. *Nucl. Phys. B* **909**, 65–72 (2016). <https://doi.org/10.1016/j.nuclphysb.2016.05.001>
6. Antonio, C., Salvatore, M.G., Gaetano, L., Quaranta, A.: Discerning the nature of neutrinos: decoherence and geometric phases. *Universe* **6**(11), 207 (2020). <https://doi.org/10.3390/universe6110207>
7. Awasthi, N., Haseli, S., Johri, U.C., Salimi, S., Dolatkah, H., Khorashad, A.S.: Quantum speed limit time for correlated quantum channel. *Quantum Inf. Process.* **19**, 308 (2020). <https://doi.org/10.1007/s11128-020-02807-1>
8. Bakhti, P., Farzan, Y., Schwetz, T.: Revisiting the quantum decoherence scenario as an explanation for the LSND anomaly. *J. High Energy Phys.* **2015**(5), 7 (2015). [https://doi.org/10.1007/JHEP05\(2015\)007](https://doi.org/10.1007/JHEP05(2015)007)
9. Balieiro Gomes, G., Guzzo, M.M., de Holanda, P.C., Oliveira, R.L.N.: Parameter limits for neutrino oscillation with decoherence in Kamland. *Phys. Rev. D* **95**, 113005 (2017). <https://doi.org/10.1103/PhysRevD.95.113005>
10. Banerjee, S., Alok, A.K., Srikanth, R., Hiesmayr, B.C.: A quantum information theoretic analysis of three flavor neutrino oscillations. *Eur. Phys. J. C* **75**(10), 487 (2015)
11. Benatti, F., Floreanini, R.: Massless neutrino oscillations. *Phys. Rev. D* **64**, 085015 (2001)
12. Benatti, F., Floreanini, R.: Open quantum dynamics: complete positivity and entanglement. *Int. J. Modern Phys. B* **19**(19), 3063–3139 (2005). <https://doi.org/10.1142/S0217979205032097>
13. Bengtsson, I., Życzkowski, K.: *Geometry of Quantum States*. Cambridge University Press, Cambridge (2006)
14. Bilenky, S.: Neutrino oscillations: from a historical perspective to the present status. *Nucl. Phys. B* **908**, 2–13 (2016). <https://doi.org/10.1016/j.nuclphysb.2016.01.025>. Neutrino Oscillations: Celebrating the Nobel Prize in Physics 2015
15. Bilenky, S.M., von Feilitzsch, F., Potzel, W.: Reply to the comment on ‘Time-energy uncertainty relations for neutrino oscillations and the Mössbauer neutrino experiment’ by E.K. Akhmedov, J. Kopp and M. Lindner. *J. Phys. G Nucl. Part. Phys.* **36**, 078002 (2009). <https://doi.org/10.1088/0954-3899/36/7/078002>
16. Bilenky, S.M., von Feilitzsch, F., Potzel, W.: Time-energy uncertainty relations for neutrino oscillations and the Mössbauer neutrino experiment. *J. Phys. G Nucl. Part. Phys.* **35**, 095003 (2008). <https://doi.org/10.1088/0954-3899/35/9/095003>
17. Bilenky, S.M., von Feilitzsch, F., Potzel, W.: Neutrino oscillations and uncertainty relations. *J. Phys. G Nucl. Part. Phys.* **38**, 115002 (2011). <https://doi.org/10.1088/0954-3899/38/11/115002>
18. Blasone, M., Dell’Anno, F., Siena, S.D., Illuminati, F.: Entanglement in neutrino oscillations. *EPL Europhys. Lett.* **85**(5), 50002 (2009)
19. Blennow, M., Smirnov, A.Y.: Neutrino propagation in matter. *Adv. High Energy Phys.* **2013**, 972485 (2013)
20. Breuer, H.P., Petruccione, F.: *The Theory of Open Quantum Systems*. Oxford University Press, Oxford (2002)
21. Brody, D.C., Longstaff, B.: Evolution speed of open quantum dynamics. *Phys. Rev. Res.* **1**, 033127 (2019). <https://doi.org/10.1103/PhysRevResearch.1.033127>
22. Buchwald, J.Z., Fox, R.: *The Oxford Handbook of the History of Physics*, Oxford Handbooks. Oxford University Press, Oxford (2014)
23. Capozzi, F., Di Valentino, E., Lisi, E., Marrone, A., Melchiorri, A., Palazzo, A.: Global constraints on absolute neutrino masses and their ordering. *Phys. Rev. D* **95**, 096014 (2017). <https://doi.org/10.1103/PhysRevD.95.096014>
24. Carpio, J., Massoni, E., Gago, A.M.: Revisiting quantum decoherence in the matter neutrino oscillation framework. [arXiv:1711.03680](https://arxiv.org/abs/1711.03680) (2017)
25. Cixin, L.: *The Dark Forest (Remembrance of Earth’s Past, 2)*. Tom Doherty Associates, LLC, New York (2008)
26. Dajka, J., Syska, J., Luczka, J.: Geometric phase of neutrino propagating through dissipative matter. *Phys. Rev. D* **83**, 097302 (2011)
27. Deffner, S.: Quantum speed limits and the maximal rate of information production. *Phys. Rev. Res.* **2**, 013161 (2020). <https://doi.org/10.1103/PhysRevResearch.2.013161>

28. Deffner, S., Campbell, S.: Quantum speed limits: from Heisenberg's uncertainty principle to optimal quantum control. *J. Phys. A Math. Theor.* **50**(45), 453001 (2017). <https://doi.org/10.1088/1751-8121/aa86c6>
29. Deffner, S., Lutz, E.: Quantum speed limit for non-Markovian dynamics. *Phys. Rev. Lett.* **111**, 010402 (2013). <https://doi.org/10.1103/PhysRevLett.111.010402>
30. Dehdashti, S., Yasar, F., Harouni, M.B., Mahdifar, A.: Quantum speed limit in the thermal spin-boson system with and without tunneling term. *Quantum Inf. Process.* **19**, 308 (2020). <https://doi.org/10.1007/s11128-020-02807-1>
31. Gago, A.M., Santos, E.M., Teves, W.J.C., Zukanovich Funchal, R.: Quantum dissipative effects and neutrinos: current constraints and future perspectives. *Phys. Rev. D* **63**, 073001 (2001). <https://doi.org/10.1103/PhysRevD.63.073001>
32. Gago, A.M., Santos, E.M., Teves, W.J.C., Zukanovich Funchal, R.: A study on quantum decoherence phenomena with three generations of neutrinos. [arxiv:hep-ph/0208166v1](https://arxiv.org/abs/hep-ph/0208166v1) (2002)
33. Gangopadhyay, D., Home, D., Roy, A.S.: Probing the Leggett–Garg inequality for oscillating neutral kaons and neutrinos. *Phys. Rev. A* **88**(2), 022115 (2013)
34. García-Pintos, L.P., del Campo, A.: Quantum speed limits under continuous quantum measurements. *New J. Phys.* **21**(3), 033012 (2019). <https://doi.org/10.1088/1367-2630/ab099e>
35. Giunti, C., Wook, K.C.: *Fundamentals of Neutrino Physics and Astrophysics*. Oxford University Press, Oxford (2007)
36. Gordon, G., Rigolin, G.: Generalized teleportation protocol. *Phys. Rev. A* **73**, 042309 (2006). <https://doi.org/10.1103/PhysRevA.73.042309>
37. Humphreys, J.: *A Course in Group Theory*. Oxford University Press, Oxford (1996)
38. Johansson, J., Nation, P., Nori, F.: QuTiP: an open-source python framework for the dynamics of open quantum systems. *Comput. Phys. Commun.* **183**(8), 1760–1772 (2012). <https://doi.org/10.1016/j.cpc.2012.02.021>
39. Johansson, J., Nation, P., Nori, F.: QuTiP 2: a python framework for the dynamics of open quantum systems. *Comput. Phys. Commun.* **184**(4), 1234–1240 (2013). <https://doi.org/10.1016/j.cpc.2012.11.019>
40. Levitin, L.B., Toffoli, T.: Fundamental limit on the rate of quantum dynamics: the unified bound is tight. *Phys. Rev. Lett.* **103**, 160502 (2009). <https://doi.org/10.1103/PhysRevLett.103.160502>
41. Lisi, E., Marrone, A., Montanino, D.: Probing possible decoherence effects in atmospheric neutrino oscillations. *Phys. Rev. Lett.* **85**, 1166–1169 (2000). <https://doi.org/10.1103/PhysRevLett.85.1166>
42. Mandelstam, L., Tamm, I.: The Uncertainty Relation Between Energy and Time in Non-relativistic Quantum Mechanics, pp. 115–123. Springer, Berlin (1991). https://doi.org/10.1007/978-3-642-74626-0_8
43. Margolus, N., Levitin, L.B.: The maximum speed of dynamical evolution. *Physica D Nonlinear Phenomena* **120**(1), 188–195 (1998). [https://doi.org/10.1016/S0167-2789\(98\)00054-2](https://doi.org/10.1016/S0167-2789(98)00054-2). Proceedings of the Fourth Workshop on Physics and Consumption
44. Molfetta, G.D., Pérez, A.: Quantum walks as simulators of neutrino oscillations in a vacuum and matter. *New J. Phys.* **18**(10), 103038 (2016). <https://doi.org/10.1088/1367-2630/18/10/103038>
45. Nguyen, D.M., Kim, S.: Quantum key distribution protocol based on modified generalization of Deutsch–Jozsa algorithm in d-level quantum system. *Int. J. Theor. Phys.* **58**(1), 71–82 (2019). <https://doi.org/10.1007/s10773-018-3910-4>
46. Nielsen, M.A., Chuang, I.L.: *Quantum Computation and Quantum Information*. Cambridge University Press, Cambridge (2010)
47. Oliveira, R.L.N., Guzzo, M.M.: Quantum dissipation in vacuum neutrino oscillation. *Eur. Phys. J. C* **69**, 493–502 (2010). <https://doi.org/10.1140/epjc/s10052-010-1388-1>
48. Ren, J.G., Xu, P., Yong, H.L., Zhang, L., Liao, S.K., Yin, J., Liu, W.Y., Cai, W.Q., Yang, M., Li, L., Yang, K.X., Han, X., Yao, Y.Q., Li, J., Wu, H.Y., Wan, S., Liu, L., Liu, D.Q., Kuang, Y.W., He, Z.P., Shang, P., Guo, C., Zheng, R.H., Tian, K., Zhu, Z.C., Liu, N.L., Lu, C.Y., Shu, R., Chen, Y.A., Peng, C.Z., Wang, J.Y., Pan, J.W.: Ground-to-satellite quantum teleportation. *Nature* **549**(7670), 70–73 (2017). <https://doi.org/10.1038/nature23675>
49. Richter, M., Dziewit, B., Dajka, J.: Leggett–Garg k_3 quantity discriminates between Dirac and Majorana neutrinos. *Phys. Rev D* **96**, 076008 (2017)
50. Richter-Laskowska, M., Lobejko, M., Dajka, J.: Quantum contextuality of a single neutrino under interactions with matter. *New J. Phys.* **20**(6), 063040 (2018). <https://doi.org/10.1088/1367-2630/aacbf9f>

51. Sakharov, A.D.: Violation of CP-invariance, C-asymmetry, and Baryon asymmetry of the universe. *Soviet Phys. Uspekhi* **34**(5), 392–393 (1991). <https://doi.org/10.1070/pu1991v034n05abeh002497>
52. Sanci, D.D., et al.: Demonstration of communication using neutrinos. *Modern Phys. Lett. A* **27**(12), 1250077 (2012). <https://doi.org/10.1142/S0217732312500770>
53. Shao, Y., Liu, B., Zhang, M., Yuan, H., Liu, J.: Operational definition of a quantum speed limit. *Phys. Rev. Res.* **2**, 023299 (2020). <https://doi.org/10.1103/PhysRevResearch.2.023299>
54. The T2K Collaboration: Constraint on the matter–antimatter symmetry-violating phase in neutrino oscillations. *Nature* **580**(7803), 339–344 (2020). <https://doi.org/10.1038/s41586-020-2177-0>
55. Wu, S., Yu, C.: Quantum speed limit based on the bound of Bures angle. *Sci. Rep.* **10**, 5500 (2020). <https://doi.org/10.1038/s41598-020-62409-w>
56. Zidan, M.: A novel quantum computing model based on entanglement degree. *Modern Phys. Lett. B* **34**(35), 2050401 (2020). <https://doi.org/10.1142/S0217984920504011>
57. Zidan, M., Eleuch, H., Abdel-Aty, M.: Non-classical computing problems: toward novel type of quantum computing problems. *Results Phys.* **21**, 103536 (2021). <https://doi.org/10.1016/j.rinp.2020.103536>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.