On the exponential Diophantine equation related to powers of two consecutive terms of Lucas sequences

Mahadi Ddamulira^{1,2} · Florian Luca^{3,4,5}

Received: 28 January 2020 / Accepted: 30 March 2020 / Published online: 23 July 2020
© The Author(s) 2020

Abstract

Let $r \geq 1$ be an integer and $\mathbf{U} := (U_n)_{n \geq 0}$ be the Lucas sequence given by $U_0 = 0$, $U_1 = 1$, and $U_{n+2} = rU_{n+1} + U_n$, for all $n \geq 0$. In this paper, we show that there are no positive integers $r \geq 3$, $x \neq 2$, $n \geq 1$ such that $U_n^x + U_{n+1}^x$ is a member of $\mathbf{U}$.

Keywords Lucas sequences · Linear forms in logarithms · Baker’s method

Mathematics Subject Classification 11B39 · 11D61 · 11J86

1 Introduction

Let $r \geq 1$ be an integer and $\mathbf{U} := (U_n)_{n \geq 0}$ be the Lucas sequence given by $U_0 = 0$, $U_1 = 1$, and

$$U_{n+2} = rU_{n+1} + U_n \quad (1)$$

Open access funding provided by Austrian Science Fund (FWF). M. D. was supported by the FWF Projects: F5510-N26—Part of the special research program (SFB), “Quasi Monte Carlo Methods: Theory and Applications” and W1230—“Doctoral Program Discrete Mathematics”. F. L. was supported by Grant RTNUM19 from CoEMaSS, Wits, South Africa.

✉ Mahadi Ddamulira
mddamulira@tugraz.at; mahadi@aims.edu.gh
Florian Luca
Florian.Luca@wits.ac.za

¹ Institute of Analysis and Number Theory, Graz University of Technology, Kopernikusgasse 24/II, 8010 Graz, Austria

² Max Planck Institute for Mathematics, Bonn, Germany

³ School of Mathematics, University of the Witwatersrand, Johannesburg, South Africa

⁴ Research Group in Algebraic Structures and Applications, King Abdulaziz University, Jeddah, Saudi Arabia

⁵ Centro de Ciencias Matemáticas UNAM, Morelia, Mexico

for all $n \geq 0$. When $r = 1$ $\mathbf{U}$ coincides with the Fibonacci sequence while when $r = 2$ $\mathbf{U}$ coincides with the Pell sequence. It is well known that

$$U_n^2 + U_{n+1}^2 = U_{2n+1} \quad \text{for all } n \geq 0. \quad (2)$$

In particular, the identity (2) tells us that the sum of the squares of two consecutive terms of $\mathbf{U}$ is also a term of $\mathbf{U}$. When $r = 1$, we even have $U_n + U_{n+1} = U_{n+2}$ for all $n \geq 0$ since $\mathbf{U}$ is the Fibonacci sequence. We thus consider the Diophantine equation

$$U_n^x + U_{n+1}^x = U_m \quad (3)$$

in nonnegative integers (n, m, x) which by Eq. (2) has the parametric solution $m = 2n + 1$ when $x = 2$ for any $r \geq 1$ and even the parametric solution $m = n + 2$ when $x = 1$ if $r = 1$. For $r = 1$ Luca and Oyono [8] proved that Eq. (3) has no positive integer solutions (n, m, x) with $n \geq 2$ and $x \geq 3$. Rihane et al. [11] studied Eq. (3) when $r = 2$ and proved that there is no positive integer solution (n, m, x) of it with $x \neq 2$. In the same spirit, Gómez Ruiz and Luca [5] studied Eq. (3) with $\mathbf{U} = \{F_n^{(k)}\}_{n \geq -(k-2)}$, which is the k -generalised Fibonacci sequence of recurrence

$$F_n^{(k)} = F_{n-1}^{(k)} + F_{n-2}^{(k)} + \cdots + F_{n-k}^{(k)} \quad \text{for all } n \geq 2$$

with the initial conditions $F_{-(k-2)}^{(k)} = F_{-(k-2)}^{(k)} = \cdots = F_0^{(k)} = 0$ and $F_1^{(k)} = 1$. When $k = 2$, this sequence coincides with the sequence of Fibonacci numbers. They proved that Eq. (3) has no positive integer solution (k, n, m, x) with $k \geq 3$, $n \geq 2$, and $x \geq 1$. Another related result involving the balancing numbers was studied by Rihane et al. in [10].

In this paper, we study Eq. (3) in nonnegative integers (r, n, m, x) treating r as an integer parameter. We may assume that $r \geq 3$ since the cases $r \in \{1, 2\}$ have been treated already in [8, 11], respectively. The solution with $(n, m) = (0, 1)$ (for any r and x) is obvious so we omit it and suppose that n is positive. Our main result is the following.

Theorem 1 *There is no positive integer solution (r, n, m, x) of Diophantine equation (3) with $r \geq 3$ and $x \neq 2$.*

2 Preliminary results

2.1 The Lucas sequence

Let

$$(\alpha, \beta) := \left(\frac{r + \sqrt{r^2 + 4}}{2}, \frac{r - \sqrt{r^2 + 4}}{2} \right)$$

be the roots of the characteristic equation $x^2 - rx - 1 = 0$ of the Lucas sequence $\mathbf{U} = (U_n)_{n \geq 0}$. We put $\Delta = r^2 + 4 = (\alpha - \beta)^2$ for the discriminant of the above quadratic equation. The Binet formula for the general term of $\mathbf{U}$ is given by

$$U_n := \frac{\alpha^n - \beta^n}{\alpha - \beta} \quad \text{for all } n \geq 0. \quad (4)$$

One may prove by induction that the inequality

$$\alpha^{n-2} \leq U_n \leq \alpha^{n-1} \quad (5)$$

holds for all positive integers n . It is also easy to show that the inequality

$$\frac{U_n}{U_{n+1}} < \frac{1}{r} \quad (6)$$

holds for all $n \geq 2$. Indeed, it follows from $U_{n+1} = rU_n + U_{n-1} > rU_n$ for $n \geq 2$. At one point of the argument we will need the companion Lucas sequence $\mathbf{V} := \{V_n\}_{n \geq 0}$ given by $V_0 = 2$, $V_1 = r$, and $V_{n+2} = rV_{n+1} + V_n$ for all $n \geq 0$. Its Binet formula is

$$V_n = \alpha^n + \beta^n \quad \text{for all } n \geq 0. \quad (7)$$

There are many relations between members of $\mathbf{U}$ and $\mathbf{V}$ such as

$$U_{2n} = U_n V_n; \quad (8)$$

$$U_{n+1}^2 - U_n U_{n+2} = (-1)^n, \quad (9)$$

$$V_n^2 - \Delta U_n^2 = 4(-1)^n, \quad (10)$$

$$2U_{m+n} = U_m V_n + U_n V_m, \quad (11)$$

$$\gcd(U_n, U_m) = U_{\gcd(m,n)}. \quad (12)$$

We record another one. All such identities follow easily from the Binet formulas (4) and (7) of $\mathbf{U}$ and $\mathbf{V}$, respectively.

Lemma 1 *If n is odd then*

$$U_n - 1 = \begin{cases} U_{(n-1)/2} V_{(n+1)/2} & \text{if } n \equiv 1 \pmod{4}, \\ U_{(n+1)/2} V_{(n-1)/2} & \text{if } n \equiv 3 \pmod{4}. \end{cases} \quad (13)$$

Both sequences $\mathbf{U}$ and $\mathbf{V}$ can be extended to negative indices either by allowing n to be negative in the Binet formulas (4) and (7) or simply by using the recurrence relation to extend $\mathbf{U}$ to negative indices via $U_{-n} = -rU_{-(n-1)} + U_{-(n-2)}$ for all $n \geq 1$. The same applies to $\mathbf{V}$. All the above formulas given in Eqs. (8), (9), (10), (11), and (13) hold when the indices are arbitrary integers, not necessarily nonnegative.

The following lemma is useful. For further details we refer the reader to the book of Koshy [6].

Lemma 2 Let $\{U_n(r)\}_{n \geq 0} \subseteq \mathbf{Z}[r]$ be the sequence of polynomials defined by $U_0(r) = 0$, $U_1(r) = 1$, and

$$U_{n+2}(r) = rU_{n+1}(r) + U_n(r) \text{ for all } n \geq 0.$$

Then,

$$U_n(r) = \sum_{\substack{0 \leq k \leq n \\ k \not\equiv n \pmod{2}}} \binom{\frac{n+k-1}{2}}{k} r^k. \quad (14)$$

Note that the summation range above is only over these $k \in [0, n]$ which have different parity than n . We record the following easy but useful consequence of the formula (14).

Lemma 3 We have $r \mid U_n$ if n is even and $r \mid U_n - 1$ if n is odd.

2.2 Logarithmic height

Let η be an algebraic number of degree d with minimal primitive polynomial over the integers

$$a_0x^d + a_1x^{d-1} + \cdots + a_d = a_0 \prod_{i=1}^d (x - \eta^{(i)}),$$

where the leading coefficient a_0 is positive and the $\eta^{(i)}$'s are the conjugates of η . The *logarithmic height* of η is given by

$$h(\eta) := \frac{1}{d} \left(\log a_0 + \sum_{i=1}^d \log \left(\max \{ |\eta^{(i)}|, 1 \} \right) \right).$$

In particular, if $\eta = p/q$ is a rational number with $\gcd(p, q) = 1$ and $q > 0$, then $h(\eta) = \log \max\{|p|, q\}$. The following are some of the properties of the logarithmic height function $h(\cdot)$, which will be used in the next sections of this paper without reference:

$$\begin{aligned} h(\eta \pm \gamma) &\leq h(\eta) + h(\gamma) + \log 2, \\ h(\eta\gamma^{\pm 1}) &\leq h(\eta) + h(\gamma), \\ h(\eta^s) &= |s|h(\eta) \quad (s \in \mathbb{Z}). \end{aligned} \quad (15)$$

2.3 Linear forms in logarithms and continued fractions

The following result on linear forms in three logarithms is due to Mignotte [9]. The result is more general (i.e. the conditions on the parameters involved are somewhat more general), but we will only quote it in the form that we need.

Theorem 2 Consider three algebraic numbers γ_1, γ_2 , and γ_3 , which are all real, greater than 1 and multiplicatively independent. Put

$$\mathcal{D} := [\mathbb{Q}(\gamma_1, \gamma_2, \gamma_3) : \mathbb{Q}].$$

Let b_1, b_2, b_3 be coprime positive integers and consider

$$\Gamma := b_2 \log \gamma_2 - b_1 \log \gamma_1 - b_3 \log \gamma_3.$$

Put

$$d_1 := \gcd(b_1, b_2) = \frac{b_1}{b'_1} = \frac{b_2}{b'_2}, \quad d_3 := \gcd(b_3, b_2) = \frac{b_2}{b''_2} = \frac{b_3}{b''_3}.$$

Let A_1, A_2 , and A_3 be real numbers such that

$$A_i \geq \max\{4, 4.296 \log \gamma_i + 2\mathcal{D}h(\gamma_i)\}, \quad i = 1, 2, 3, \quad \text{and} \quad \Omega := A_1 A_2 A_3 \geq 100.$$

Put

$$b' := \left(\frac{b'_1}{A_2} + \frac{b'_2}{A_1} \right) \left(\frac{b''_3}{A_2} + \frac{b''_2}{A_3} \right) \quad \text{and} \quad \log \mathcal{B} := \max \left\{ 0.882 + \log b', \frac{10}{\mathcal{D}} \right\}.$$

Then, either

$$\log |\Gamma| > -790.95 \Omega \mathcal{D}^2 (\log \mathcal{B})^2,$$

or one of the following conditions holds:

(i) there exist two positive integers r_0 and s_0 such that

$$r_0 b_2 = s_0 b_1$$

with

$$r_0 \leq 5.61 A_2 (\mathcal{D} \log \mathcal{D})^{\frac{1}{3}} \quad \text{and} \quad s_0 \leq 5.61 A_1 (\mathcal{D} \log \mathcal{D})^{\frac{1}{3}};$$

(ii) there exist integers r_1, s_1, t_1 , and t_2 , with $r_1 s_1 \neq 0$, such that

$$(t_1 b_1 + r_1 b_3) s_1 = r_1 b_2 t_2, \quad \gcd(r_1, t_1) = \gcd(s_1, t_2) = 1$$

which also satisfy

$$\begin{aligned} |r_1 s_1| &\leq 5.61 \delta A_3 (\mathcal{D} \log \mathcal{D})^{\frac{1}{3}}, \quad |s_1 t_1| \leq 5.61 \delta A_1 (\mathcal{D} \log \mathcal{D})^{\frac{1}{3}}, \\ |r_1 t_2| &\leq 5.61 \delta A_2 (\mathcal{D} \log \mathcal{D})^{\frac{1}{3}}, \end{aligned}$$

where

$$\delta := \gcd(r_1, s_1).$$

Moreover, when $t_1 = 0$ we can take $r_1 = 1$, and when $t_2 = 0$ we can take $s_1 = 1$.

At some point we will need to treat linear forms in two logarithms of algebraic numbers. To set the stage, let γ_1 and γ_2 be real algebraic numbers which are positive and let $\mathcal{D} := [\mathbb{Q}(\gamma_1, \gamma_2) : \mathbb{Q}]$. Let b_1, b_2 be nonzero integers, let B_1 and B_2 be real numbers larger than 1 such that

$$\log B_i \geq \max \left\{ h(\gamma_i), \frac{|\log \gamma_i|}{\mathcal{D}}, \frac{1}{\mathcal{D}} \right\}, \quad \text{for } i = 1, 2,$$

and put

$$b' = \frac{|b_1|}{\mathcal{D} \log B_2} + \frac{|b_2|}{\mathcal{D} \log B_1}.$$

Let

$$\Gamma := b_1 \log \gamma_1 + b_2 \log \gamma_2. \quad (16)$$

The following result of Laurent et al. is Corollary 2 in [7].

Theorem 3 *With the above notations assuming furthermore that γ_1 and γ_2 are multiplicatively independent we have*

$$\log |\Gamma| > -24.34\mathcal{D}^4 \left(\max \left\{ \log b' + 0.14, \frac{21}{\mathcal{D}}, \frac{1}{2} \right\} \right)^2 \log B_1 \log B_2. \quad (17)$$

Note that the fact that $\Gamma \neq 0$ is already guaranteed by the condition that γ_1 and γ_2 are multiplicatively independent together with the fact that b_1, b_2 are nonzero integers.

During the calculations we get upper bounds on our variables which are too large, thus we need to reduce them. To do so we use some results from the theory of continued fractions.

For the treatment of linear forms homogeneous in two integer variables we use a well-known classical result in the theory of Diophantine approximation due to Legendre.

Lemma 4 *Let τ be an irrational number, $p_0/q_0, p_1/q_1, p_2/q_2, \dots$ be the sequence of convergents of the continued fraction expansion of τ and M be a positive integer. Let N be a nonnegative integer such that $q_N > M$. Putting $a(M) := \max\{a_i : i = 0, 1, 2, \dots, N\}$ the inequality*

$$\left| \tau - \frac{u}{v} \right| > \frac{1}{(a(M) + 2)v^2}$$

holds for all pairs (u, v) of positive integers with $0 < v < M$. Furthermore, if

$$\left| \tau - \frac{u}{v} \right| < \frac{1}{2v^2},$$

then $u/v = p_k/q_k$ for some $k \geq 0$.

For a nonhomogeneous linear form in two integer variables we use a slight variation of a result due to Dujella and Pethő (see [4, Lemma 5a]). For a real number X , we write $\|X\| := \min\{|X - n| : n \in \mathbb{Z}\}$ for the distance from X to the nearest integer.

Lemma 5 *Let M be a positive integer, p/q be a convergent of the continued fraction expansion of the irrational number τ such that $q > 6M$, and A, B, μ be some real numbers with $A > 0$ and $B > 1$. Furthermore, let $\varepsilon := \|\mu q\| - M\|\tau q\|$. If $\varepsilon > 0$, then there is no solution to the inequality*

$$0 < |u\tau - v + \mu| < AB^{-w}$$

in positive integers u, v , and w with

$$u \leq M \quad \text{and} \quad w \geq \frac{\log(Aq/\varepsilon)}{\log B}.$$

3 Proof of Theorem 1

3.1 The cases $n = 1$ or $x = 1$

We assume that $n \geq 1$, as the solution with $n = 0$ is trivial. Since $U_{n+1} < U_{n+1} + U_n < U_{n+2}$, it follows that the Diophantine equation (3) has no solution with $x = 1$. Let us assume that $n = 1$. We then get that

$$U_m = 1 + r^x. \quad (18)$$

In particular, $U_m \equiv 1 \pmod{r}$. Lemma 3 shows that $m \equiv 1 \pmod{2}$ and now Lemma 1 shows that

$$r^x = U_m - 1 = U_{(m-\delta)/2} V_{(m+\delta)/2} \quad \text{where } \delta \in \{\pm 1\}, \quad \delta \equiv m \pmod{4}.$$

We now recall the Primitive Divisor Theorem of Carmichael [3] (see [1] for the most general statement) for the sequence $\mathbf{U}$. It states that if $\ell > 12$, there is a prime factor p of U_ℓ which is primitive in the sense that $p \nmid U_k$ for any positive $k < \ell$. So, assume $m + \delta \geq 14$. Since $m + \delta$ is even we have $U_{m+\delta} = U_{(m+\delta)/2} V_{(m+\delta)/2}$ by the formula (8). Since $U_{m+\delta}$ has a primitive prime factor p , the primitive prime p must be a divisor of $V_{(m+\delta)/2}$, which in turn must divide $r = U_2$, a contradiction. Thus, $m + \delta \leq 12$, therefore $m \leq 13$. It thus follows that

$$r^x = U_m - 1 < \alpha^{m-1} \leq \alpha^{12} < (r^2 + 4)^6,$$

so, if $x \geq 13$, then

$$3^{x-12} \leq r^{x-12} < (1 + 4/r^2)^6 \leq (1 + 4/3^2)^6 < 9.1,$$

which gives $x \leq 14$. Thus, $m \leq 13$, $x \leq 14$. For each choice of the pair (m, x) with the components in the above ranges, the equation $U_m(r) - 1 - r^x = 0$ is a polynomial equation in r . After a simple computer search, we found no other solutions to Eq. (18) apart from the solution $(n, m, x) = (1, 3, 2)$ which has $x = 2$ so it is part of the parametric family of solutions indicated at Eq. (2).

So, from now on we assume that $n \geq 2$ and $x \geq 3$.

3.2 Calculations when $n \in [2, 100]$ and $x \in [3, 100]$

Using the Eq. (3) and the inequality (5), we get

$$\alpha^{(n-1)x} < U_{n+1}^x < U_n^x + U_{n+1}^x = U_m \leq \alpha^{m-1},$$

and

$$\alpha^{m-2} < U_m = U_n^x + U_{n+1}^x < (U_n + U_{n+1})^x < U_{n+2}^x < \alpha^{(n+1)x}.$$

From the above inequalities, we get the following result which we record for future reference.

Lemma 6 *The inequalities*

$$(n-1)x + 1 < m < (n+1)x + 2 \quad (19)$$

hold for all $x \geq 3$ and $n \geq 2$.

We next consider Eq. (14) given in Lemma 2. We write Eq. (3) as

$$\left(\sum_{\substack{0 \leq k \leq n \\ k \not\equiv n \pmod{2}}} \binom{\frac{n+k-1}{2}}{k} r^k \right)^x + \left(\sum_{\substack{0 \leq k \leq n+1 \\ k \not\equiv n+1 \pmod{2}}} \binom{\frac{n+k}{2}}{k} r^k \right)^x = \sum_{\substack{0 \leq k \leq m \\ k \not\equiv m \pmod{2}}} \binom{\frac{m+k-1}{2}}{k} r^k. \quad (20)$$

Assume first that n is even. Then Eq. (20) becomes

$$\begin{aligned} & \left(\frac{n}{2} r + \binom{\frac{n+2}{2}}{3} r^3 + \dots \right)^x + \left(1 + \binom{\frac{n+2}{2}}{2} r^2 + \binom{\frac{n+4}{2}}{4} r^4 + \dots \right)^x \\ &= 1 + \binom{\frac{m+1}{2}}{2} r^2 + \binom{\frac{m+3}{2}}{3} r^4 + \dots, \end{aligned}$$

which is equivalent to

$$\binom{\frac{n+2}{2}}{2} r^2 x + \binom{\frac{n+4}{2}}{4} r^4 x + \dots \equiv \binom{\frac{m+1}{2}}{2} r^2 + \binom{\frac{m+3}{2}}{3} r^4 + \dots \pmod{r^x}. \quad (21)$$

The above relation implies that

$$r^{\min\{x,4\}-2} \left| \binom{\frac{m+1}{2}}{2} - x \binom{\frac{n+2}{2}}{2} \right|. \quad (22)$$

Similarly, when n is odd, one is led to the analogous divisibility relation

$$r^{\min\{x,4\}-2} \left| \binom{\frac{m+1}{2}}{2} - x \binom{\frac{n+1}{2}}{2} \right|. \quad (23)$$

So, fixing $n \in [2, 100]$ and $x \in [3, 100]$, inequalities (19) give some range for m . For each (n, x, m) , divisibility relations (22) and (23) (according to whether n is even or odd) give us some possibilities for $r \geq 3$ and now one checks whether relation (3) holds for this candidate (n, x, m, r) . A computer search with Mathematica in this range for n and x which ran for a few hours found no solutions. For the search we didn't actually checked that formula (20) holds but we checked that Eq. (20) does not hold modulo T , where T is the product of the first 20 primes. The Mathematica function `powermod` allowed us to compute the powers of r modulo T arising from the binomial formula rather quickly.

From now on, we assume that $n \geq 2$, $x \geq 3$ and $\max\{n, x\} > 100$.

3.3 A small linear form in three logs

We rewrite Eq. (3) as

$$\frac{\alpha^m}{\alpha - \beta} - U_{n+1}^x = U_n^x + \frac{\beta^m}{\alpha - \beta}. \quad (24)$$

Dividing both sides of the above equation by U_{n+1}^x and using the inequality (6), we obtain

$$\left| \alpha^m (\alpha - \beta)^{-1} U_{n+1}^{-x} - 1 \right| = \left(\frac{U_n}{U_{n+1}} \right)^x + \frac{\beta^m}{(\alpha - \beta) U_{n+1}^x} < 2 \left(\frac{U_n}{U_{n+1}} \right)^x < \frac{2}{r^x}. \quad (25)$$

Put

$$\Lambda := \alpha^m (\alpha - \beta)^{-1} U_{n+1}^{-x} - 1 \quad \text{and} \quad \Gamma := m \log \alpha - \log(\alpha - \beta) - x \log U_{n+1}. \quad (26)$$

We observe that $\Lambda = e^\Gamma - 1$, where Λ and Γ are given by (26). Since $|\Lambda| \leq 2/27$, we have that $e^{|\Gamma|} \leq 27/25$ and using the inequality (25) we obtain

$$|\Gamma| = \left| m \log \alpha - \log(\sqrt{r^2 + 4}) - x \log U_{n+1} \right| \leq e^{|\Gamma|} |e^{|\Gamma|} - 1| \leq \frac{27|\Lambda|}{25} < \frac{2.2}{r^x}. \quad (27)$$

We record the above inequality for future reference.

Lemma 7 *With Γ given by formula (26), inequality (27) holds.*

We want to apply Theorem 2 with the following data:

$$\gamma_1 := \alpha - \beta = \sqrt{r^2 + 4}, \quad \gamma_2 := \alpha, \quad \gamma_3 := U_{n+1}, \quad b_1 := 1, \quad b_2 := m, \quad b_3 := x.$$

We need to check that γ_1 , γ_2 , and γ_3 are multiplicatively independent. This we do in the next subsection.

3.4 Checking that γ_1 , γ_2 , γ_3 are multiplicatively independent

Well, assume they are not and let i, j, k be integers not all zero such that

$$\gamma_1^i \gamma_2^j \gamma_3^k = 1.$$

Squaring and rearranging the above relation we get $\gamma_2^{2j} = (\gamma_1^2)^{-i} \gamma_3^{-2k} \in \mathbb{Q}$. However, γ_2^{2j} is also a unit, so an algebraic integer whose reciprocal is also an algebraic integer, and it is also positive, so it must be 1. Thus, $j = 0$. It now follows that i and k are both nonzero (since if one of them is, so is the other one) and further $\gamma_3 = \gamma_1^{-i/k}$. In particular, all prime factors of U_{n+1} are prime factors of $\Delta := r^2 + 4$. But this is also contemplated by the Primitive Divisor Theorem of Carmichael since primes dividing Δ are not considered primitive. In particular, U_{n+1} does not have primitive prime factors so $n + 1 \leq 12$. In fact, Theorem C in [1] together with Table 1 there show that either $n + 1 \in \{2, 3, 4, 6\}$ or $n + 1 \in \{5, 12\}$ but in this last case, the only such Lucas sequence $\mathbf{U}$ for which either one of U_5 or U_{12} does not have primitive prime factors is the sequence of Fibonacci numbers, which is not our case. Thus, $n + 1 \in \{2, 3, 6\}$. Further, for each prime p let $z(p)$ be the index of appearance of p in $\mathbf{U}$ defined as the smallest positive integer k such that $p \mid U_k$. This always exists for our sequence $\mathbf{U}$ since α is a quadratic unit. It has the additional property that if ℓ is a positive integer then $p \mid U_\ell$ if and only if $z(p) \mid \ell$. It is also well-known and easy to prove that if $p \mid \Delta$, then $z(p) = p$. Since also $z(p) \mid n + 1$ and $n + 1 \in \{2, 3, 6\}$, it follows that the only possibilities for p are $p = 2, 3$. Hence, $r^2 + 4 = 2^a 3^b$. However, 3 cannot divide $r^2 + 4$ for any positive integer r (because -4 is not a quasartic residue modulo 3), so $b = 0$ and $2^a = r^2 + 4$. Thus, $r = 2r_0$ is even, $a \geq 3$ and the equation simplifies to $2^{a-2} = r_0^2 + 1$. Hence, r_0 is odd, so $r_0^2 \equiv 1 \pmod{8}$, therefore $2 \parallel r_0^2 + 1$, which leads to $a = 3$, $r_0 = 1$, which gives $r = 2$, is not our case. Hence, indeed γ_1 , γ_2 , γ_3 are multiplicatively independent.

3.5 Applying Theorem 2

Since $\gamma_1, \gamma_2, \gamma_3 \in \mathbb{Q}(\alpha)$, we have $\mathcal{D} = 2$. We also have

$$r < \alpha < \sqrt{r^2 + 4} < r + 1.$$

So, we bound the heights of α and $\sqrt{r^2 + 4}$ in terms of $\log(r + 1)$. Since

$$h(\gamma_1) = h(\alpha - \beta) = \frac{1}{2} \log(r^2 + 4) < \log(r + 1), \quad h(\gamma_2) = \frac{1}{2} \log \alpha < \frac{1}{2} \log(r + 1),$$

and

$$h(\gamma_3) = \log U_{n+1} < \log \alpha^n = n \log \alpha < n \log(r + 1),$$

we can take

$$A_1 := 8.296 \log(r + 1), \quad A_2 := 6.296 \log(r + 1), \quad A_3 := 8.296n \log(r + 1).$$

Thus $\Omega := A_1 A_2 A_3 > 433n(\log(r + 1))^3 > 100$ since $n \geq 2$ and $r \geq 3$. Then,

$$\begin{aligned} b' &= \left(\frac{1}{6.296 \log(r + 1)} + \frac{m}{8.296 \log(r + 1)} \right) \left(\frac{x}{6.296 \log(r + 1)} + \frac{m}{8.296n \log(r + 1)} \right) \\ &< \frac{m}{(\log(r + 1))^2} \left(\frac{1}{6.296} + \frac{1}{8.296} \right) \frac{x}{\log(r + 1)} \left(\frac{1}{6.296} + \frac{2}{8.296} \right) \\ &< \frac{0.12mx}{(\log(r + 1))^2}. \end{aligned} \quad (28)$$

In the above chain of inequalities, we used the fact that $m < (n + 1)x + 2 < 2nx$ [see inequality (19)] since $n \geq 2$ and $x \geq 3$. Further, putting

$$\log \mathcal{B} := \max \left\{ 0.882 + \log \left(\frac{0.12mx}{(\log(r + 1))^2} \right), 5 \right\},$$

we have that either the inequality

$$\log |\Gamma| > -790.95 \times 434n(\log(r + 1))^3 \times 2^2 \times (\log \mathcal{B})^2 \quad (29)$$

holds, or the other possibilities (i), (ii) from Theorem 2 hold. We treat (i) and (ii) later and deal with the above inequality (29) at this stage. If $\log \mathcal{B} = 5$, then

$$x < mx < \frac{e^{5-0.882}}{0.12} (\log(r + 1))^2 < 512(\log(r + 1))^2. \quad (30)$$

On the other hand, if $\log \mathcal{B} > 5$, then

$$\log \mathcal{B} := 0.882 + \log \left(\frac{0.12mx}{(\log(r+1))^2} \right) < \log \left(\frac{0.3mx}{(\log(r+1))^2} \right), \quad (31)$$

where in the above inequality we used the fact that $e^{0.882} \times 0.12 < 0.3$. Thus, we get that

$$\begin{aligned} \log |\Gamma| &> -790.95 \times 434n(\log(r+1))^3 \times 2^2 \times \left(\log \left(\frac{0.3mx}{(\log(r+1))^2} \right) \right)^2 \\ &> -1.374 \times 10^6 n(\log(r+1))^3 \left(\log \left(\frac{0.3mx}{(\log(r+1))^2} \right) \right)^2. \end{aligned}$$

Comparing this inequality with Eq. (27), we get that

$$x \log r - \log 2.2 < 1.374 \times 10^6 n(\log(r+1))^3 \left(\log \left(\frac{0.3mx}{(\log(r+1))^2} \right) \right)^2,$$

which implies, via the inequality

$$\frac{\log(r+1)}{\log r} = 1 + \frac{\log(1+1/r)}{\log r} < 1 + \frac{1}{r \log r}, \quad (32)$$

that

$$\begin{aligned} x &< \frac{\log 2.2}{\log r} + 1.374 \times 10^6 n \left(\frac{\log(r+1)}{\log r} \right) (\log(r+1))^2 \left(\log \left(\frac{0.3mx}{(\log(r+1))^2} \right) \right)^2 \\ &< \frac{\log 2.2}{\log r} + 1.374 \times 10^6 n \left(1 + \frac{1}{r \log r} \right) (\log(r+1))^2 \left(\log \left(\frac{0.3mx}{3(\log(r+1))^2} \right) \right)^2 \\ &< 1.38 \times 10^6 n \left(1 + \frac{1}{r \log r} \right) (\log(r+1))^2 \left(\log \left(\frac{0.3mx}{(\log(r+1))^2} \right) \right)^2. \end{aligned} \quad (33)$$

Using the inequality (19), we know that $m < (n+1)x + 2 < (n+1)(x+1)$ (because $n \geq 2$), and substituting this in (33) we get that

$$x < 1.38 \times 10^6 n \left(1 + \frac{1}{r \log r} \right) (\log(r+1))^2 \left(\log \left(\frac{0.3(n+1)(x+1)^2}{(\log r)^2} \right) \right)^2. \quad (34)$$

We now turn our attention to the possibilities (i) and (ii). In case (i), there are positive integers r_0, s_0 , which may be assumed to be coprime, such that $r_0 b_2 = s_0 b_1$. So, we get $r_0 m = s_0$ and since r_0, s_0 are coprime, we take $r_0 = 1, s_0 = m$, and we get

$$m = s_0 < 5.61 A_1 (\mathcal{D} \log \mathcal{D})^{1/3} < 5.61 \times 8.296 \times (2 \log 2)^{1/3} \log(r+1)$$

$$\begin{aligned} &< 5.61 \times 8.296 \times (2 \log 2)^{1/3} \log(r+1) \\ &< 52 \log(r+1). \end{aligned}$$

Since $m > (n-1)x + 1 > x$, this situation gives

$$x < 52 \log(r+1). \quad (35)$$

This was in situation (i). In situation (ii), we have integers r_1, s_1, t_1, t_2 with $r_1 s_1 \neq 0$ and

$$(t_1 b_1 + r_1 b_3) s_1 = r_1 b_2 t_2, \quad \gcd(r_1, t_1) = \gcd(s_1, t_2) = 1.$$

Thus, for us, we have

$$(t_1 + r_1 x) s_1 = r_1 m t_2, \quad \gcd(r_1, t_1) = \gcd(s_1, t_2) = 1.$$

Reducing the above equation modulo r_1 we get $t_1 s_1 \equiv 0 \pmod{r_1}$ and since $\gcd(t_1, r_1) = 1$, we get that $r_1 \mid s_1$. So, we put $s_1 = r_1 s'_1$ and simplify both sides of the above equation by r_1 to get

$$(t_1 + r_1 x) s'_1 = m t_2.$$

Consequently, for us $\delta = \gcd(r_1, s_1) = r_1$. Hence,

$$\begin{aligned} |t_1 s'_1| &< 5.61 A_1 (\mathcal{D} \log \mathcal{D})^{1/3} < 5.61 \times 8.296 \log(r+1) \times (2 \log 2)^{1/3} < 52 \log(r+1), \\ |t_2| &< 5.61 A_2 (\mathcal{D} \log \mathcal{D})^{1/3} < 5.61 \times 6.296 (2 \log 2)^{1/3} \log(r+1) < 40 \log(r+1), \\ |r_1 s'_1| &< 5.61 A_3 (\mathcal{D} \log \mathcal{D})^{1/3} < 5.61 \times 8.296 (2 \log 2)^{1/3} n \log(r+1) < 52n \log(r+1). \end{aligned} \quad (36)$$

Assume first that $t_2 = 0$. Then

$$x = |t_1|/|r_1| \leq |t_1| < 52 \log(r+1),$$

which is the same as (35). Assume next that $t_2 \neq 0$. We return to inequality (27) and multiply both sides by t_2 and get

$$|m t_2 \log \gamma_2 - t_2 \log \gamma_1 - x t_2 \log \gamma_3| < \frac{2.2 |t_2|}{r^x}.$$

We substitute $m t_2$ by $t_1 s'_1 + (r_1 s'_1) x$ inside the left-hand side above and then the left-hand side above becomes

$$\left| \log \left(\frac{\gamma_2^{t_1 s'_1}}{\gamma_1^{t_2}} \right) + x \log \left(\frac{\gamma_2^{r_1 s'_1}}{\gamma_3^{t_2}} \right) \right| < \frac{2.2 |t_2|}{r^x}. \quad (37)$$

Inequality (37) is of the form

$$|\Gamma_1| < \frac{2.2|t_2|}{r^x}, \quad \text{where } \Gamma_1 := \log \eta_1 + x \log \eta_2, \quad (38)$$

and

$$\eta_1 := \frac{\gamma_2^{t_1 s'_1}}{\gamma_1^{t_2}}, \quad \eta_2 := \frac{\gamma_2^{r_1 s'_1}}{\gamma_3^{t_2}}.$$

We check that η_1 and η_2 are multiplicatively independent. If not, there are integers i, j not both zero such that $\eta_1^i \eta_2^j = 1$. This gives

$$\left(\frac{\gamma_2^{t_1 s'_1}}{\gamma_1^{t_2}} \right)^i \left(\frac{\gamma_2^{r_1 s'_1}}{\gamma_3^{t_2}} \right)^j = 1.$$

If $i \neq 0$, this gives a multiplicative dependence among $\gamma_1, \gamma_2, \gamma_3$ with the exponent of γ_1 being the nonzero integer $-t_2 i$, a contradiction with the main result of Sect. 3.4. Thus, $i = 0$, so $j \neq 0$, and we get again a multiplicative relation among γ_2, γ_3 (the exponent of γ_3 being the nonzero integer $-t_2 j$), which is the same contradiction. Thus, indeed η_1 and η_2 are multiplicatively independent and they are also positive. So, we are in position to apply Theorem 3 to the left-hand side of inequality (38). We compute $\log B_i$ for these choices. We have, by the properties (15),

$$\begin{aligned} h(\eta_1) &\leq |t_1 s'_1| h(\gamma_2) + |t_2| h(\gamma_1) \leq (52 \log(r+1))(1/2 \log(r+1)) \\ &\quad + (40 \log(r+1)) \log(r+1) \\ &= 66(\log(r+1))^2, \\ h(\eta_2) &\leq |r_1 s'_1| h(\gamma_2) + |t_2| h(\gamma_3) \leq 52n \log(r+1)(1/2 \log(r+1)) \\ &\quad + (40 \log(r+1))(n \log(r+1)) \\ &= 66n(\log(r+1))^2. \end{aligned}$$

Since $|\log \gamma_i|/2 \leq h(\gamma_i)$ holds for $i = 1, 2, 3$, it follows, by the absolute value inequality, that the same inequalities are satisfied by the numbers $|\log \eta_i|/2$ for $i = 1, 2$. Thus, since $\mathcal{D} = 2$, we can take

$$\log B_1 := 66(\log(r+1))^2, \quad \log B_2 := 66n(\log(r+1))^2.$$

We bound

$$\frac{1}{2 \log B_2} + \frac{x}{2 \log B_1} = \frac{1}{132(\log(r+1))^2} \left(\frac{1}{n} + x \right) < \frac{x+1}{132(\log(r+1))^2}.$$

Hence, we take

$$b' := \frac{x+1}{132(\log(r+1))^2}.$$

Now Theorem 3 gives

$$\begin{aligned} \log |\Lambda| &> -24.34 \times 2^4 (\max\{\log b' + 0.14, 10.5\})^2 \times (66(\log(r+1))^2) \\ &\quad \times (66n(\log(r+1))^2) \\ &> -1.627 \times 10^6 n (\log(r+1))^4 M^2, \end{aligned} \quad (39)$$

where $M := \max\{\log b' + 0.4, 10.5\}$. In case $M = 10.5$, we get

$$b' < e^{10.5-0.4} < 24400,$$

which gives

$$x + 1 < 24400 \times 132 (\log(r+1))^2 < 3.3 \times 10^6 (\log(r+1))^2. \quad (40)$$

Finally, suppose that

$$M = \log b' + 0.4 = \log(e^{0.4}b') < \log(1.5b') = \log\left(\frac{x+1}{88(\log(r+1))^2}\right).$$

Comparing inequality (39) with inequality (38), we get

$$x \log r - \log(2.2|t_2|) < 1.627 \times 10^6 n (\log(r+1))^4 \left(\log\left(\frac{x+1}{88(\log(r+1))^2}\right) \right)^2.$$

Since $2.2|t_2| < 88 \log(r+1)$, we get that

$$\begin{aligned} x &< \frac{\log(88 \log(r+1))}{\log r} \\ &\quad + 1.627 \times 10^6 \left(\frac{\log(r+1)}{\log r} \right) (\log(r+1))^3 \left(\log\left(\frac{x+1}{88(\log(r+1))^2}\right) \right)^2. \end{aligned}$$

The first summand in the right-hand side is < 5 for all $r \geq 3$. Using inequality (32), we get that

$$x < 1.63 \times 10^6 n \left(1 + \frac{1}{r \log r} \right) (\log(r+1))^3 \left(\log\left(\frac{x+1}{88(\log(r+1))^2}\right) \right)^2. \quad (41)$$

To summarise, either we are in the first situation of Theorem 2 and $\log \mathcal{B} = 5$, in which case inequality (30) holds, or $\log \mathcal{B} > 5$ in which case inequality (34) holds, or we are in the exceptional case (i) for which inequality (35) holds, which is contained in inequality (30), or we are in the exceptional situation (ii) in which case either inequality (40) holds, or inequality (63) holds. Since inequality (30) is contained in inequality (40), it follows, using the inequality $1/88 < 0.12$, that we proved the following result.

Lemma 8 *One of the following inequalities holds:*

$$x < 3.3 \times 10^6 (\log(r+1))^2, \quad (42)$$

$$x < 1.38 \times 10^6 n \left(1 + \frac{1}{r \log r}\right) (\log(r+1))^2 \left(\log \left(\frac{0.3(n+1)(x+1)^2}{(\log(r+1))^2}\right)\right)^2, \quad (43)$$

$$x < 1.63 \times 10^6 n \left(1 + \frac{1}{r \log r}\right) (\log(r+1))^3 \left(\log \left(\frac{0.12(x+1)}{(\log(r+1))^2}\right)\right)^2. \quad (44)$$

3.6 More inequalities in terms of n and x

We put

$$\kappa := nx + 1 - m \quad (45)$$

Later we shall show that κ is positive except possibly if $r = 3$. In this section, assuming that it is positive, we show how it gives some lower bounds for x in terms of n .

Lemma 9 *The following holds :*

- (i) $\kappa \neq 1$,
- (ii) *If $\kappa = 2$ and $n \geq 3$ then $x \geq r^{\max\{2, n-3\}}$,*
- (iii) *If $\kappa \geq 3$, then $\kappa \geq n/2$.*

Proof (i). If $\kappa = 1$, then $m = nx$. So Eq. (3) becomes

$$U_n^x + U_{n+1}^x = U_{nx}.$$

If p is prime dividing U_n (which exists since $n > 1$), then $p \mid U_n^x$ and $p \mid U_n \mid U_{nx}$, so from the above equation we get $p \mid U_{n+1}$, a contradiction since $\gcd(U_n, U_{n+1}) = U_{\gcd(n, n+1)} = 1$ by relation (12).

(ii) In this case $m = nx - 1$ so Eq. (3) becomes

$$U_n^x + U_{n+1}^x = U_{nx-1}.$$

In particular, $U_{nx-1} - U_{n+1}^x \equiv 0 \pmod{U_n^2}$. We study this congruence. In what follows for three algebraic integers a, b, c , we write $a \equiv b \pmod{c}$ if $(a - b)/c$ is an algebraic integer. Write

$$U_n = \frac{\alpha^n - \beta^n}{\alpha - \beta} \quad \text{as} \quad \alpha^n = \beta^n + \sqrt{\Delta} U_n.$$

Then

$$\alpha^{nx} = (\beta^n + \sqrt{\Delta} U_n)^x \equiv \beta^{nx} + x \beta^{n(x-1)} \sqrt{\Delta} U_n \pmod{\Delta U_n^2}.$$

Thus,

$$U_{nx-1} = \frac{\alpha^{nx} \alpha^{-1} - \beta^{nx-1}}{\sqrt{\Delta}}$$

$$\begin{aligned}
&\equiv \frac{(\beta^{nx} + x\beta^{n(x-1)}\sqrt{\Delta}U_n)\alpha^{-1} - \beta^{nx-1}}{\alpha - \beta} \pmod{\sqrt{\Delta}U_n^2} \\
&\equiv \frac{\beta^{nx}\alpha^{-1} - \beta^{nx}\beta^{-1}}{\alpha - \beta} + x\beta^{n(x-1)}\alpha^{-1}U_n \pmod{U_n^2} \\
&\equiv \beta^{nx} + x\beta^{n(x-1)}\alpha^{-1}U_n \pmod{U_n^2}.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
U_{n+1}^x &= \left(\frac{\alpha^{n+1} - \beta^{n+1}}{\sqrt{\Delta}} \right)^x \\
&\equiv \left(\frac{(\beta^n + \sqrt{\Delta}U_n)\alpha - \beta^{n+1}}{\sqrt{\Delta}} \right)^x \pmod{U_n^2} \\
&\equiv (\beta^n + U_n\alpha)^x \pmod{U_n^2} \\
&\equiv \beta^{nx} + x\beta^{n(x-1)}\alpha U_n \pmod{U_n^2}.
\end{aligned}$$

Thus,

$$\begin{aligned}
U_{nx-1} - U_{n+1}^x &\equiv (\beta^{nx} + x\beta^{n(x-1)}\alpha^{-1}U_n) - (\beta^{nx} + x\beta^{n(x-1)}\alpha U_n) \pmod{U_n^2} \\
&\equiv x\beta^{n(x-1)}(\alpha^{-1} - \alpha)U_n \pmod{U_n^2} \\
&\equiv -x\beta^{n(x-1)}rU_n \pmod{U_n^2}.
\end{aligned}$$

In the last step above, we used the fact that $\alpha^{-1} - \alpha = -\beta - \alpha = -r$. Since the expression $U_{nx-1} - U_{n+1}^x$ is divisible by U_n^2 , we get that $U_n^2 \mid \beta^{n(x-1)}xrU_n$. Since β is a unit, we get that $U_n \mid xr$. For $n = 2$, this gives us nothing since $U_2 = r$. For $n = 3$, $U_3 = r^2 + 1$ is coprime to r , so $U_3 \mid x$, which gives $x \geq r^2 + 1 > r^2$. For $n = 4$, we have that $U_4 = r(r^2 + 2)$ divides rx , so $r^2 + 2 \mid x$ giving $x \geq r^2 + 2 > r^2$. Finally, for $n \geq 5$, we have that $U_n > \alpha^{n-2} > r^{n-2}$ and so $x \geq U_n/r \geq r^{n-3}$. This proves (ii).

(iii) We may assume that $n \geq 7$, otherwise the conclusion is trivial. Recall that $V_n = \alpha^n + \beta^n$. Relations Eqs. (9) and (10) are

$$U_{n+1}^2 - U_n U_{n+2} = (-1)^n \quad \text{and} \quad V_n^2 - \Delta U_n^2 = 4(-1)^n.$$

In particular, $U_{n+1}^2 \equiv (-1)^n \pmod{U_n}$ and $V_n^2 \equiv 4(-1)^n \pmod{U_n}$. We also use the fact that $U_{-m} = (-1)^{m-1}U_m$, $V_{-m} = (-1)^m V_m$ and relation (11) which is

$$2U_{m+n} = U_m V_n + U_n V_m.$$

Armed with these facts, writing $m = nx - (\kappa - 1)$ and

$$U_n^x + U_{n+1}^x = U_{nx-(\kappa-1)},$$

we multiply both sides of the above equation by 2 and write

$$2U_n^x + 2U_{n+1}^x = 2U_{nx-(\kappa-1)} = U_{nx}V_{-(\kappa-1)} + V_{nx}U_{-(\kappa-1)}.$$

We square both sides of the above equation and reduce it modulo U_n taking into account that $U_n \mid U_{nx}$ and $V_{nx}^2 \equiv 4(-1)^{nx} \pmod{U_{nx}} \equiv 4(-1)^{nx} \pmod{U_n}$, and get

$$\begin{aligned} 4(-1)^{nx} &\equiv 4(U_{n+1}^2)^x \pmod{U_n} \equiv (U_{nx} V_{-(\kappa-1)} + V_{nx} U_{-(\kappa-1)})^2 \pmod{U_n} \\ &\equiv V_{nx}^2 U_{-(\kappa-1)}^2 \pmod{U_n} \equiv V_{nx}^2 U_{\kappa-1}^2 \pmod{U_n} \equiv 4(-1)^{nx} U_{\kappa-1}^2 \pmod{U_n}. \end{aligned}$$

Thus, $U_n \mid 4(U_{\kappa-1}^2 - 1)$. The right-hand side is nonzero since $\kappa > 2$. If $\kappa - 1$ is odd, then

$$4(U_{\kappa-1}^2 - 1) = 4U_{\kappa-2}U_{\kappa}$$

by relation (9) (with $n + 1 := \kappa - 1$). Since $n \geq 7$ and $\mathbf{U}$ is not the Fibonacci sequence, it follows that U_n has a primitive divisor p , which must divide one of $U_{\kappa-2}$ or U_{κ} . Thus, $z(p) = n$ divides one of $\kappa - 2$ or κ , so we get $\kappa \geq n$, which is a better conclusion than the one desired. If $\kappa - 1$ is even and $U_n \mid 2(U_{\kappa-1}^2 - 1)$, we then get

$$\alpha^{n-2} < U_n < 2U_{\kappa-1}^2 < r(\alpha^{\kappa-1})^2 < \alpha^{2\kappa-1},$$

so $2\kappa - 2 \geq n - 2$, therefore $\kappa \geq n/2$. The above argument was based on the fact that $r \geq 2$. In particular, if $r \geq 4$, then the same argument gives again that

$$\alpha^{n-2} < U_n < 4U_{\kappa-1}^2 < r(\alpha^{\kappa-1})^2 < \alpha^{2\kappa-1},$$

so $\kappa \geq n/2$. So, the only case when the above arguments fail are when $r = 3$ and $4 \mid U_n$. It then follows that n is even (in fact, n is a multiple of 6, but we shall not need that), so $r \mid U_n \mid 4(U_{\kappa-1}^2 - 1)$. But $U_{\kappa-1}$ is a multiple of r (since $\kappa - 1$ is even), so $U_{\kappa-1}^2 - 1$ is coprime to r . Thus, $r \mid 4$, which is false. This finishes the proof of (iii). $\square$

Corollary 4 *If $\kappa > 0$, then $x > n/2$.*

Proof By Lemma 9, if $\kappa = 2$, then $x \geq r^{\min\{2, n-3\}} \geq 3^{\min\{2, n-3\}} > n/2$ for any $n \geq 2$. If $\kappa \geq 3$, then

$$1 + nx - m = \kappa \geq n/2,$$

which leads to $1 + nx - n/2 \geq m$. Comparing this with the lower bound $m > (n-1)x + 1$ given by inequality (19), we get $x > n/2$. $\square$

3.7 Another inequality among r, n, m, x

In this section, we return to inequality (27) and rewrite it in order to deduce a good approximation of $\log r$ by a rational number whose denominator is a multiple of r^2 . Let's get to work. We need approximations of $\log \alpha$ and $\log \sqrt{r^2 + 4}$ in terms of $\log r$.

Lemma 10 *For $r \geq 3$, we have the following approximations:*

$$\log \sqrt{r^2 + 4} = \log r + \frac{2}{r^2} + \zeta, \quad |\zeta| < \frac{5.81}{r^4}; \quad (46)$$

$$\log \alpha = \log r + \frac{1}{r^2} + \zeta', \quad |\zeta'| < \frac{3.64}{r^4}. \quad (47)$$

Proof We have

$$\log \sqrt{r^2 + 4} = \frac{1}{2} \log(r^2 + 4) = \log r + \frac{1}{2} \log \left(1 + \frac{4}{r^2} \right).$$

With $z := 4/r^2$, we have $|z| \leq 4/9$ and

$$\log(1 + z) = z + \zeta_1, \quad |\zeta_1| \leq \sum_{k \geq 2} \frac{z^k}{k} = \frac{z^2}{2} \left(1 + (2/3)z + (2/4)z^2 + \cdots \right).$$

The expression in parenthesis above is smaller than

$$c_1 := 1 + (2/3)(4/9) + (2/4)(4/9)^2 + \cdots = (-4/9 - \log(1 - 4/9)) \times 2 \times (9/4)^2.$$

Hence,

$$\begin{aligned} \log \sqrt{r^2 + 4} &= \log r + \frac{2}{r^2} + \frac{\zeta_1}{2} := \log r + \frac{2}{r^2} + \zeta, \quad |\zeta| = \frac{|\zeta_1|}{2} \\ &< \left(\frac{c_1}{4} \right) z^2 = \frac{4c_1}{r^4} < \frac{5.81}{r^4}. \end{aligned}$$

For α , we write

$$\begin{aligned} \log \alpha &= \log r + \log \left(\frac{1}{2} + \sqrt{\frac{1}{4} + \frac{1}{r^2}} \right) = \log r + \log(1 + z_1), \\ z_1 &:= \sqrt{\frac{1}{4} + \frac{z}{4}} - \frac{1}{2}. \end{aligned}$$

Note that $|z_1| \leq 1/r^2$. Thus,

$$\log(1 + z_1) = z_1 + \zeta_2, \quad |\zeta_2| \leq \frac{|z_1|^2}{2} (1 + (2/3)|z_1| + \cdots) \leq \frac{c_2}{2r^4},$$

where by the previous arguments,

$$c_2 = (-\lambda_0 - \log(1 - \lambda_0)) \times 2 \times \lambda_0^{-2}, \quad \text{with } \lambda_0 := \sqrt{\frac{1}{4} + \frac{1}{9}} - \frac{1}{2}.$$

It remains to expand z_1 . For this, we have

$$z_1 = \frac{1}{2} (\sqrt{1 + z} - 1) = \frac{1}{2} \left(\frac{z}{2} + \zeta_3 \right), \quad |\zeta_3| \leq \sum_{k \geq 2} \left| \binom{k}{1/2} \right| z^k.$$

Since $|\binom{k}{1/2}| \leq 1/4$ for all $k \geq 1$, it follows that

$$|\zeta_3| \leq \frac{1}{4} \sum_{k \geq 2} z^k = \frac{z^2}{4(1-z)} \leq \frac{36}{5r^4}.$$

Hence,

$$\begin{aligned} \log \alpha &= \log r + \frac{z}{4} + \left(\zeta_2 + \frac{\zeta_3}{2} \right) =: \log r + \frac{1}{r^2} + \zeta', \\ |\zeta'| &< \left(\frac{c_2}{2} + \frac{36}{10} \right) \frac{1}{r^2} < \frac{3.64}{r^4}. \end{aligned}$$

□

The following estimate is the main result of this section.

Lemma 11 *If $r \geq 4$, then $\kappa > 0$. Furthermore,*

$$x > \frac{\kappa r^2 \log r + 1}{1 + 5/r}. \quad (48)$$

Proof We shall use the approximations given in Lemma 10 but we also need an approximation of $\log U_{n+1}$. We have

$$\log U_{n+1} = \log \left(\frac{\alpha^{n+1}}{\alpha - \beta} \left(1 - \left(\frac{\beta}{\alpha} \right)^{n+1} \right) \right) = (n+1) \log \alpha - \log(\sqrt{r^2 + 4}) + \zeta'', \quad (49)$$

where

$$\zeta'' = \log \left(1 - \left(\frac{\beta}{\alpha} \right)^{n+1} \right).$$

Since $\beta = -\alpha^{-1}$, it follows that $|\beta/\alpha| = 1/\alpha^2$. Thus,

$$|\zeta''| \leq \frac{1}{\alpha^{2n+2}} \left(1 + \sum_{k \geq 1} \frac{1}{k(\alpha^{2n+2})^k} \right) \leq \frac{1}{\alpha^{2n+2}} \left(1 + \frac{1}{2(1 - 1/\alpha^{2n+2})} \right) < \frac{1.51}{\alpha^{2n+2}}, \quad (50)$$

where for the last inequality we used the fact that $\alpha > r \geq 3$ and $n \geq 2$. Inserting estimates (46) and (47) together with Eq. (49) into inequality (27), we get

$$\begin{aligned} |\Gamma| &= \left| m \log \alpha - \log(\sqrt{r^2 + 4}) - x((n+1) \log \alpha - \log(\sqrt{r^2 + 4}) + \zeta'') \right| \\ &= \left| (m - x(n+1)) \log \alpha + (x-1) \log(\sqrt{r^2 + 4}) - x \zeta'' \right| \\ &= \left| (m - x(n+1)) \left(\log r + \frac{1}{r^2} + \zeta' \right) + (x-1) \left(\log r + \frac{2}{r^2} + \zeta \right) - x \zeta'' \right| \end{aligned}$$

$$= \left| (m - nx - 1) \log r + \frac{(m - nx - 1) + (x - 1)}{r^2} + (m - x(n + 1))\zeta' + (x - 1)\zeta + x\zeta'' \right|.$$

We recognise the coefficient of $\log r$ as the number we denoted $-\kappa$ in (45). Using inequality (27), we get

$$\left| -\kappa \log r + \frac{-\kappa + (x - 1)}{r^2} \right| < \frac{2.2}{r^x} + |m - x(n + 1)||\zeta'| + |x - 1||\zeta| + x|\zeta''|.$$

Inequality (19) shows that $m - x(n + 1) \in [-2x + 2, 2]$. In particular, $|m - x(n + 1)| \leq 2(x - 1)$. We thus get, by estimates (46) and (47) together with Eq. (49), that

$$\left| -\kappa \log r + \frac{-\kappa + (x - 1)}{r^2} \right| < \frac{4}{r^x} + \frac{7.28(x - 1)}{r^4} + \frac{5.81(x - 1)}{r^4} + \frac{1.51x}{r^6}.$$

Since $x \geq 3$ and $r \geq 3$, we get that the last term satisfies

$$\frac{1.51x}{r^6} \leq \left(\frac{1.51 \times (3/2)}{3^2} \right) \frac{(x - 1)}{r^4} < \frac{0.26(x - 1)}{r^4}.$$

Hence,

$$\left| -\kappa \log r + \frac{-\kappa + (x - 1)}{r^2} \right| < \frac{2.2}{r^x} + \frac{13.35(x - 1)}{r^4}. \quad (51)$$

Assume that $\kappa \leq 0$. We then get that

$$\frac{x - 1}{r^2} \leq \frac{2.2}{r^x} + \frac{13.35(x - 1)}{r^4},$$

which implies that

$$1 \leq \frac{13.35}{r^2} + \frac{2.2}{r^{x-2}(x - 1)}.$$

If $r \geq 5$, the right-hand side is

$$\leq \frac{13.35}{25} + \frac{2.2}{5 \times 4} = \frac{13.54}{16} < 1,$$

a contradiction. Similarly, if $r \geq 4$ and $x \geq 4$, then the right side is

$$\leq \frac{13.25}{16} + \frac{2.2}{16 \times 3} < \frac{14}{16} < 1.$$

Thus, if $r \geq 5$ or $r = 4$ and $x \geq 4$, then $\kappa > 0$ and now Lemma 9 applies. We will show at the end of this proof that $\kappa > 0$ for $(r, x) = (4, 3)$ as well. Multiplying both sides of

estimate (51) by r^2 , we get

$$| -\kappa(r^2 \log r + 1) + (x - 1) | < (x - 1) \left(\frac{13.35}{r^2} + \frac{2.2}{r^{x-2}(x-1)} \right).$$

Hence,

$$\kappa(r^2 \log r + 1) < (x - 1) \left(1 + \left(\frac{13.35}{r^2} + \frac{2.2}{r^{x-2}(x-1)} \right) \right) \leq (x - 1) \left(1 + \frac{5}{r} \right),$$

which gives estimate (48). It remains to treat the case $(r, x) = (4, 3)$. By inequality (19), we have $m < 3(n + 1) + 1 = 3n + 4$ so $\kappa = 3n + 1 - m \geq -2$. So, the only instances in which $\kappa \leq 0$ is possible are when $m = 3n + 3$, $m = 3n + 2$, $3n + 1$. Well, let us show that this is not possible by proving that

$$U_n^3 + U_{n+1}^3 < U_{3n+1}.$$

Using the Binet formula (4), this is implied by

$$\alpha^{-1} \left(1 + \frac{1}{\alpha^{6n}} \right)^3 + \alpha^2 \left(1 + \frac{1}{\alpha^{6n+6}} \right)^3 < \Delta \left(1 - \frac{1}{\alpha^{6n+3}} \right),$$

with $\alpha = 2 + \sqrt{5}$ and $\Delta = 20$. The function of n in the left is decreasing and the function with n in the right is increasing, and the inequality holds at $n = 1$ (the left-hand side there is < 18.5 and the right side is > 19.5), so it holds for all $n \geq 1$. Thus, $\kappa \geq 2$ for $r = 4$ as well. $\square$

3.8 The case $n \leq 100$

We first seek bounds on r . Having the bounds in r and n , we get bounds on x using Lemma 8. Finally, for a fixed r we use Baker–Davenport on estimate Eq. (27) to lower x . The hope is that in all cases $x \leq 100$, a case which has already been treated.

We prove the following result.

Lemma 12 *When $n \leq 100$, we have $r \leq 1.5 \times 10^6$.*

Proof We assume $r > 10^6$. Then $x > \kappa r^2 \log r / 1.01 \geq nr^2 \log r / 2.02$ by Lemmas 9 and 11. We go through the three possibilities of Lemma 8. In case (i), we get

$$r^2 \log r \leq 1.01 \times 3.3 \times 10^6 (\log(r + 1))^2,$$

which gives $r < 5500$, a contradiction. Assume we are in case (ii). Then

$$x + 1 < 1.38 \times 10^6 \times 1.001 (\log(r + 1))^2 \log \left(\frac{0.3(n + 1)(x + 1)^2}{(\log(r + 1))^2} \right). \quad (52)$$

The factor 1.001 is an upper bound on the factor $1 + 1/(r \log r)$ which is valid since r is large. Now

$$x + 1 > x > \frac{nr^2 \log r}{2.02} > nr^2 \log(r + 1) \left(\frac{\log r}{2.02 \log(r + 1)} \right) > \frac{nr^2 \log(r + 1)}{2.03},$$

where the last inequality holds since $r > 10^6$. Put $y := (x + 1)/(n \log(r + 1))$. Then the above inequality is $y > r^2/2.03$. Inequality (52) can be rewritten in terms of y as

$$\begin{aligned} y &< 1.38 \times 10^6 \log(r + 1) \times 1.001 \left(\log(0.3n^2(n + 1)y^2) \right)^2 \\ &< 1.39 \times 10^6 \log(r + 1) (2 \log y + \log(30,030))^2. \end{aligned}$$

We look at the function

$$f(y) := \frac{y}{(2 \log y + \log(30,030))^2}.$$

Its derivative is

$$\frac{2 \log y + \log(30,030) - 4}{(2 \log y + \log(30,030))^3} > 0,$$

so our function $f(y)$ is increasing. Since $f(y) < 1.39 \times 10^6 \log(r + 1)$, and $y > r^2/2.03$, it follows that $f(r^2/2.03) < 1.39 \times 10^6 \log(r + 1)$. This gives

$$r^2 < 1.39 \times 2.03 \times 10^6 (\log(r + 1)) (2 \log(r^2/2.03) + \log(30,030))^2,$$

which gives $r < 370000$, a contradiction. Assume we are in case (iii). We use the same substitution $y := (x + 1)/(n \log(r + 1))$. We then get

$$\begin{aligned} y &< 1.64 \times 10^6 (\log(r + 1))^2 (\log(0.12n^2y^2))^2 \leq 1.64 \\ &\times 10^6 (\log(r + 1))^2 (2 \log y + \log(12))^2. \end{aligned}$$

The function $g(y) := y/(2 \log y + \log(12))^2$ is also increasing, so we deduce that

$$r^2 < 2.03 \times 1.64 \times 10^6 (\log(r + 1))^2 (2 \log(r^2/2.03) + \log(12))^2,$$

and this gives $r < 1.5 \times 10^6$. □

Having bounds on n and r , inequalities (i), (ii) and (iii) from Lemma 8 become

$$\begin{aligned} x &< 3.3 \times 10^6 (\log(1.5 \times 10^6))^2 < 7 \times 10^8, \\ x &< 1.39 \times 10^6 \times 100 (\log(1.5 \times 10^6))^2 \left(\log(0.3 \times 101(x + 1)^2) \right)^2 \\ &< 3.7 \times 10^{10} \log(30.3(x + 1))^2, \end{aligned}$$

$$\begin{aligned} x &< 1.64 \times 10^6 \times 100 \times (\log(1.5 \times 10^6))^3 (\log(0.12(x+1)))^2 \\ &< 6.5 \times 10^{11} \log(0.12(x+1))^2. \end{aligned}$$

Any one of these inequalities implies that $x < 3 \times 10^{15}$. Now we do Baker–Davenport on estimates (27) for $n \in [2, 100]$, $r \in [3, 1,500,000]$, and $x < 3 \times 10^{15}$. This also gives $m < 3 \times 10^{17}$ via inequality (19). We return to inequality (27) and rewrite it as follows:

$$\left| x \frac{\log U_{n+1}}{\log \alpha} - m + \frac{\log(\sqrt{r^2+4})}{\log \alpha} \right| < \frac{2.2}{r^x \log \alpha}. \quad (53)$$

Then, we apply Lemma 5 on Eq. (53) with the data:

$$M := 3 \times 10^{17}, \quad \tau := \frac{\log U_{n+1}}{\log \alpha}, \quad \mu := \frac{\log(\sqrt{r^2+4})}{\log \alpha}, \quad A := \frac{2.2}{\log \alpha}, \quad \text{and} \quad B := r.$$

A computer search in *Mathematica* reveals that $x \leq 81$, which is a contradiction. This computation lasted 16 hours on a cluster of four 16 GB RAM computers.

3.9 The case $n > 100$

Estimate (27) together with estimate (49) give

$$|\Gamma| = |m \log \alpha - \log(\sqrt{r^2+4} - x((n+1) \log \alpha - \log(\sqrt{r^2+4} + \zeta''))| < \frac{2.2}{r^x},$$

which implies, via estimate (50), that

$$|(m - x(n+1)) \log \alpha + (x-1) \log(\sqrt{r^2+4})| < \frac{2.2}{r^x} + \frac{1.51x}{\alpha^{2n+2}}. \quad (54)$$

If $r \geq 4$ then $\kappa > 0$, so by Lemma 9, we have $\kappa \geq 2$. If $\kappa \geq 3$, then

$$x \geq \frac{\kappa(r^2 \log r + 1)}{1 + 5/r} > \frac{n4^2 \log 4}{4.5} > 2n + 2.$$

The same conclusion holds for $\kappa = 2$ since then $x \geq r^{\min\{2, n-3\}} \geq 4^{\min\{2, n-3\}} \geq 2n + 2$ for all $n \geq 2$. We thus get

$$|(m - x(n+1)) \log \alpha + (x-1) \log(\sqrt{r^2+4})| < \frac{4}{r^{2n+2}} + \frac{1.51x}{\alpha^{2n+2}} < \frac{1.51x + 2.2}{\alpha^{2n+2}}. \quad (55)$$

We prove the following lemma.

Lemma 13 *For $n > 100$, we have $r^{n-2} > x$.*

Note that Lemmas 9 and 13 show that the case $\kappa = 2$ cannot occur for $r \geq 4$ provided that $n > 100$.

Proof Assume $x \geq r^{n-2}$. We use the bounds given by Lemma 8 on x . In case (i), we get

$$r^{99} \leq r^{n-2} \leq 3.3 \times 10^6 (\log(r+1))^2.$$

The function $y \mapsto y^{99}/(\log(y+1))^2$ is increasing for all $y \geq 3$ as one can check by computing its derivative. Thus, if the above inequality holds for r , it should hold $r = 3$ as well, which is false. In case (ii), we have

$$x < 1.38 \times 10^6 n \left(1 + \frac{1}{r \log r}\right) (\log(r+1))^2 \left(\log\left(\frac{0.3(n+1)(x+1)^2}{(\log(r+1))^2}\right)\right)^2.$$

The expression $1 + 1/(r \log r)$ is smaller than 1.304 at $r = 3$. Since $1.38 \times 1.304 < 1.8$, we get

$$x < 1.8 \times 10^6 (\log(r+1))^2 \left(\log\left(\frac{0.3(n+1)(x+1)^2}{(\log(r+1))^2}\right)\right)^2.$$

If $x \leq n$, then we get $r^{n-2} \leq n$ for $r \geq 3$ and $n \geq 101$, which is false. Thus, $n < x$, so we may use $0.3(n+1) < n < x+1$ to get

$$\begin{aligned} x+1 &< 1.8 \times 10^6 n (\log(r+1))^2 (\log((x+1)^3))^2 \\ &< 1.8 \times 3^2 \times 10^6 (\log(r+1))^2 (\log(x+1))^2 \\ &< 1.7 \times 10^7 (\log(r+1))^2 (\log(x+1))^3. \end{aligned}$$

The function $(x+1)/(\log(x+1))^3$ is increasing for $x+1 > e^3$, which is the case for us since $x \geq r^{99} \geq 3^{99}$. Hence, the above inequality should hold for $x+1$ replaced by r^{n-2} , and that gives

$$r^{n-2} \leq 1.7 \times 10^7 (\log(r+1))^2 (\log(r^{n-2}))^2 < 1.7 \times 10^7 n^2 (\log(r+1))^4.$$

Since $r^{n/3} \geq 3^{n/3} > n$ holds for $n > 100$, we get that

$$r^{n/3-2} < 1.7 \times 10^7 (\log(r+1))^4,$$

so

$$r^{95} \leq r^{n-6} < \left(1.7 \times 10^7 (\log(r+1))^2\right)^3 < 5 \times 10^{21} (\log(r+1))^{12}.$$

The function $y \mapsto y^{95}/(\log(y+1))^6$ is increasing for $y \geq 3$, so the last inequality should hold also for r replaced by 3, which is false. A similar argument works if x is in case (iii) of Lemma 8. We don't give further details. $\square$

From Lemma 13 we conclude that if $r \geq 4$, then inequality (55) leads to

$$|(m - x(n+1)) \log \alpha + (x-1) \log(\sqrt{r^2+4})| < \frac{1}{r^n}. \quad (56)$$

We put

$$\Gamma_2 := (x - 1) \log(\sqrt{r^2 + 4}) - ((n + 1)x - m) \log \alpha.$$

We apply Theorem 3 to find a lower bound on $\log |\Gamma_2|$ with the data:

$$t := 2, \quad \gamma_1 := \sqrt{r^2 + 4}, \quad \gamma_2 := \alpha, \quad b_1 := x - 1, \quad b_2 := m - (n + 1)x. \quad (57)$$

Since $\gamma_1, \gamma_2 \in \mathbb{Q}(\alpha)$, we take again $\mathbb{K} := \mathbb{Q}(\alpha)$ with degree $\mathcal{D} := 2$. The fact that γ_1 and γ_2 are multiplicatively independent has already been checked.

We take

$$\max \left\{ h(\gamma_1), \frac{|\log \gamma_1|}{2}, \frac{1}{2} \right\} = \frac{1}{2} \log(r^2 + 4) < \log(r + 1) := \log B_1,$$

and

$$\max \left\{ h(\gamma_2), \frac{|\log \gamma_2|}{2}, \frac{1}{2} \right\} = \frac{1}{2} \log \alpha < \frac{1}{2} \log(r + 1) := \log B_2.$$

Thus,

$$b' := \frac{|b_1|}{D \log B_2} + \frac{|b_2|}{D \log B_1} = \frac{x - 1}{2 \log(r + 1)} + \frac{|(n + 1)x - m|}{\log(r + 1)} < \frac{2.5x}{\log(r + 1)},$$

where we used the fact that $m - (n + 1)x \in [-2x + 2, 2]$. By Theorem 3, we get that

$$\log |\Gamma_2| > -195 \left(\max \left\{ \log \left(\frac{2.5x}{\log r} \right), 10.5 \right\} \right)^2 (\log(r + 1))^2. \quad (58)$$

We want an upper bound on r . So, assume $r \geq 10^6$. If $\log((2.5x)/\log(r + 1)) < 10.5$, then

$$x < \frac{e^{10.5} \log(r + 1)}{2.5} < 15,000 \log(r + 1).$$

Since $x > nr^2 \log r / (2 + 10/r) \geq 101r^2 \log r / 2.02 = 50r^2 \log r$, we get

$$50r^2 \log r \leq x \leq 15,000(\log(r + 1)),$$

so

$$r^2 < \left(\frac{15,000}{50} \right) \left(\frac{\log(r + 1)}{\log r} \right) < 300 \times 1.01 = 303,$$

so $r \leq 17$, a contradiction.

Assume next that $\log(2.5x/\log(r+1)) > 10.5$. We then get

$$\log |\Gamma_2| > -195(\log(r+1))^2 \left(\log \left(\frac{2.5x}{\log(r+1)} \right) \right)^2.$$

Comparing the above inequality with estimate (56), we get

$$n \log r < 195(\log(r+1))^2 \left(\log \left(\frac{2.5x}{\log(r+1)} \right) \right)^2,$$

so

$$\begin{aligned} n &< 195 \left(\frac{\log(r+1)}{\log r} \right) \log(r+1) \left(\log \left(\frac{2.5x}{\log(r+1)} \right) \right)^2 \\ &< 196 \log(r+1) \left(\log \left(\frac{2.5x}{\log(r+1)} \right) \right)^2, \end{aligned}$$

where we used the fact that $r \geq 10^6$ so $\log(r+1)/\log r < 1.0001$.

We now use the bounds on x given by Lemma 8. In case (i), we have

$$50r^2 \log r \leq x \leq 3.3 \times 10^6 (\log(r+1))^2,$$

so

$$r^2 < \left(\frac{3.3 \times 10^6}{50} \right) \times \left(\frac{\log(r+1)}{\log r} \right) \log(r+1) < 6.7 \times 10^4 \log(r+1),$$

so $r < 670$, a contradiction. In case (ii), we have

$$x+1 < 1.39 \times 10^6 n (\log(r+1))^2 \left(\log \left(\frac{0.3(n+1)(x+1)^2}{(\log(r+1))^2} \right) \right)^2.$$

In case $(x+1)/(\log(r+1)) \leq n$, we have

$$x+1 < 196(\log(r+1))^2 \left(\frac{2.5x}{\log(r+1)} \right)^2.$$

So, putting $y := x/\log(r+1)$, we get $y < 196 \log(r+1) \log(2.5y)^2$. Note that

$$y = \frac{(x+1)}{\log(r+1)} > \frac{nr^2 \log r}{2.01 \log(r+1)} \geq \frac{101r^2}{2.01(\log(r+1)/\log r)} > \frac{101r^2}{2.02} = 50r^2.$$

In the above inequalities we used the fact that $r \geq 10^6$. The function $y \mapsto y/\log(2.5y)^2$ is increasing for $y > e^2/2.5$, which is our case. Hence, the inequality $y < 196 \log(r+1) \log(2.5y)^2$ should hold with y replaced by $50r^2$, which yields

$50r^2 < 196 \log(r+1) \log(2.5 \times 50r^2)^2$ and gives $r < 50$, a contradiction. Thus, $n < (x+1)/\log(r+1)$. Since $0.3(n+1) < n$, we conclude that

$$x+1 < 1.39 \times 10^6 (\log(r+1))^2 \left(196 \log(r+1) \left(\log \left(\frac{2.5x}{\log(r+1)} \right) \right)^2 \right) \\ \times \left(\log \left(\left(\frac{(x+1)}{\log(r+1)} \right)^3 \right) \right)^2.$$

Putting again $y := (x+1)/\log(r+1)$, we get

$$y < 1.39 \times 10^6 \times 196 \times (\log(r+1))^2 \log(2.5y)^2 (3 \log y)^2 < 2.46 \times 10^9 (\log(2.5y))^4.$$

The function $y \mapsto y/(\log(2.5y))^4$ is increasing for our range for $y > 50r^2 > 50 \times 10^{16}$, so we get that the above inequality should hold by replacing y by $50r^2$. Thus,

$$50r^2 < 2.46 \times 10^9 (\log(r+1))^2 (\log(2.5 \times 50r^2))^4,$$

so $r < 2.6 \times 10^8$. A similar argument holds when x is in case (iii). In that case, we may again suppose that $n < (x+1)/\log(r+1)$. We get

$$x+1 < 1.64 \times 10^6 (\log(r+1))^3 \left(196 \log(r+1) \left(\log \left(\frac{2.5x}{\log(r+1)} \right) \right)^2 \right) \\ \times \left(\log \left(\frac{0.12(x+1)}{(\log(r+1))^2} \right) \right)^2,$$

so

$$y < 3.3 \times 10^8 (\log(r+1))^3 (\log(2.5y))^2 (\log(0.12y))^2.$$

Imposing that the above inequality holds for y replaced by $50r^2$, we get

$$50r^2 < 1.3 \times 10^9 (\log(r+1))^3 (\log(2.5 \times 50r^2)) (\log(0.12 \times 50r^2))^2,$$

so $r < 4.3 \times 10^8$.

To summarise, we have proved the following.

Lemma 14 *If $r \geq 4$, then $r < 4.3 \times 10^8$.*

Having bounds for r it is easy to find bounds for x . For example,

$$n < 195 \log(r+1) (\log(2.5x))^2 < 195 \log(4.3 \times 10^8) (\log(2.5x))^2 < 3900 (\log(2.5x))^2.$$

Next, if x is in case (i), then

$$x < 3.3 \times 10^6 (\log(r+1))^2 < 3.3 \times 10^6 (\log(4.3 \times 10^8))^2 < 1.4 \times 10^9.$$

If x is in case (ii), then

$$\begin{aligned} x &< 1.39 \times 10^6 (\log(4.3 \times 10^8))^2 (3900(\log(2.5x))^2) \\ &\quad \times (\log(0.3 \times 3900(\log(2.5x))^2(x+1)^2))^2 \\ &< 2.2 \times 10^{12} (\log(2.5x))^2 (\log(1200(x+1)^2 \log(2.5x)))^2, \end{aligned}$$

which gives $x < 5 \times 10^{19}$. Finally, if x is as in case (iii), then

$$\begin{aligned} x &< 1.64 \times 10^6 (\log(4.3 \times 10^8))^3 \times 3900(\log(2.5x))^2 \times (\log(0.12(x+1)))^2 \\ &< 5.1 \times 10^{13} (\log(2.5x))^2 (\log(0.12(x+1)))^2, \end{aligned}$$

so $x < 3 \times 10^{20}$.

Thus, $r < 4.3 \times 10^8$ and $x < 3 \times 10^{20}$. Inequality (56) gives that

$$\left| \frac{\log \sqrt{r^2 + 4}}{\log \alpha} - \frac{x(n+1) - 1}{x - 1} \right| < \frac{1}{r^n(x-1) \log \alpha} < \frac{1}{16(x-1)^2}, \quad (59)$$

where for the last inequality we used that $r^n = r^2 r^{n-2} \geq 16x$ by Lemma 13. In particular, the ratio $(x(n+1)-1)/(x-1)$ is a convergent of $\log \sqrt{r^2 + 4} / \log \alpha$. Since $x < 3 \times 10^{20} < F_{100}$, it follows that $(x(n+1)-1)/(x-1) = p_k/q_k$ for some $k \in [0, 99]$. So, we apply Lemma 4 on Eq. (59) with the data:

$$M := 3 \times 10^{20}, \quad \tau := \frac{\log \sqrt{r^2 + 4}}{\log \alpha}, \quad u := x(n+1) - 1, \quad \text{and} \quad v := x - 1.$$

With the help of a computer search in *Mathematica*, we checked all these possibilities over all the values for $4 \leq r \leq 4.3 \times 10^8$ and found that $n \leq 46$, which is a contradiction. This computation lasted 6 hours on a cluster of four 16 GB RAM computers.

3.10 The case $r = 3$

The case $r = 3$ is special since there we don't know that $\kappa > 0$ so some of the inequalities used for the case $r \geq 4$ do not apply. In the case $n \leq 100$, Lemma 8 gives

$$x < 3.3 \times 10^6 \times (\log 4)^2 < 6.4 \times 10^6,$$

or

$$\begin{aligned} x &< 1.38 \times 10^8 \times 100 \left(1 + \frac{1}{3 \log 3} \right) (\log 4)^2 \left(\log \left(\frac{0.3 \times 101(x+1)^2}{(\log 4)^2} \right) \right), \\ &< 3.5 \times 10^{10} (\log(16(x+1)^2))^2, \end{aligned}$$

which gives $x < 1.7 \times 10^{14}$, or

$$\begin{aligned}
 x &< 1.63 \times 10^6 \times 100 \left(1 + \frac{1}{3 \log 3}\right) (\log 4)^3 \left(\log \left(\frac{0.12(x+1)}{(\log 4)^2}\right)\right)^2 \\
 &< 5.7 \times 10^8 (0.07(x+1))^2,
 \end{aligned}$$

which gives $x < 3.3 \times 10^{11}$. Now we can do Baker–Davenport on estimate (27) and get better bounds on x . In case $n > 100$, estimate (54) together with Lemma 13 hold and give

$$\begin{aligned}
 |(m - x(n+1)) \log \alpha + (x-1) \log(\sqrt{r^2+4})| &< \frac{2.2}{r^x} + \frac{1.51x}{\alpha^{2n+2}} \\
 &< \frac{2.2}{r^x} + \frac{1.51}{r^{n+2}} < \frac{1}{3^{\min\{n-1, x-1\}}}.
 \end{aligned} \tag{60}$$

We keep the notation r and α although this section only applies to $r = 3$ for which $\alpha = (3 + \sqrt{13})/2$. Put $\ell := \min\{n-1, x-1\}$. The lower bound estimate (58) still applies and gives

$$\ell \log 3 < -\log |\Gamma_2| < 195 \left(\max \left\{ \log \left(\frac{2.5x}{\log 4} \right), 10.5 \right\} \right)^2 (\log 4)^2.$$

When $\ell = n-1$, we get

$$(n-1) \log 3 < 375 \left(\max \left\{ \log \left(\frac{2.5x}{\log 4} \right), 10.5 \right\} \right)^2.$$

In case the maximum above is 10.5 we get $x < 16,000$ and $n < 40,000$. If the maximum above is not at 10.5, we then get

$$n < \frac{375}{\log 3} \left(\log \left(\frac{2.5x}{\log 4} \right) \right)^2 + 1 < 350(\log(2x))^2.$$

Going via the possibilities (i), (ii), (iii), we get

$$\begin{aligned}
 x &< 3.3 \times 10^6 (\log 4)^2 < 7 \times 10^6; \\
 x &< 1.38 \times 10^6 \left(350(\log(2x))^2 \right) \left(1 + \frac{1}{3 \log 3} \right) (\log 4)^2 \\
 &\quad \times \left(\log \left(\frac{0.3 \times 351(\log 2x)^2(x+1)^2}{(\log 4)^2} \right) \right)^2 \\
 &< 1.3 \times 10^9 (\log 2x)^2 (\log(55(x+1)^2))^2,
 \end{aligned}$$

which gives $x < 2 \times 10^{16}$, or finally

$$\begin{aligned}
 x &< 1.63 \times 10^6 (350(\log(2x))^2) \left(1 + \frac{1}{3 \log 3} \right) (\log 4)^3 \left(\log \left(\frac{0.12(x+1)}{(\log 4)^2} \right) \right)^2 \\
 &< 2 \times 10^9 (\log(2x))^2 (\log(0.07(x+1)))^2,
 \end{aligned}$$

which gives $x < 3.1 \times 10^{15}$. So, in all instances, $x < 2 \times 10^{16}$, and now

$$n < 350(\log(2x))^2 < 6 \times 10^5.$$

Since Lemma 13 still applies, it follows that estimate (60) gives

$$\left| \frac{\log \sqrt{r^2 + 4}}{\log \alpha} - \frac{x(n+1) - m}{x-1} \right| < \frac{1}{3^{n-1}(x-1)(\log 3)} < \frac{1}{3(x-1)^2},$$

so again $(x(n+1) - m)/(x-1) = p_k/q_k$ is a convergent of $\log \sqrt{r^2 + 4}/\log \alpha$ with $x < 2 \times 10^{16} < F_{80}$, so $k \in [0, \dots, 16]$. So, everything works fine if $\ell = n-1$.

In case $\ell = x-1$, one gets

$$(x-1) \log 3 < 375 \left(\max \left\{ \log \left(\frac{2.5x}{\log 4} \right), 10.5 \right\} \right)^2,$$

which gives $x < 5 \times 10^4$. And one wonders how one should finish it off. We can expand another linear form in logarithms which is small, or we may recall the following main theorem from [2].

Theorem 5 Assume that $s \notin \{1, 2, 4\}$ is minimal such that $U_m \mid U_{n+1}^s - U_n^s$. Then $m < 20,000s^2$.

In our instance, since $U_m = U_n^x + U_{n+1}^x$, one checks that the minimal s is exactly $2x$. Thus, $m < 80,000x^2$, and since $m > (n-1)x$ by estimate (19), we get $n \leq 80,000x < 4 \times 10^9$. Thus, $n < 4 \times 10^9$ and $x < 5 \times 10^4$. It is still a large range and we need to reduce it.

We consider the element

$$y := \frac{x}{\alpha^{2n}}.$$

Lemma 13 together with the fact that $\alpha > r$ implies that

$$y < \frac{1}{\alpha^n}, \quad (61)$$

where the last inequality holds for all $n > 100$. Now, we write

$$U_n^x = \frac{\alpha^{nx}}{(r^2 + 4)^{x/2}} \left(1 - \frac{(-1)^n}{\alpha^{2n}} \right)^x,$$

and

$$U_{n+1}^x = \frac{\alpha^{(n+1)x}}{(r^2 + 4)^{x/2}} \left(1 - \frac{(-1)^{n+1}}{\alpha^{2(n+1)}} \right)^x.$$

If n is odd, then

$$1 < \left(1 - \frac{(-1)^n}{\alpha^{2n}}\right)^x = \left(1 + \frac{1}{\alpha^{2n}}\right)^x < e^y < 1 + 2y,$$

because y is very small. On the other hand, if n is even, then

$$1 > \left(1 - \frac{(-1)^n}{\alpha^{2n}}\right)^x = \exp\left(x \log\left(1 - \frac{1}{\alpha^{2n}}\right)\right)^x > e^{-2y} > 1 - 2y,$$

because y is very small. Thus, the following inequalities hold in both cases,

$$\left|U_n^x - \frac{\alpha^{nx}}{(r^2 + 4)^{x/2}}\right| < \frac{2y\alpha^{nx}}{(r^2 + 4)^{x/2}},$$

and

$$\left|U_{n+1}^x - \frac{\alpha^{(n+1)x}}{(r^2 + 4)^{x/2}}\right| < \frac{2y\alpha^{(n+1)x}}{(r^2 + 4)^{x/2}}.$$

Now, we return to (3) and rewrite it as

$$\begin{aligned} \frac{\alpha^m - \beta^m}{(r^2 + 4)^{1/2}} &= U_m = U_n^x + U_{n+1}^x \\ &= \frac{\alpha^{nx}}{(r^2 + 4)^{x/2}} + \frac{\alpha^{(n+1)x}}{(r^2 + 4)^{x/2}} \\ &\quad + \left(U_n^2 - \frac{\alpha^{nx}}{(r^2 + 4)^{x/2}}\right) + \left(U_{n+1}^x - \frac{\alpha^{(n+1)x}}{(r^2 + 4)^{x/2}}\right), \end{aligned}$$

or

$$\begin{aligned} &\left|\frac{\alpha^m}{(r^2 + 4)^{1/2}} - \frac{\alpha^{nx}(1 + \alpha^x)}{(r^2 + 4)^{x/2}}\right| \\ &= \left|\frac{\beta^m}{(r^2 + 4)^{1/2}} + \left(U_n^x - \frac{\alpha^{nx}}{(r^2 + 4)^{x/2}}\right) + \left(U_{n+1}^x - \frac{\alpha^{(n+1)x}}{(r^2 + 4)^{x/2}}\right)\right| \\ &< \frac{1}{\alpha^m} + \left|U_n^x - \frac{\alpha^{nx}}{(r^2 + 4)^{x/2}}\right| + \left|U_{n+1}^x - \frac{\alpha^{(n+1)x}}{(r^2 + 4)^{x/2}}\right| \\ &< \frac{1}{\alpha^m} + 2y \left(\frac{\alpha^{nx}(1 + \alpha^x)}{(r^2 + 4)^{x/2}}\right). \end{aligned}$$

Multiplying both sides of the above inequality by $\alpha^{-(n+1)x}(r^2 + 4)^{x/2}$, we obtain that

$$\begin{aligned} \left| \alpha^{m-(n+1)x} (r^2 + 4)^{(x-1)/2} - (1 + \alpha^{-x}) \right| &< \frac{(r^2 + 4)^{x/2}}{\alpha^{m+(n+1)x}} + 2y(1 + \alpha^{-x}) \\ &< \frac{1}{2\alpha^n} + 3y < \frac{3}{\alpha^n}, \end{aligned} \quad (62)$$

so we may divide both sides of it by $1 + \alpha^{-x}$ and get

$$\left| \alpha^{m-(n+1)x} (r^2 + 4)^{(x-1)/2} (1 + \alpha^{-x})^{-1} - 1 \right| < \frac{3}{\alpha^n}.$$

Since $n > 100$, the left-hand side is small (say smaller than $1/2$), so we can pass to a logarithmic form and get

$$|(m - (n + 1)x) \log \alpha + (x - 1) \log(\sqrt{r^2 + 4}) - \log(1 + \alpha^{-x})| < \frac{6}{\alpha^n}. \quad (63)$$

For us, the parameter x is in $[3, 50000]$. Given x , we have $m < 2 \times 10^{14}$. So, this is a suitable inequality to apply Baker-Davenport to. To do so, we rewrite Eq. (63) as

$$\left| ((n + 1)x - m) \frac{\log \alpha}{\log(\sqrt{r^2 + 4})} - (x - 1) + \frac{\log(1 + \alpha^{-x})}{\log(\sqrt{r^2 + 4})} \right| < \frac{6}{\alpha^n \log(\sqrt{r^2 + 4})}. \quad (64)$$

Then, we apply Lemma 5 on Eq. (64) with the data:

$$\begin{aligned} M &:= 2 \times 10^{14}, \quad \tau := \frac{\log \alpha}{\log(\sqrt{r^2 + 4})}, \quad \mu := \frac{\log(1 + \alpha^{-x})}{\log(\sqrt{r^2 + 4})}, \\ A &:= \frac{6}{\log(\sqrt{r^2 + 4})}, \quad B := \alpha. \end{aligned}$$

A computer search in *Mathematica* reveals that $n \leq 94$, which is the final contradiction. This computations lasted a few hours on a cluster four 16 GB RAM computers. $\square$

Acknowledgements This paper was written when both authors visited the Max Planck Institute for Mathematics Bonn, in 2019. They thank this institution for the hospitality and the fruitful working environment.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bilu, Yu., Hanrot, G., Voutier, P.M.: Existence of primitive divisors of Lucas and Lehmer numbers. With an appendix of M. Mignotte. *J. Reine Angew. Math.* **539**, 75–122 (2001)
2. Bilu, Yu., Komatsu, T., Luca, F., Pizarro-Madariaga, A., Stănică, P.: On a divisibility relation for Lucas sequences. *J. Number Theory* **163**, 1–18 (2016)
3. Carmichael, R.D.: On the numerical factors of the arithmetic forms $\alpha^n \pm \beta^n$. *Ann. of Math.* **15**, 30–70 (1913)
4. Dujella, A., Pethő, A.: A generalization of a theorem of Baker and Davenport. *Quart. J. Math. Oxford* **49**, 291–306 (1998)
5. Gómez Ruíz, C.A., Luca, F.: An exponential Diophantine equation related to the sum of powers of two consecutive k -generalized Fibonacci numbers. *Colloq. Math.* **137**, 171–188 (2014)
6. Koshy, T.: *Fibonacci and Lucas Numbers with Applications*. Wiley, Hoboken (2001)
7. Laurent, M., Mignotte, M., Nesterenko, Yu.: Formes linéaires en deux logarithmes et déterminants d'interpolation. *J. Number Theory* **55**, 285–321 (1995)
8. Luca, F., Oyono, R.: An exponential Diophantine equation related to powers of two consecutive Fibonacci numbers. *Proc. Jpn. Acad.* **137**(Ser. A), 45–50 (2014)
9. Mignotte, M.: A kit for linear forms in three logarithms. Preprint (2008)
10. Rihane, S.E., Faye, B., Luca, F., Togbé, A.: An exponential Diophantine equation related to the difference between powers of two consecutive Balancing numbers. *Ann. Math. Inf.* **50**, 167–177 (2019)
11. Rihane, S.E., Faye, B., Luca, F., Togbé, A.: On the exponential Diophantine equation $P_n^x + P_{n+1}^x = P_m$. *Turk. J. Math.* **43**, 1640–1649 (2019)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.