Combined Radio and Space-Based Solar Observations: From Techniques to New Results – Preface

Eduard P. Kontar¹ · Alexander Nindos²

Published online: 19 June 2018
© Springer Nature B.V. 2018

Abstract The phenomena observed at the Sun have a variety of unique radio signatures that can be used to diagnose the processes in the solar atmosphere. The insights provided by radio observations are further enhanced when they are combined with observations from space-based telescopes. This Topical collection demonstrates the power of combination methodology at work and provides new results on i) type I solar radio bursts and thermal emission to study active regions; ii) type II and IV bursts to better understand the structure of coronal mass ejections; and iii) non-thermal gyro-synchrotron and/or type III bursts to improve the characterisation of particle acceleration in solar flares. The ongoing improvements in time, frequency, and spatial resolutions of ground-based telescopes reveal new levels in the complexity of solar phenomena and pose new questions.

Keywords Radio bursts · Flares · Active regions · Corona · Coronal mass ejections

CESRA, the Community of European Solar Radio Astronomers,¹ organises triennial workshops on the investigations of solar atmosphere processes using radio and other observations. The 2016 workshop² had a special emphasis on the complementarity of current and future space-based observations with ground-based radio observations. It was the place to discuss

¹See webpage <http://cesra.net>.

²See <http://cesra2016.sciencesconf.org>.

Combined Radio and Space-based Solar Observations: From Techniques to New Results
Guest Editors: Eduard Kontar and Alexander Nindos

✉ E.P. Kontar
eduard@eris.astro.gla.ac.uk

A. Nindos
anindos@uoi.gr

¹ School of Physics and Astronomy, University of Glasgow, Glasgow, G12 8QQ, UK

² Physics Department, University of Ioannina, 45110 Ioannina, Greece

the new exciting science opportunities that arise from radio instruments such as the *Atacama Large Millimeter/submillimeter Array* (ALMA³), the *Expanded Owens Valley Solar Array* (EOVSA⁴), the *Expanded Very Large Array* (EVLA⁵), the *Low Frequency array* (LOFAR⁶), the *Mingantu Spectral Radioheliograph* (MUSER, see Yan *et al.*, 2009, for details), and the developments of the *Square Kilometre Array* (SKA⁷). The workshop discussions have focused on particle acceleration and transport, the radio diagnostics of coronal mass ejections, fine structures in solar radio bursts, and the radio aspects of space weather. This volume provides a snapshot of the developments and challenges that were discussed during the workshop. This Topical collection covers four sub-topics:

- i) Solar Radio Emission Modelling (Lyubchik *et al.*, 2017; Rodger and Labrosse, 2017; Stupishin *et al.*, 2018; Zaitsev and Stepanov, 2017)
- ii) Solar Flares and Solar Energetic Particles (Anastasiadis *et al.*, 2017; Benz, Battaglia, and Güdel, 2017; Altyntsev *et al.*, 2017)
- iii) Fine Structures in Solar Radio Emission (Mohan and Oberoi, 2017; Mugundhan, Hariharan, and Ramesh, 2017)
- iv) Coronal Mass Ejections (Al-Hamadani, Pohjolainen, and Valtonen, 2017; Kumari *et al.*, 2017; Long *et al.*, 2017; Melnik *et al.*, 2018; Miteva, Samwel, and Costa-Duarte, 2018)

1. Solar Radio Emission Modelling

The complexity of the solar atmosphere as well as the increasing quality of solar radio observations necessitates the development of new more complex models. The modelling of solar atmosphere parameters above sunspots using RATAN-600 microwave observations (Stupishin *et al.*, 2018) inferred the upper transition-region structure of sunspots. The method presented is based on iterative correction of the temperature–height profile in the transition region and lower corona, allowing us to test time-independent models of density and temperature as a function of height. Anticipating future observations in the sub-THz range, Rodger and Labrosse (2017) studied how the ratio of brightness temperatures at two frequencies can be used to estimate the optical thickness and the emission measure for prominences. Highlighting that there is no generally accepted theory that would explain high brightness temperatures in type I storms, Lyubchik *et al.* (2017) proposed a new model to explain type I solar radio bursts associated with active regions. The model is based on the turbulence of kinetic-scale Alfvén waves that produce an asymmetric plateau in the electron velocity and a high level of Langmuir waves, leading to plasma emission. The model proposed by Zaitsev and Stepanov (2017) suggests that the electron acceleration and storage of energetic particles in solar magnetic loops can be better explained by a mechanism based on oscillations of the electric current. Specifically, the model aims to explain synchronous pulsation in a wide frequency interval that is hard to achieve by the sausage and kink magnetohydrodynamic modes.

³See <http://www.almaobservatory.org>.

⁴See <http://ovsa.njit.edu>.

⁵See <http://www.aoc.nrao.edu/evla/>.

⁶See <http://www.lofar.org>.

⁷See <https://www.skatelescope.org>.

2. Solar Flares and Solar Energetic Particles

Solar flares are well known for the efficient acceleration of non-thermal electrons and hence are a source of various solar radio bursts (see Nindos *et al.*, 2008, for a review). However, Benz, Battaglia, and Güdel (2017) have demonstrated that there are exceptions to this rule and presented observations of a radio-quiet solar pre-flare, which suggests that acceleration to relativistic energies, if any, should be occurring with low efficiency and does not lead to observable radio emission. Studying optically thin gyrosynchrotron emission, Altyntsev *et al.* (2017) reported an unusual flare, whose emission displayed an apparently ordinary-mode polarisation, in contrast to the classical theory that favours the extraordinary mode. This apparent ordinary-wave emission in the optically thin mode has been attributed to radio wave propagation across the quasi-transverse layer that changes the radio-wave polarisation.

Solar flares are often associated with energetic particle events (SEPs) observed near Earth. These SEP events are an important element of space weather, and there is growing interest in developing reliable forecasting systems. Anastasiadis *et al.* (2017) presented an integrated prediction system for solar flares and SEP events. The system is based on statistical methods and demonstrated promising results for the expected SEP characteristics.

3. Fine Structures in Solar Radio Emission

High-frequency resolution solar radio telescopes have enabled detailed imaging and spectroscopic studies of the fine structures of solar radio emission (Kontar *et al.*, 2017). For example, type III bursts that sometimes show fine structures (stria) in dynamic spectra (de La Noe and Boischot, 1972) can be used to study density fluctuations. Assuming that the individual stria bandwidth is determined by the amplitude of density fluctuations, Mugundhan, Hariharan, and Ramesh (2017) used the striations in a type III radio burst to determine the electron density variations along the path of the electron beams. The observations of solar bursts in time, frequency, and two spatial coordinates naturally lead to 4D data. Using the Murchison Widefield Array (MWA⁸) radio telescope data, Mohan and Oberoi (2017) implemented a formalism to generate 4D data cubes based on brightness temperature maps.

4. Coronal Mass Ejections

Coronal mass ejections (CMEs) are often associated with type II and type IV radio bursts observed over a wide radio frequency range. Simultaneous radio and white-light observations of CMEs can be used to poorly constrain the strength of the solar coronal magnetic field above $\approx 2R_{\odot}$ (Kumari *et al.*, 2017). Assuming the plasma emission mechanism in type IV radio burst to be associated with a behind-the-limb CME, Melnik *et al.* (2018) have estimated the densities of plasma in the core of the CME. However, the relation between CMEs, global extreme ultraviolet (EUV) waves, and type II solar is not always clear. Long *et al.* (2017) studied over 160 global EIT waves observed in EUV and found no clear relationship between global waves and type II radio bursts. The relation between CMEs and type II events is even more complicated when the Sun launches multiple CMEs. Al-Hamadani, Pohjolainen, and Valtonen (2017) studied type II solar radio bursts that occurred during a multiple CME event and demonstrated that the last type II burst had enhanced emission in a wider

⁸See <http://www.mwatelescope.org/>.

bandwidth, which should be consistent with the CME-CME interaction. To understand the physics of the relation between the phenomena at the Sun and the satellite-damaging proton events at 1 AU, identification of solar flare CMEs that are responsible for the proton events is required. The statistical relationships found by Miteva, Samwel, and Costa-Duarte (2018) serve as a useful tool to diagnose the dependencies and test solar flare and CME models.

The CESRA 2016 workshop took place in Orléans, France. The members of the Scientific Organizing Committee were M. Bartà (Czech Republic), K.-L. Klein (France; co-chair), E.P. Kontar (UK), M. Kretzschmar (France; co-chair), C. Marqué (Belgium), A. Nindos (Greece), S. Pohjolainen (Finland; co-chair), A. Warmuth (Germany), and M.K. Georgoulis (Greece; as president of the European Solar Physics Division). The workshop received financial support from the University of Orléans, the Observatoire de Paris, and the CNRS/INSU.

The Topical collection Editors would like to thank both the authors and the referees of the articles.

References

- Al-Hamadani, F., Pohjolainen, S., Valtonen, E.: 2017, Formation of radio type ii bursts during a multiple coronal mass ejection event. *Solar Phys.* **292**(12), 183. [DOI](#).
- Altıntsev, A., Meshalkina, N., Myshyakov, I., Pal'shin, V., Fleishman, G.: 2017, Flare sol2012-07-06: on the origin of the circular polarization reversal between 17 GHz and 34 GHz. *Solar Phys.* **292**(9), 137. [DOI](#).
- Anastasiadis, A., Papaioannou, A., Sandberg, I., Georgoulis, M., Tziotziou, K., Kouloumvakos, A., Jiggins, P.: 2017, Predicting flares and solar energetic particle events: the forspet tool. *Solar Phys.* **292**(9), 134. [DOI](#).
- Benz, A.O., Battaglia, M., Güdel, M.: 2017, Observations of a radio-quiet solar preflare. *Solar Phys.* **292**(10), 151. [DOI](#).
- de LaNoë, J., Boisshot, A.: 1972, The type III B burst. *Astron. Astrophys.* **20**, 55. [ADS](#).
- Kontar, E.P., Yu, S., Kuznetsov, A.A., Emslie, A.G., Alcock, B., Jeffrey, N.L.S., Melnik, V.N., Bian, N.H., Subramanian, P.: 2017, Imaging spectroscopy of solar radio burst fine structures. *Nat. Commun.* **8**, 1515. [DOI](#). [ADS](#).
- Kumari, A., Ramesh, R., Kathiravan, C., Wang, T.J.: 2017, Strength of the solar coronal magnetic field – a comparison of independent estimates using contemporaneous radio and white-light observations. *Solar Phys.* **292**(11), 161. [DOI](#).
- Long, D.M., Murphy, P., Graham, G., Carley, E.P., Pérez-Suárez, D.: 2017, A statistical analysis of the solar phenomena associated with global euv waves. *Solar Phys.* **292**(12), 185. [DOI](#).
- Lyubchik, O., Kontar, E.P., Voitenko, Y.M., Bian, N.H., Melrose, D.B.: 2017, Solar plasma radio emission in the presence of imbalanced turbulence of kinetic-scale Alfvén waves. *Solar Phys.* **292**(9), 117. [DOI](#).
- Melnik, V.N., Brazhenko, A.I., Konovalenko, A.A., Dorovskyy, V.V., Rucker, H.O., Panchenko, M., Frantsuzenko, A.V., Shevchuk, M.V.: 2018, Decameter type iv burst associated with a behind-the-limb cme observed on 7 November 2013. *Solar Phys.* **293**(3), 53. [DOI](#).
- Miteva, R., Samwel, S.W., Costa-Duarte, M.V.: 2018, The wind/epact proton event catalog (1996–2016). *Solar Phys.* **293**(2), 27. [DOI](#).
- Mohan, A., Oberoi, D.: 2017, 4d data cubes from radio-interferometric spectroscopic snapshot imaging. *Solar Phys.* **292**(11), 168. [DOI](#).
- Mugundhan, V., Hariharan, K., Ramesh, R.: 2017, Solar type iiib radio bursts as tracers for electron density fluctuations in the corona. *Solar Phys.* **292**(11), 155. [DOI](#).
- Nindos, A., Aurass, H., Klein, K.-L., Trottet, G.: 2008, Radio emission of flares and coronal mass ejections. Invited review. *Solar Phys.* **253**, 3. [DOI](#). [ADS](#).
- Rodger, A., Labrosse, N.: 2017, Solar prominence modelling and plasma diagnostics at alma wavelengths. *Solar Phys.* **292**(9), 130. [DOI](#).
- Stupishin, A.G., Kaltman, T.I., Bogod, V.M., Yasnov, L.V.: 2018, Modeling of solar atmosphere parameters above sunspots using ratan-600 microwave observations. *Solar Phys.* **293**(1), 13. [DOI](#).
- Yan, Y., Zhang, J., Wang, W., Liu, F., Chen, Z., Ji, G.: 2009, The Chinese spectral radioheliograph – CSRH. *Earth Moon Planets* **104**, 97. [DOI](#). [ADS](#).
- Zaitsev, V.V., Stepanov, A.V.: 2017, Acceleration and storage of energetic electrons in magnetic loops in the course of electric current oscillations. *Solar Phys.* **292**(10), 141. [DOI](#).