

Identification of *LEC1*, *LIL* and *Polycomb Repressive Complex 2* genes and their expression during the induction phase of *Medicago truncatula* Gaertn. somatic embryogenesis

Anna Orłowska¹ · Rafał Igielski¹ · Katarzyna Łągowska¹ · Ewa Kępczyńska¹

Received: 14 October 2016 / Accepted: 18 December 2016 / Published online: 29 December 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Although somatic embryogenesis (SE), a multi-step process starting from somatic tissues and ending with somatic embryos, has been applied to numerous plants including *Medicago* sp., the molecular basis of development reprogramming in somatic cells toward the embryogenic pathway is still incompletely known. Though recent analysis of the proteome and transcriptome has led to the identification and characterization of new genes involved in SE, lot of these genes are up-regulated only in the late developmental stages. Consequently, this work was aimed at finding out if and when the genetic program changed during the SE induction phase in both the highly embryogenic line M9-10a of *Medicago truncatula* cv. Jemalong and its non-embryogenic predecessor line, M9. Based on multipoint (day 0, 2, 7, 14 and 21) gene expression qPCR analysis of two embryogenesis marker genes *LEC1* and *LIL* and selected genes encoding proteins of PRC2 complex (*CLF*, *SWN*, *FIE*, *MSI1*, *VRN2*) it was possible to distinguish two periods during the induction phase. The first week was related to dedifferentiation with no visible changes in explant morphology and lack of transcripts of *LEC1* and *LIL*; the expression of PRC2 members, however, increased in the embryogenic line. The next two weeks were regarded as the expression phase involving the beginning of a rapid callus growth and the appearance of products for *LEC* genes, which was observed in the embryogenic line only.

However, the callus formation was observed in the non-embryogenic line as well, but the *LEC1* and *LIL* genes were not expressed and the transcription of *PRC2* genes was at stable level. It was only the *SWN* expression that decreased at the beginning of induction and did not change in the subsequent days in both lines. Our results indicate that *LEC1* and *LIL*, known as marker genes of the late developmental stages, may be associated with the acquisition of embryogenic competency by somatic cells (prime events in the SE induction). The *PRC2* complex genes are also expressed during embryogenic callus tissue formation on *Medicago truncatula* leaf explants.

Keywords Expression pattern · Induction of somatic embryogenesis · Legumes · Epigenetic regulation

Abbreviations

CLF	Curly Leaf
FIE	Fertilization-Independent Endosperm
LIL	Leafy Cotyledon1-like
LEC1	Leafy Cotyledon1
MSI1	Multicopy Suppressor of Ira1
PRC2	Polycomb Repressive Complex2
SE	Somatic embryogenesis
SWN	Swinger
VRN2	Vernalization2

Electronic supplementary material The online version of this article (doi:10.1007/s11240-016-1161-8) contains supplementary material, which is available to authorized users.

✉ Ewa Kępczyńska
ekepcz@univ.szczecin.pl

¹ Department of Plant Biotechnology, Faculty of Biology, University of Szczecin, Wąska 13, 71-415 Szczecin, Poland

Introduction

Somatic embryogenesis (SE) is a multi-step in vitro regeneration process in which embryos are formed from somatic cells without the fusion of gametes (Williams and Maheswaran 1986; Zimmerman 1993). To date, such vegetative propagation techniques have been regarded as

the lowest-cost methods for producing uniform, identical plants, particularly of those species that are difficult to propagate in a natural way or in other conventional manner. The process is also used to study the embryonic development regulation at morphological, biochemical, genetic and epigenetic levels (Willemssen and Scheres 2004; Elhiti et al. 2013; Smertenko and Bozhkov 2014). Competence to somatic embryogenesis is highly correlated with the genotype. At last two embryogenic lines, 2HA and M9-10a, are known in *Medicago truncatula* (Nolan et al. 1989; Neves et al. 1999) and are considered as models for study SE in this species. M9-10a line was directly derived from non-embryogenic (M9) as effect of somaclonal variation. This is a good model to compare progress of SE without potential background that result from comparison different genetic origin lines (Almeida et al. 2012). Generally, SE takes a few weeks starting from a single cell or a cell group of somatic explants. Its fundamental phases are distinguished at the morphological level, and are divided into a number of stages: induction of proembryogenic structures, followed by embryo formation, their maturation and dehydration. SE terminates with embryos ready to regenerate a new plant. Each of these phases is regulated by various intrinsic (e.g. developmental stage of the starting explant and hormone levels) as well as extrinsic factors. The latter one particularly a number of physical and chemical treatments applied at appropriate schedules seems to be the critical for successful and efficient somatic embryogenesis. However, among all these factors, plant growth regulators appear to play crucial role in SE. The 2,4-D artificial auxin, the popular SE inducer, is sufficient in many species to initiate a genetic program that leads to the establishment of cell lineages with an altered gene transcription pattern, and a different morphology and developmental fate (Gaspar et al. 1996; Karami et al. 2009). Different groups of genes are known to be expressed and essential during somatic embryogenesis, including *WUSCHEL* (*WUS*; Zuo et al. 2002), *AGAMOUS LIKE-15* (*AGL15*; Harding et al. 2003; Zheng et al. 2013), *BABY-BOOM* (*BBM*; Boutilier et al. 2002) and *LEAFY COTYLEDON* (*LECs*, i.e. *LIL*, *LEC1*, *LEC2*, *FUSCA3*; Gaj et al. 2005; Ledwoń and Gaj 2011).

The *LEC* genes are generally known to play multiple roles during zygotic embryogenesis in the early and late phase of embryonic development when the molecular and cellular environment necessary for the process is being established (Jia et al. 2013). Expression of *LEC1* and its close homolog *LIL* plays an important role in zygotic embryogenesis regulating embryo identity and development (Kwong et al. 2003). The role of these genes is mainly to control seed maturation, to inhibit premature germination, and to identify embryonic organs (Meinke 1992; West et al. 1994). *LECs* are also considered as belonging to the somatic embryogenesis marker genes. One of them,

LEC1, is a CCAAT-binding (CBF) transcription factor with a HAP3 subunit (Lee et al. 2003). The *LEC1* genes expression pattern was analyzed during somatic embryogenesis from different initial explants representing different genetic program such as immature embryos of *Zea mays* and *Arabidopsis thaliana* (Zhang et al. 2002; Gaj et al. 2005), *Coffea canephora* seedling leaves (Nic-Can et al. 2013), *Daucus carota* hypocotyl (Yazawa et al. 2004), and *Medicago sativa* protoplasts (Domoki et al. 2006). LEAFY COTYLEDON 1-LIKE B-domain of HAP3 shared 83% similarity to that of *LEC1*, which means that they define distinct class of HAP3 subunits (Kwong et al. 2003). The *LIL* is crucial for proper embryogenesis performance. Its expression was analyzed during SE induction from single-node *Helianthus annuus* (Fambrini et al. 2006), *Theobroma cacao* staminodes (Alemanno et al. 2008) as well as the stamen and nodal *Vitis vinifera* (Schellenbaum et al. 2008; Maillot et al. 2009). Overexpression of the *LEC1* gene in *Arabidopsis* induced the embryonic process and expressed a set of genes important for embryonic development (Lotan et al. 1998). On the other hand, *LIL* is not regarded as a SE marker, but its overexpression could rescue *lec1* mutant and it is up-regulated by *LEC2*, another key SE transcription factor (Guo et al. 2013).

Epigenetic regulation can be another important player during SE induction, capable of controlling genome-wide changes in gene expression during the cell fate transition from a somatic explant (e.g. leaf blade) to the embryogenic callus (He et al. 2012). To date, three levels of epigenetic control of SE are known: the micro RNA (miRNA) pathway, the DNA methylation, and histone post-translational modifications (Smertenko and Bozhkov 2014). Histone modifications are driven by the Polycomb group proteins (PcG) and are essential for cell fate determination, cellular differentiation and transition through successive stages of development from seed to seed in plants (Hennig and Derkacheva 2009; Bemer and Grossniklaus 2012). In plants they form two main conserved protein complexes: the Polycomb Repressive Complex 1 (PRC1) and the well-characterized PRC2. PRC2 catalyzes trimethylation of histone 3 (H3K27me3) lysine 27 through the SET-domain protein. PRC1 binds to the H3K27me3 and ubiquitinates lysine 119 of histone H2A (H2AK27ub) resulting in a compacted chromatin state (Margueron and Reinberg 2011; He et al. 2012; Gleason and Kramer 2013). The PRC2 complex consists of numerous proteins: two WD40-domain proteins, MULTICOPY SUPPRESSOR OF IRA1 (MSI1) and FERTILIZATION-INDEPENDENT ENDOSPERM (FIE); one of SET-domain proteins: CURLY LEAF (CLF), SWINGER (SWN) or MEDEA (MEA); and one of VEFS-domain proteins: VERNALIZATION2 (VRN2), FERTILIZATION-INDEPENDENT SEED2 (FIS2) or EMBRYONIC FLOWER2 (EMF2). Current data available

for *Arabidopsis thaliana* suggest the existence of different PRC2 complexes in plants: VRN-PRC2, FIS-PRC2 and EMF-PRC2. VRN-PRC2 promotes flowering after vernalization by silencing the expression of the flowering repressor *FLOWERING LOCUS C (FLC)* (Gendall et al. 2001; Jiang et al. 2008; Butenko and Ohad 2011). FIS-PRC2 is the main complex involved in regulation of gametophyte and endosperm development. It is responsible for inhibiting the endosperm development in the absence of fertilization. FIS-PRC2 also suppresses the endosperm and embryo cell proliferation after fertilization (Guitton et al. 2004; Köhler and Makarevich 2006). EMF-PRC2 regulates the transition from the vegetative to the generative phase through suppression of important flowering regulators (Yoshida et al. 2001; Chanvivattana et al. 2004). Both EMF-PRC2 and VRN-PRC2 complexes can together control the development of the sporophyte and have a major role in ensuring differentiation and repressing stem cell genes (Bemer and Grossniklaus 2012). The loss of *Arabidopsis MSII* function results in numerous disorders in the vegetative and embryogenic development, leads to defects in shoot apical meristems, floral meristems, primordia and embryo abortion at different developmental stages (Hennig et al. 2003; Köhler et al. 2003; Bouveret et al. 2006; Schönrock et al. 2006; Steinbach and Hennig 2014). In *msi* mutant plants displayed defects in cell division on early stages of embryogenesis and over-proliferation with deficiency in differentiation of embryos (Köhler et al. 2003; Guitton et al. 2004), indicating an important role of *MSII* in the zygotic embryo development control. Together with *MSII*, *FIE* is an important epigenetic regulator of the reproductive program in *Arabidopsis*. Mutation in the *FIE* gene and the absence of its product results in defects during embryonic development and 50% embryo abortion, similarly to what has been observed in the *msi* mutant (Spillane et al. 2000; Yadegari et al. 2000; Katz et al. 2004). Plants that lost the *FIE* expression show a normal body plan of *A. thaliana* seedlings, but form floral buds very fast and develop copious ectopic cells and organs (Kinoshita et al. 2001; Bouyer et al. 2011).

Information on the contribution of PRC2 to the regulation of somatic embryogenesis induction phase in plants is very scarce. It is only *Arabidopsis* mutants with reduced or inhibited expression of PRC2 complex components that have been studied. These mutations revealed unorganized cell divisions and callus-like tissue development with ability to form somatic embryos (Chanvivattana et al. 2004; Makarevich et al. 2006; Bouyer et al. 2011; Ikeuchi et al. 2015; Mozgova et al. 2015).

To date, identification of *LEC1*, *LIL* and genes coding protein components of the PRC2 complex and their expression during the induction phase of *Medicago truncatula* SE has not been described. Therefore, the work reported

here was aimed at investigate the changes in the expression of two somatic embryogenesis marker genes (*LEC1*, *LIL*) and genes from the PRC2 complex (*CLF*, *SWN*, *FIE*, *MSII*, *VRN2*) genes during the SE induction phase comparing their expression in two genotypes of *M. truncatula* cv. Jemalong: the embryogenic (M9-10a) and the non-embryogenic (M9) line, which could allow obtaining new insights into the involvement of these genes into SE regulation.

Materials and methods

Plant material

Seeds of two genotypes of *Medicago truncatula* cv. Jemalong: the high embryogenic (M9-10a) and the non-embryogenic (M9) line, were kindly provided by Pedro Manuel Feveireiro from Instituto de Tecnologia Quimica e Biologica (ITQB), Portugal. The seeds were used to produce mother plants. Before sowing, the seeds were scarified in 2 ml Eppendorf tubes in 1 ml H₂SO₄ (96%) on ice and shaken for 8 min. After scarification, the seeds were rinsed ten times in cold sterile water and immersed for 30 min in 1 ml gibberellin A₃ (100 µM) solution, then rinsed once with sterile water. Next, the seeds were placed on sterile 15 cm Petri dishes (100 seeds per dish) lined with moisturized filter paper. They were stratified in darkness at 4 °C for 2 days and then transferred to 20 °C for 1 day. Those seeds with well-developed embryonic root (1 cm) were seeded into a mixture of sand, soil, perlite and vermiculite (1:1:1:1). The plants were grown in a growth room at 24 ± 1 °C under a 16 h photoperiod of 70 µM m⁻² s⁻¹ GreenLED (Philips).

Tissue culture protocol

Initial explants for callus induction in both lines of *M. truncatula* were obtained from well-developed trifoliolate leaves from the second to third nodes of 60 day-old mother plants. The leaves were surface-sterilized in 1% sodium hypochlorite (using Domestos) for 5 min and then rinsed three times in sterile water. For callus formation induction, three square-shaped explants of 1 cm side length were cut off from each trifoliolate leaf and placed on Petri dishes (ø 55 mm) filled with the SH medium (Schenk and Hildebrandt 1972) supplemented with 0.5 µM 2,4-D and 1 µM zeatin with 30 g l⁻¹ sucrose. The medium was solidified with 2.5 g l⁻¹ gerlit and adjusted to pH 5.7. The culture was kept at 28 ± 1 °C in darkness for 21 days. Subsequently, the callus tissue was transferred to the MS differentiation medium (Murashige and Skoog 1962) with 30 g l⁻¹ sucrose and 2.5 g l⁻¹ gerlit, pH 5.8. The explants were cultured for 14 days at 24 ± 1 °C under a 16 h photoperiod of 70 µM m⁻² s⁻¹ GreenLED (Philips).

RNA isolation and cDNA synthesis

Samples were collected at four time points (day 2, 7, 14, and 21) during the SE induction phase in both lines, except for day 0 when leaves from intact plants were used. Three biological samples, each consisting of 7 trifoliolate explants representing each time point pooled together were subjected to RNA isolation. Total RNA was isolated from 50 mg of frozen tissues in 1 ml TRIzol Reagent (ZymoResearch) using Direct-zol™ RNA-MiniPrep Kit (ZymoResearch) according to the manufacturer's instructions. DNA contamination was removed by using DNase I (ZymoResearch). RNA was eluted in 30 µl DNase/RNase Free-water. The purity and concentration of RNA was checked with BioSpec-nano (Shimadzu) and by electrophoresis in 2% agarose gel. First-strand cDNA of each sample was synthesized from 500 ng total RNA in a 20 µl reaction volume using the High-Capacity cDNA Reverse Transcription Kit (LifeTechnologies) according to the appropriate protocol and then used for quantitative PCR (qPCR).

Sequence analysis

To perform *Medicago truncatula* gene identification, *Arabidopsis thaliana* reference amino acid sequences of *LEC1*, *LIL*, and *PRC2* genes were obtained from the TAIR data base (<http://www.arabidopsis.org/>) and used to BLAST search against the JCVI Mt 4.0v1 data base (<http://www.jcvi.org/medicago/>) of *M. truncatula*. The candidate genes are summarized in Table 1. Locations of specific domains were confirmed in the InterPro data base. The sequence alignments analysis was used to check the homology of protein sequences. The similar protein sequences obtained from the NCBI data base for *Arabidopsis thaliana*, *Brassica napus*, *Cicer arietinum*, *Glycine max*, *Medicago truncatula*, *Nicotiana tomentosiformis*, *Theobroma cacao*, *Phaseolus vulgaris*, *Ricinus communis*, *Solanum lycopersicum*, and *Vitis vinifera* were used to perform ClustalW alignments. The sequence alignments analysis was performed using the Geneious 6.1 software (<http://www.geneious.com>, Kearse et al. 2012). Phylogenetic unrooted trees were building with the Neighbor-Joining method and Jukes-Cantor genetic distance model. The trees were resampled 1000 times using the bootstrap method.

Quantitative real-time PCR

Gene-specific primers for quantitative real-time PCR (qPCR) were designed using the PrimerExpress® Software v3.0 (LifeTechnologies). All the sequences and parameters are given in Table 1. qPCR was performed with the SYBR Select Master Mix (Applied Biosystems) using the STEP ONE Real-time PCR System (LifeTechnologies) following

the manufacturer's instructions. The 10 µl reaction mixture contained 5 µl SYBR Select Master Mix, 0.2 µl 10 mM primers, 1 µl cDNA template, and 3.8 µl DNase/RNase-free distilled water. The expression profile of selected genes in the M9 and M9-10a lines during the SE induction phase was performed using 1:5 cDNA dilution. Analyses for both lines were run on separate plates. Additionally, the Inter-Plate Calibrator analysis was performed on each plate according to the GenEX user guide to compare profiles on one plot. To confirm the changes in transcription level at day 7 and 14, 1:3 cDNA dilutions was used and the analyses for both lines were performed on a single plate. Three biological replicates of each time point and three technical ones were analyzed. The qPCR reaction conditions were as follows: initiation at 95 °C for 2 min, followed by 40 cycles of amplification with 15 s at 95 °C for denaturation and 1 min at 60 °C for annealing. The final extension was performed at 60 °C for 1 min. The dissociation curves were analyzed to check for gene-specific amplification; no unspecific products were detected. The reaction efficiency was 95–100%, as tested using a standard curve for each primer pair. Based on the existing bibliography (Kakar et al. 2008; Mantiri et al. 2008; Pérez et al. 2015) selected 5 candidate reference genes and constructed onsite geNorm and NormFinder evaluation within them (supplement 1). For further analysis used *ACTIN2* as reference gene. For each gene, the relative transcript abundance was calculated and expressed as factor change using $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen 2001), normalized to *ACTIN2* and relative to the lowest observed transcription (for day profiles) or relative to expression in the non-embryogenic line M9 (for distinct day comparisons). Computer analyses were performed using the GenEX software (MultiD Analyses AB, Sweden).

All the experiments were carried out in triplicate. Change in gene expression among induction phase days were analyzed using the GenEX software (MultiD Analyses AB, Sweden). The results are expressed as mean \pm SD. Statistical analyses were performed using the Student's t-test and ANOVA. Differences between the mean values were considered to be significant at $p < 0.01$ or $p < 0.05$.

Results

Identification of *LEC1*, *LIL*, and *PRC2* complex genes in *Medicago truncatula*

Prior to the expression analysis of *LEC1*, *LIL*, and *PRC2* genes it was necessary to carry out the sequence alignment analysis of proteins encoded by these genes.

Analyses of *LEC1* and *LIL* were performed using their amino acid sequences. The phylogenetic tree showed that

Table 1 Description of analyzed genes, their primer sequence and product size

Gene name	Gene short	<i>Arabidopsis thaliana</i> accession number	Accession number nucleotide/ protein	Genome identifier	Primer sequence	Length (bp)	Tm (°C)	Amplicon length (bp)
<i>Leafy Cotyledon1 - Like</i>	L1L	NP_199578.2	XM_013603204	Medtr4g133952	F: AGATGAACA CGAGGCAGC AAGT	22	60.97	100
			XP_013458658		R: GCAATTGGC ATAAAACGG TCTT	22	59.99	
<i>Leafy Cotyledon1</i>	LEC1	NP_173612.2	XM_003589718	Medtr1g039040	F: AGTGAAGGT GAACCTGCT TCTGT	23	59.12	101
			XP_003589766		R: GGCATTGAT GAAAACGAT GAAGA	23	60.93	
<i>Curly Leaf</i>	CLF	NP_179919.1	XM_003611648	Medtr5g016870	F: CGAAGAGTC AATGCCGAA GTC	21	59.63	101
			XP_003611696		R: AGCTGACCC AACAGTTCC TACAA	23	59.84	
<i>Swinger</i>	SWN	NP_567221.1	XM_003591348	Medtr1g086980	F: CAAGCATCA AGATTCCAC GTATG	23	59.29	101
			XP_003591396		R: ATTGATCGT CAGCCATTC TCTGA	23	60.74	
<i>Vernalization2</i>	VRN2	NP_974563.1	XM_003611313	Medtr5g013150	F: GCTTTAAGG GTTTGCGAT TTCA	22	60.21	101
			XP_003611361		R: CGTTCACTG CTTGGTAAT CATTG	23	59.33	
<i>Multicopy Suppressor of Ira1</i>	MSI1	NP_200631.1	XM_003608461	Medtr4g096880	F: TCTCATGCT CGCTCAAGT TCAA	22	60.82	101
			XP_003608509		R: AACCTCCA ACTTCAGGA CGAT	22	60.85	
<i>Fertilization Independent Endosperm</i>	FIE	NP_188710.1	XM_013610904	Medtr1g028310	F: CGCAGCCGA CATACTCA GAA	21	61.07	101
			XP_013466358		R: GCACATGCC TTGAAATGG AAAT	22	60.82	

LEC1 and L1L to separate into two distinct clades (Fig. 1). *Medicago truncatula* LEC1 formed one clade alongside other LEC1 from the family *Fabaceae*: *Cicer arietinum*, *Glycine max* and *Phaseolus vulgaris*. The *MtLEC1* showed a 69% identity with *AtLEC1*. *MtL1L*, too, formed one clade alongside other LEC1 from the family *Fabaceae*, but the *MtL1L* similarity to *AtL1L* was 61%.

On the phylogenetic tree SWN and CLF to grouped separately (Fig. 2a). Both SWN and CLF assembled together with the *Fabaceae* proteins examined. *MtSWN* was more

closely related to *AtSWN* than to *AtCLF*. *MtCLF* showed a 47% identity with *MtSWN*. *MtMSI1* was a close ortholog of *AtMSI1* (Fig. 2b). Both *MSI1* and *FIE* grouped into one clade with *MSI1* and *FIE* of *Cicer arietinum*, *Glycine max*, and *Phaseolus vulgaris*, respectively (Fig. 2b, c). The *MtFIE* amino acid sequence showed a 72% affinity to *AtFIE*.

The presence of a protein-specific domain indicates its membership in a particular group of PRC2 complex. The domain-searching analysis revealed that, among the




Fig. 1 Unrooted phylogenetic tree based on deduced amino acid sequences of *Medicago truncatula* LEC1 and L1L. At *Arabidopsis thaliana*, Bn *Brassica napus*, Ca *Cicer arietinum*, Gm *Glycine max*, Mt *Medicago truncatula*, Nt *Nicotiana tomentosiformis*, Tc *Theobroma cacao*, Pv *Phaseolus vulgaris*, Rc *Ricinus communis*, Sl *Solanum lycopersicum*

Medicago PRC2 proteins analyzed, only CLF and SWN (described also as EZA1) have the SET domain (Fig. 3). Both MS11 and FIE contain the WD40 domain, but differ in the number of WD-40 repeats and the presence of the RBBP4 domain in the MS11 protein.

Callus induction in Jemalong M9 and M9-10a lines and its ability to form the embryos

The callus development in leaf explants of two *Medicago truncatula* cv. Jemalong genotypes, the non-embryogenic (M9) and embryogenic (M9-10a), after 2, 7, 14 and 21 days on the SH induction medium and the somatic embryos production after 14 days on the MS differentiation medium are shown in Fig. 4. During the first two days, the explants became swollen, but no visible differences between the two lines were observed. After a one-week induction the explants curled up at the cut edges and produced callus tissue (Fig. 4). The weight of the callus obtained from the M9-10a explants was 1.5-fold higher than that of the M9 line (Fig. 5). The last two weeks were the period of a rapid callus growth. The M9 leaf explant calluses had compact and even structure with green-yellow colour, while the M9-10a explant callus texture was granular, very loose, and had a light-yellow colour (Fig. 4). After 14 days, the M9-10a line leaf explants produced nearly twice as

much callus as the M9 line did (Fig. 5). On day 21, all the M9-10a leaf explants were covered by the callus tissue, whereas the M9 leaf explants showed callus to have developed less expansively (Fig. 4). The weight of the callus obtained from the M9-10a explants was more than two-fold higher than that of the M9 line (Fig. 5). The 21 day-old explants with well-developed callus tissue were placed on hormone-free MS differentiation medium. After the subsequent two weeks, somatic embryos developed only on the M9-10a embryogenic explants.

LEC1 and L1L expression during somatic embryogenesis

The relative expression of *LEC1* and *L1L* genes was analyzed during the induction phase at distinct time-points in the non-embryogenic (M9) and embryogenic (M9-10a) lines. In primary explants, the expression level of *LEC1* (Fig. 6a.1) and *L1L* (Fig. 6b.1) was elevated in both lines, and their expression did not change during the entire period in M9. In contrast, expression of *LEC1* in the M9-10a tissues increased: during the last two weeks of induction the expression was 7- and 25-fold higher on days 14 and 21, respectively, compared to the lowest expression observed at day 2. A direct comparison of *LEC1* expression in M9 and M9-10a on day 7 revealed no significant difference between the two lines (Fig. 6a.2). However, the gene's expression in M9-10a was nine-fold higher than that observed in M9 on day 14. The expression profile revealed that transcripts of *L1L* gene was expressed between days 2 and 7 in M9-10a (Fig. 6b.1). The expression grew to reach even 129- and 307-fold increase on days 14 and 21, respectively, compared to the lowest expression in the primary leaf explants of M9-10a. A direct comparison of *L1L* expression in M9 and M9-10a on day 7 and 14 showed a 129- and 168-fold increase in M9-10a, respectively, compared to the expression in M9 normalized to 1 (Fig. 6b.2). Both genes are expressed in embryogenic calluses during the induction phase only; moreover, both are highly expressed during the last two weeks of the process.

Expression analysis of genes encoding PRC2 complex proteins


Since PRC2 is known to take part in transition from the vegetative to the generative phase in *Arabidopsis thaliana* ontogenesis, it was interesting to find out whether the proteins: CLF, SWN, VRN2, MS11 and FIE, belonging to this complex, are putative involved in acquisition of embryogenic competence during the SE induction phase in *M. truncatula*. The relative expression of genes encoding these proteins was analyzed during the induction phase at distinct time-points. The expression in the M9-10a

Fig. 2 Unrooted phylogenetic tree based on deduced amino acid sequences of *Medicago truncatula* CLF and SWN (a), MSI1 (b) and FIE (c). At *Arabidopsis thaliana*, Bn *Brassica napus*, Ca *Cicer arietinum*, Gm *Glycine max*, Mt *Medicago truncatula*, Nt *Nicotiana tomentosiformis*, Tc *Theobroma cacao*, Pv *Phaseolus vulgaris*, Rc *Ricinus communis*, Sl *Solanum lycopersicum*

primary explants (day 0) in all the relative gene profiling experiments was assumed to be 1. The *CLF* expression in M9 during induction was slightly up-regulated with regard to that on 0 day (Fig. 7a.1), while the *CLF* expression in M9-10a was highly up-regulated; it was three-fold higher (at day 7 and 14) than the expression in the primary explant. After one- and two-week long induction (days 7 and 14), the expression was 1.8- and 1.6-fold higher in M9-10a (Fig. 7a.2). The expression profiles for *SWN* in both lines (Fig. 7b.1) revealed a similar pattern with a distinct drop down on the second day of induction. Moreover, the expression level of *SWN* on days 7 and 14 remained the same in the two lines (Fig. 7b.2). The expression profiles of *VRN2* and *FIE* genes were very similar in the two lines analyzed (Fig. 7c.1, d.1). The expression during first week of induction increased four- and six-fold for *VRN2* and *FIE*, respectively. From 7 day their expression was on this same level in both lines (Fig. 7c.2, d.2). The expression profile of *MSI1* (Fig. 7e.1) in M9 remained on a similar level, whereas in M9-10a it was highly up-regulated, the expression being 11-, 9-, and 14-fold higher (for days 7, 14, and 21, respectively) than it was at the beginning of induction. Comparison of the *MSI1* expression in both lines on days 7 and 14 showed that in M9-10a was twice as high as that in M9 (Fig. 7e.2). It is worth pointing out that the expression of *VRN2*, *MSI1* and *FIE* denoted in the primary leaf explants was about three-fold higher in M9 than in M9-10a, which might be regarded as a crucial initial background sufficient to regulate the subsequent processes.

Discussion

The ability to produce embryos from somatic cells of plant explants, known as the somatic embryogenesis (SE), has been studied for a long time, but still remains incompletely understood. At present, only just residual data concerning the regulation of the process at the epigenetic level carried out by the Polycomb proteins are available, and all concern *Arabidopsis*. It is known that acquisition of embryogenic competency is related to transition from the somatic to the embryogenic state with activation of transcription factors among other from the LEC group which the prominent function is the specification of cotyledon identity (Altamura et al. 2016). Since the Polycomb Repressive Complex 2 (PRC2) proteins takes part in transition from the vegetative






Fig. 3 A schematic view of the domain location of Polycomb Repressive Complex 2 proteins of *Arabidopsis thaliana* (At) and *Medicago truncatula* (Mt)

Fig. 4 The leaf-to-callus transition on SH induction medium (21 days) and somatic embryo production on MS differentiation medium (14 days) during the somatic embryogenesis of *Medicago truncatula* non-embryogenic (M9) and embryogenic lines (M9-10a) (black dots on the timeline—sampling time points)





Fig. 5 The growth rate of callus on leaf explants of *Medicago truncatula* non-embryogenic (M9) and embryogenic (M9-10a) lines during the induction phase on SH medium

to the reproductive program in *Arabidopsis* (Mozgova et al. 2015), it was interesting to investigate both the expression of genes encoding these proteins and coding the transcription factors from the LEC group during the SE induction in *Medicago truncatula* cv. Jemalong.

Two lines: the non-embryogenic (M9) and embryogenic (M9-10a) were examined during a three-week long induction phase. The first week of induction ended with

the residual callus on cut edges of the embryogenic line explants; the changes visible during that period were slight only. Dedifferentiation and the formation of totipotent cells during the initial days is an effect of molecular mechanisms and activation of required transcription factors without any remarkable morphological changes (Rose et al. 2013). During the last two weeks of induction, a rapid growth of callus was observed primarily in the embryogenic line, but the difference in the callus structure between the two lines was not an obvious factor determining the embryogenic potential. It would be more appropriate to call this stage of the process the expression phase (according to Almeida et al. 2012) rather than the induction one. When calluses were transferred to a hormone-free medium, the somatic embryos started to develop on the embryogenic line explants only, which resulted in phenotypes strong enough to differentiate between the lines tested. To sum up, the induction phase in the embryogenic line involves two distinct stages: dedifferentiation and expression, but it is not possible to determine when the formation of totipotent cells starts or ends.

Several genes are known to be involved in SE induction in plants, the most interesting among them being the LEC-group genes. The sequence alignment analysis of LEC1 and L1L in *Medicago truncatula* confirmed that, like in other




Fig. 6 Relative expression patterns of *LEC1* and *LEC1-LIKE* genes in the leaf explants of *Medicago truncatula* non-embryogenic (M9) and embryogenic (M9-10a) lines during the induction phase on SH medium. Transcript levels for *LEC1* and *LEC1-LIKE* genes were estimated by Real-time PCR and normalized to that of *ACTIN2*. Bars show standard deviation. Dotted line indicate and white bars M9-10a line, solid line and grey bars-M9 line. **a-b.1** Line graphs show the dynamic expression of particular genes during the 21 day of induction phase and indicates the fold changes of its expression relative to the lowest value for which assumed value of 1. Statistical analyses,


two-way ANOVA with confidence interval 0.05, significance between groups indicated as * for $P \leq 0.05$, ** for $P \leq 0.01$, *** for $P \leq 0.001$ and **** for $P \leq 0.0001$. Bars indicate \pm SD. **a-b.2** Column graphs show the difference in the level of transcription at 7 and 14 day between M9 and M9-10a line expressed as fold changes in relation to the obtained value in M9 line taken as 1. Statistical analyses, two-tailed t-test with confidence interval 0.05. Asterisks represent significance level respectively: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$ and **** $P \leq 0.0001$. Bars indicate \pm SD

vascular plants, these transcription factors grouped into two separated clades on phylogenetic tree, despite their overlapping role (Cagliari et al. 2014). The expression of *LIL* was observed to be induced only in the embryogenic line during the first week of induction and continued to increase steeply for the next two weeks, while *LEC1* was only just expressed during this latter period. Moreover, the expression level of *LIL* was much higher than that of *LEC1*. As both transcription factors can be functionally substituted to some extent, it is not clear if the presence of both is necessary during SE. The expression profile indicates that activation of both genes overlapped with the expression phase, which might suggest that the genetic program differs from that of the dedifferentiation phase. According to Huang et al. (2015), the high expression of *LEC1* during *Arabidopsis* zygotic embryogenesis is caused by the release of the vegetative tissue from the suppressing environment,

which might suggest that during the first two weeks of *Medicago truncatula* SE induction a switch from the vegetative to embryonic program takes place. Furthermore, our results indicate that *LEC1* and *LIL* may be used as good gene markers for SE in *Medicago truncatula*.

The Polycomb Repressive Complex 2 belongs to epigenetic factors which, together with PRC1, play a key role in plant development by acting through the regulation of gene silencing. Two members of the PRC2 complex: CLF and SWN, belong to the SET-domain proteins involved in histone H3 lysine 27 trimethylation. They are active in meristems and dividing cells throughout the plant vegetative and reproductive development (Chanvivattana et al. 2004). The phylogenetic tree of *Arabidopsis thaliana* and *Vitis vinifera* showed that CLF and SWN belonged to two distinct clades, but were much more similar to each other than to another SET-domain protein, MEDEA (MEA) (Chanvivattana et al.

Fig. 7 Relative expression patterns genes of the *PRC2* complex in the leaf explants of *Medicago truncatula* non-embryogenic (M9) and embryogenic (M9-10a) lines during the induction phase on SH medium (a–e). Transcript levels for each gene were estimated by Real-time PCR and normalized to *ACTIN2*. Bars show standard deviation. Dotted line indicate and white bars M9-10a line, solid line and grey bars—M9 line. **a–b.1** Line graphs show the dynamic expression of particular genes during the 21 day of induction phase and indicates the fold changes of its expression relative to the lowest value for which assumed value of 1. Statistical analyses, two-way ANOVA with confidence interval 0.05, significance between groups indicated as * for $P \leq 0.05$, ** for $P \leq 0.01$, *** for $P \leq 0.001$ and **** for $P \leq 0.0001$. Bars indicate \pm SD. **a–b.2** Column graphs show the difference in the level of transcription at 7 and 14 day between M9 and M9-10a line expressed as fold changes in relation to the obtained value in M9 line taken as 1. Statistical analyses, two-tailed t-test with confidence interval 0.05. Asterisks represent significance level respectively: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$ and **** $P \leq 0.0001$. Bars indicate \pm SD



2004; Almada et al. 2011). The analysis of *M. truncatula* CLF and SWN showed, too, that these two proteins grouped separately. *MtMEA* was excluded from the sequence alignment analysis due to the lack of *AtMEA* close orthologous in any available *Medicago truncatula* data base. *MEA* was shown to occur in the family *Brassicaceae* only (Spillane et al. 2007). Recent research revealed that CLF and SWN might be essential for the callus tissue formation in plants. During the SE induction in *M. truncatula*, *CLF* expression increase significantly in consecutive days of the process in the embryogenic line (M9-10a). In the non-embryogenic line (M9), expression of this gene was stable in all the stages samples analyzed. A completely different pattern of gene expression was shown by *SWN*: its transcription dropped sharply between days 0 and 2, and remained at low levels until the end of the induction. This result suggests that the increase of the *CLF* expression overlapped with the dedifferentiation of somatic cells, callus growth and acquisition of the embryogenic potential. In our case, the expression of *SWN* decreased, while that of *CLF* increased during induction in the embryogenic line. These results are consistent with previous conclusions that CLF and SWN may play interchangeable roles in the PRC2 complex of *Arabidopsis* (Chanvivattana et al. 2004). Also Chanvivattana et al. (2004) found that the lack of expression of the *SWN* and *CLF* genes led to spontaneous emergence of a callus-like structure on whole seedlings. However, a different result was reported by He et al. (2012), where *Arabidopsis thaliana* double mutants *clf/swn* failed to develop the callus tissue from leaf explants, but it was observed on root explants. Those suggest that these genes are required for the leaf, but not the root and seedling, during callus formation. Probably, CLF is also an active component of PRC2 during the SE induction phase in *Medicago truncatula*.

The VRN2 protein is another component of the PRC2 complex and, together with other proteins, may form a VRN-PRC2 complex. It regulates the transition from the vegetative to the reproductive development (Bemer and Grossniklaus 2012). During the induction phase, the *VRN2* gene expression increased gradually with the strong callus tissue development in the embryogenic line (M9-10a) of *M. truncatula*, whereas the expression in the non-embryogenic line (M9) was stable and the weight of the callus was lower than that in the M9-10a line. Different results were obtained during the development of *Arabidopsis emf2/vrn2* mutants which the expression of these genes during the callus-like tissue formation was absent (Schubert et al. 2005).

The two other components of the PRC2 complex, i.e. MSII and FIE bind together with a high efficiency and can work as a complex (Köhler et al. 2003). The amino acid identity and the number of WD repeats observed in MSII and FIE confirmed close identity of the protein in *M. truncatula* and *A. thaliana*. The two lines of *M. truncatula*

tested: the embryogenic and the non-embryogenic differed in their *MSII* expression during the induction phase of somatic embryogenesis. An increased expression of the gene was observed in the embryogenic line's callus tissue, which results in embryo formation during the differentiation phase. On the other hand, the *MSII* transcript level was stable in the non-embryogenic line and embryos did not appear. A strong increase in this gene's expression during the first week of the induction phase in the embryogenic line may suggest its role in a switch from the vegetative to the embryonic development pattern. These results may suggest that *MSII* is required for explant somatic cells to acquire the embryogenic potential and for embryo development, as in zygotic embryogenesis. In our experiment expression of FIE was on this same level in M9 and M9-10a line. Also studies on *fie* mutants showed that, they have the ability to disorganized growth, to form callus-like structures, and to develop somatic embryos at a high frequency (Kinoshita et al. 2001; Bouyer et al. 2011).

Two components of PRC2 complex, CLF and SWN are known to insert H3K27me3 marks in histones. Analysis of mutants showed that the lack of *clf/swn* activity resulted in an up-regulation of *LEC1* and *LEC2* genes in *Arabidopsis thaliana* (Makarevich et al. 2006). Regulation of *LEC1* by epigenetic mechanisms was also proposed for *Coffea canephora* (Nic-Can et al. 2013) where the presence of H3K27me3 marks within the *LEC1* histone-associated region was a possible cause of reduced transcriptional activity of the gene during early stages of induction somatic embryogenesis. In these experiments, *LEC1* expression increased with progress of somatic embryogenesis induction, which was consistent with our results where *LEC1* and *LIL* transcripts started to appear between days 7 and 21. This might be the effect of histone demethylation carried out by Trithorax group proteins, which removed H3K27me3 marks inserted by PRC2 proteins (Pien and Grossniklaus 2007; Köhler and Hennig 2010). This hypothesis, however, needs to be experimentally tested. Another possible mechanism might be related to differences of *PRC2* genes expression in the primary explants. *VRN2*, *MSII* and *FIE* showed a lower expression in the embryogenic line leaf explants, which might have resulted in the ability to form embryogenic callus. The higher expression in the non-embryogenic line may play an inhibitory role in this process, so - because of the difference in the expression of these genes—primary explants may possess a different level of chromatin trimethylation and a changed ability to step in SE. Involvement of *FIE* and *MSII* genes in determining the H3K27me3 level has been confirmed in *A. thaliana* (Bouyer et al. 2011; Derkacheva et al. 2013).

Our results showed that in leaf explants of *Medicago truncatula* embryogenic line during the prime events of somatic embryogenesis (induction phase) a strong

up-regulation of *LEC1* and *LIL* genes with accompanied increased of *CLF* and *MSI1* expression.

Acknowledgements This work was partly supported by the National Scientific Centre (NCN) Grant No. NN303801340. We are indebted to dr Teresa Radziejewska for linguistic assistance.

Author contributions AO, RI and KŁ conducted experiments and carried out the statistical analysis. EK contributed to the discussion of result. AO and EK wrote the manuscript, RI participated in the writing part of the manuscript. All the authors read the final version of the paper.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Alemanno L, Devic M, Niemenak N, Sanier C, Guillemot J, Rio M, Verdeil J-L, Montoro P (2008) Characterization of *leafy cotyledon1-like* during embryogenesis in *Theobroma cacao* L. *Planta* 227:853–866
- Almada R, Cabrera N, Casaretto JA, Peña-Cortés H, Riz-Lara S, González Villanueva E (2011) Epigenetic repressor-like genes are differentially regulated during grapevine (*Vitis vinifera* L.) development. *Plant Cell Rep* 30:1959–1968
- Almeida AM, Parreira JR, Santos R, Duque AS, Francisco R, Tome DFA, Ricardo CP, Coehlo AV, Fevereiro P (2012) A proteomics study of the induction of somatic embryogenesis in *Medicago truncatula* using 2DE and MALDI-TOF/TOF. *Physiol Plant* 146:236–249
- Altamura MM, Della Rovere F, Fattorini L, D'Angeli S, Falasca G (2016) Recent advances on genetic and physiological bases of in vitro somatic embryo formation. *Methods Mol Biol*. doi:10.1007/978-1-4939-3061-6_3
- Bemer M, Grossniklaus U (2012) Dynamic regulation of Polycomb group activity during plant development. *Curr Opin Plant Biol* 15:523–529
- Boutillier K, Offringa R, Sharma VK, Kieft H, Ouellet T, Zhang L, Hattori J, Liu C-M, van Lammeren AAM, Miki BL, Custers JBM, van Lookeren Campagne MM (2002) Ectopic expression of BABY BOOM triggers a conversion from vegetative to embryonic growth. *Plant Cell* 14(8):1737–1749
- Bouveret R, Schönrock N, Grissem W, Hennig L (2006) Regulation of flowering time by *Arabidopsis* MSI1. *Development* 133:1693–1702
- Bouyer D, Roudier F, Heese M, Andersen ED, Gey D, Nowack MK, Goodrich J, Renou JP, Grini PE, Colot V, Schnittger A (2011) Polycomb Repressive Complex 2 controls the embryo-to-seedling phase transition. *PLOS Genet*. doi:10.1371/journal.pgen.1002014
- Butenko Y, Ohad N (2011) Polycomb-group mediated epigenetic mechanisms through plant evolution. *Biochim et Biophys Acta* 1809:395–406
- Cagliari A, Turchetto-Zolet AC, Korbes AP, dos Santos Maraschin F, Margis R, Margis-Pinheiro M (2014) New insights on the evolution of Leafy cotyledon1 (*LEC1*) type genes in vascular plants. *Genomics* 103:380–387
- Chanvivattana Y, Bishopp A, Schubert D, Stock C, Moon YH, Sung ZR, Goodrich J (2004) Interaction of Polycomb-group proteins controlling flowering in *Arabidopsis*. *Development* 131(21):5263–5276
- Derkacheva M, Steinbach Y, Wildhaber T, Mozgová I, Mahrez W, Nanni P, Bischof S, Grissem W, Hennig L (2013) *Arabidopsis* MSI1 connects LHP1 to PRC2 complexes. *EMBO J* 32:2073–2085
- Domoki M, Györgyey J, Bíró J, Pasternak TP, Zvara Á, Bottka S, Puskás LG, Dudits D, Fehér A (2006) Identification and characterization of genes associated with the induction of embryonic competence in leaf-protoplast-derived alfalfa cells. *Biochim et Biophys Acta* 1759:543–551
- Elhiti M, Stasolla C, Wang A (2013) Molecular regulation of plant somatic embryogenesis. *In Vitro Cell Dev-PI* 49:631–642
- Fambrini M, Durante C, Cionini G, Geri C, Giorgetti L, Michelotti V, Salvini M, Pugliesi C (2006) Characterization of *LEAFY COTYLEDON1-LIKE* gene in *Helianthus annuus* and its relationship with zygotic and somatic embryogenesis. *Dev Genes Evol* 216:253–264
- Gaj MD, Zhang S, Harada JJ, Lemaux PG (2005) Leafy cotyledon genes are essential for induction of somatic embryogenesis of *Arabidopsis*. *Planta* 222(6):977–988
- Gaspar T, Kevers C, Penel C, Greppin H, Reid DM, Thorpe TA (1996) Plant hormones and plant growth regulators in plant tissue culture. *In Vitro Cell Dev-PI* 32:272–289
- Gendall AR, Levy YY, Wilson A, Dean C (2001) The *VERNALIZATION 2* gene mediates the epigenetic regulation of vernalization in *Arabidopsis*. *Cell* 107:525–535
- Gleason EJ, Kramer EM (2013) Conserved roles for Polycomb Repressive Complex 2 in the regulation of lateral organ development in *Aquilegia x coerulea* “Origami”. *BMC Plant Biol* 13:185 doi:10.1186/1471-2229-13-185
- Guitton A-E, Page DR, Chambrier P, Lionnet C, Faure J-E, Grossniklaus U, Berger F (2004) Identification of new members of Fertilization Independent Seed Polycomb Group pathway involved in the control of seed development in *Arabidopsis thaliana*. *Development* 131:2971–2981
- Guo F, Liu C, Xia H, Bi Y, Zhao C, Zhao S, Hou L, Li F, Wang X (2013) Induced expression of *AtLEC1* and *AtLEC2* differentially promotes somatic embryogenesis in transgenic Tobacco plants. *PLOS One* 8:e71714
- Harding EW, Tang W, Nichols KW, Fernandez DE, Perry SE (2003) Expression and maintenance of embryonic potential is enhanced through constitutive expression of *AGAMOUS-Like 15*. *Plant Physiol* 133(2):653–663
- He C, Chen X, Huang H, Lin X (2012) Reprogramming of H3K27me3 is critical for acquisition of pluripotency from cultured *Arabidopsis* tissues. *PLOS Genet* 8:e1002911
- Hennig L, Derkacheva M (2009) Diversity of Polycomb group complexes in plants: same rules, different players? *Trends Genet* 25(9):414–423
- Hennig L, Taranto P, Walser M, Schönrock N, Grissem W (2003) *Arabidopsis* MSI1 is required for epigenetic maintenance of reproductive development. *Development* 130:2555–2565
- Huang M, Hu Y, Liu X, Li Y, Hou X (2015) *Arabidopsis* LEAFY COTYLEDON1 controls cell fate determination during post-embryonic development. *Front Plant Sci* 6:955
- Ikeuchi M, Iwase A, Rymen B, Harashima H, Shibata M, Ohnuma M, Breuer C, Morao AK, de Lucas M, De Veylder L, Goodrich J, Brady SM, Roudier F, Sugimoto K (2015) PRC2 represses dedifferentiation of mature somatic cells in *Arabidopsis*. *Nat Plants*. doi:10.1038/nplants.2015.89
- Jia H, McCarty DR, Suzuki M (2013) Distinct roles of LAFL network genes in promoting the embryonic seedling fate in the absence of VAL repression. *Plant Physiol* 163:1293–1305

- Jiang D, Wang Y, Wang Y, He Y (2008) Repression of *FLOWERING LOCUS C* and *FLOWERING LOCUS T* by the *Arabidopsis* polycomb Repressive complex 2 COMPONENTS. PLOS One. doi:10.1371/journal.pone.0003404
- Kakar K, Wandrey M, Czechowski T, Gaertner T, Scheible W-R, Stitt M, Torres-Jerez I, Xiao Y, Redman JC, Wu HC, Cheung F, Town CD, Udvardi MK (2008) A community resource for high-throughput quantitative RT-PCR analysis of transcription factor gene expression in *Medicago truncatula*. Plant Methods 4:1
- Karami O, Aghavaishi B, Pour AM (2009) Molecular aspects of somatic-to-embryogenic transition in plants. J Chem Biol 2:177–190
- Katz A, Oliva M, Mosquna A, Hakim O, Ohad N (2004) FIE and CURLY LEAF polycomb proteins interact in the regulation of homeobox gene expression during sporophyte development. Plant J 37:707–719
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Mentjies P, Drummond A (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28(12):1647–1649
- Kinoshita T, Harada JJ, Goldberg RB, Fischer RL (2001) Polycomb repression of flowering during early plant development. PNAS 98(24):14156–14161
- Köhler C, Hennig L (2010) Regulation of cell identity by plant Polycomb and trithorax group proteins. Curr Opin Genet Dev 20(5):541–547
- Köhler C, Makarevich G (2006) Epigenetic mechanisms governing seed development in plants. EMBO Rep 7(12):1223–1227
- Köhler C, Hennig L, Bouveret R, Gheyselincx J, Grossniklaus U, Gruissem W (2003) *Arabidopsis* MSII is a component of the MEA/FIE Polycomb group complex and required for seed development. EMBO J 22(18):4804–4814
- Kwong RW, Bui AQ, Lee H, Kwong LW, Fischer RL, Goldberg RB, Harada JJ (2003) LEAFY COTYLEDON1-LIKE defines a class of regulators essential for embryo development. Plant Cell 15:5–18
- Ledwoń A, Gaj MD (2011) *LEAFY COTYLEDON1*, *FUSCA3* expression and auxin treatment in relation to somatic embryogenesis induction in *Arabidopsis*. Plant Growth Regul 65:157–167
- Lee H, Fischer RL, Goldberg RB, Harada JJ (2003) *Arabidopsis* LEAFY COTYLEDON1 represents a functionally specialized subunit of the CCAAT binding transcription factor. PNAS 100:2152–2156
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. Methods 25:402–408
- Lotan T, Ohto M, Yee KM, West MAL, Lo R, Kwong RW, Yamagishi K, Fischer RL, Goldberg RB, Harada JJ (1998) *Arabidopsis* LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells. Cell 93:1195–1205
- Maillot P, Lebel S, Schellenbaum P, Jacques A, Walter B (2009) Differential regulation of *SERK*, *LECI-Like* and *Pathogenesis-Related* genes during indirect secondary somatic embryogenesis in grapevine. Plant Physiol Bioch 47:743–752
- Makarevich G, Leroy O, Akinci U, Schubert D, Clarenz O, Goodrich J, Grossniklaus U, Köhler C (2006) Different Polycomb group complexes regulate common target genes in *Arabidopsis*. EMBO Rep 7(9):947–952
- Mantiri FR, Kurdyukov S, Lohar DP, Sharopova N, Saeed NA, Wang X-D, VandenBosch KA, Rose RJ (2008) The transcription factor MtSERF1 of the ERF subfamily identified by transcriptional profiling is required for somatic embryogenesis induced by auxin plus cytokinin in *Medicago truncatula*. Plant Physiol 146(4):1622–1636
- Margueron R, Reinberg D (2011) The Polycomb complex PRC2 and its mark in life. Nature 469:343–349
- Meinke DW (1992) A homeotic mutant of *Arabidopsis thaliana* with leafy cotyledons. Science 258:1647–1650
- Mozgova I, Köhler C, Hennig L (2015) Keeping the gate closed: functions of the Polycomb repressive complex PRC2 in development. Plant J 83:121–132
- Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with Tobacco tissue cultures. Physiol Plant 15(3):473–497
- Neves LO, Duque SRL, Almeida JS, Feveireiro PS (1999) Repetitive somatic embryogenesis in *Medicago truncatula* ssp. *Narbonensis* and *M. truncatula* Gaertn cv. *Jemalong*. Plant Cell Rep 18:398–405
- Nic-Can GI, López-Torres A, Barredo-Poll F, Wrobel K, Loyola-Vargas VM, Rojas-Herrera R, De-la-Peña C (2013) New Insights into somatic embryogenesis: *LEAFY COTYLEDON1*, *BABY BOOM1* and *WUSCHEL-RELATED HOMEBOX4* are epigenetically regulated in *Coffea canephora*. PLOS One 8:e72160
- Nolan KE, Rose RJ, Gorst JR (1989) Regeneration of *Medicago truncatula* from tissue culture: increased somatic embryogenesis using explants from regenerated plants. Plant Cell Rep 8:278–281
- Peréz M, Cañal MJ, Toorop PE (2015) Expression analysis of epigenetic and abscisic acid-related genes during maturation of *Quercus suber* somatic embryos. Plant Cell Tiss Org Cult 121(2):353–366
- Pien S, Grossniklaus U (2007) Polycomb group and trithorax group proteins in *Arabidopsis*. Biochim Biophys Acta 1769(5–6):375–382
- Rose JR, Sheahan MB, Tiew TW-Y (2013) Connecting stress to development in the induction of somatic embryogenesis. In: Aslam J, Srivastava PS, Sharma MP (ed) Somatic embryogenesis and gene expression, 1st edn. Narosa Publishing House, New Delhi, pp 146–156
- Schellenbaum P, Jacques A, Maillot P, Bertsch C, Mazet F, Farine S, Walter B (2008) Characterization of *VvSERK1*, *VvSERK2*, *VvSERK3* and *VvLIL* genes and their expression during somatic embryogenesis of grapevine (*Vitis vinifera* L.). Plant Cell Rep 27:1799–1809
- Schenk RU, Hildebrandt AC (1972) Medium and techniques for induction and growth of monocotyledonous and dicotyledonous plant cell cultures. Can J Botany 50(1):199–204
- Schönrock N, Bouveret R, Leroy O, Borghi L, Köhler C, Gruissem W, Hennig L (2006) Polycomb-group proteins repress the floral activator *AGL19* in the *FLC*-independent vernalization pathway. Genes Dev 20(12):1667–1678
- Schubert D, Clarenz O, Goodrich J (2005) Epigenetic control of plant development by Polycomb-group proteins. Curr Opin Plant Biol 8:533–561
- Smertenko A, Bozhkov PV (2014) Somatic embryogenesis: life and death processes during apical-basal patterning. J Exp Bot 65:1343–1360
- Spillane C, MacDougall C, Stock C, Köhler C, Vielle-Calzada J-P, Nunes SM, Grossniklaus U, Goodrich J (2000) Interaction of the *Arabidopsis* Polycomb group proteins FIE and MEA mediates their common phenotypes. Curr Biol 10(23):1535–1538
- Spillane C, Schmid KJ, Laouéillé-Dupart S, Pien S, Escobar-Restrepo J-M, Baroux C, Gagliardini V, Page DR, Wolfe KH, Grossniklaus U (2007) Positive darwinian selection at the imprinted MEDEA locus in plants. Nature 448:349–352
- Steinbach Y, Hennig L (2014) *Arabidopsis* MSII functions in photoperiodic flowering time control. Front Plant Sci 5:77
- West MAL, Yee KM, Danao J, Zimmerman JL, Fischer RL, Goldberg RB, Harada JJ (1994) LEAFY COTYLEDON1 is an essential

- regulator of late embryogenesis and cotyledon identity in *Arabidopsis*. *Plant Cell* 6:1731–1745
- Willemsen V, Scheres B (2004) Mechanisms of pattern formation in plant embryogenesis. *Annu Rev Genet* 38:587–614
- Williams EG, Maheswaran G (1986) Somatic embryogenesis: factors influencing coordinated behavior of cells as an embryogenic group. *Ann Bot* 57:433–462
- Yadegari R, Kinoshita T, Lotan O, Cohen G, Katz A, Choi Y, Katz A, Nakashima K, Harada JJ, Goldberg RB, Fischer RL, Ohad N (2000) Mutations in the FIE and MEA genes that encode interacting Polycomb proteins cause parent-of-origin effects on seed development by distinct mechanisms. *Plant Cell* 12:2367–2381
- Yazawa K, Takahata K, Kamada H (2004) Isolation of the gene encoding Carrot leafy cotyledon1 and expression analysis during somatic and zygotic embryogenesis. *Plant Physiol Bioch* 42:215–223
- Yoshida N, Yanai Y, Chen L, Kato Y, Hiratsuka J, Miwa T, Sung ZR, Takahashi S (2001) EMBRYONIC FLOWER2, a novel Polycomb group protein homolog, mediates shoot development and flowering in *Arabidopsis*. *Plant Cell* 13:2471–2481
- Zhang S, Wong L, Meng L, Lemaux PG (2002) Similarity of expression patterns of *knotted1* and *ZmLEC1* during somatic and zygotic embryogenesis in maize (*Zea mays* L.). *Planta* 215:191–194
- Zheng Q, Zheng Y, Perry SE (2013) AGAMOUS-Like15 promotes somatic embryogenesis in *Arabidopsis* and Soybean in part by the control of ethylene biosynthesis and response. *Plant Physiol* 161:2113–2127
- Zimmerman JL (1993) Somatic embryogenesis: a model for early development in higher plants. *The Plant Cell* 5:1411–1423
- Zuo J, Niu Q-W, Frugis G, Chua N-H (2002) The *WUSCHEL* gene promotes vegetative-to-embryogenic transition in *Arabidopsis*. *Plant J* 30(3):349–335