



# Effect of photoperiod and plant growth regulators on in vitro mass bulblet proliferation of *Narcissus tazetta* L. (Amaryllidaceae), a potential source of galantamine

Majid Rahimi Khonakdari<sup>1</sup> · Hassan Rezadoost<sup>2</sup> · Roya Heydari<sup>1</sup> · Mohammad Hossein Mirjalili<sup>1</sup>

Received: 7 March 2020 / Accepted: 14 May 2020 / Published online: 20 May 2020  
© Springer Nature B.V. 2020

## Abstract

*Narcissus tazetta* L., a bulbous plant belongs to the Amaryllidaceae family, contains alkaloid galantamine (GAL) with acetylcholinesterase inhibitory activity which has been recently considered to treat Alzheimer's disease (AD). In the current work, the effect of photoperiod (16/8 h light/dark and 24 h dark) and various concentrations of NAA, BAP, and GA<sub>3</sub> (0, 0.5, 1 and 2 mg l<sup>-1</sup>) on the in vitro mass bulblet regeneration of *N. tazetta* was studied. The GAL production ability of the regenerated bulblets was assessed by HPLC-UV-MS. Light treatments significantly affected the number of bulblet and leaf, the ratio of bulblet/leaf, and leaf length. The maximum number of bulblet (31.0 ± 1.58) and leaf (13.3 ± 1.33) was recorded from the cultures fortified with NAA and BAP (2 mg l<sup>-1</sup>) kept in 16/8 h light/dark, while the maximum leaf length (2.1 ± 0.92 cm) was measured on the MS medium containing 0.5 mg l<sup>-1</sup> NAA and 2 mg l<sup>-1</sup> BAP incubated in the same photoperiod. The average ratio of bulblet proliferation per explant was significantly different between studied photoperiod (1.1 ± 0.86) and 24 h dark (0.62 ± 0.31). The regenerated bulblets contained 40 and 20 μg g<sup>-1</sup> DW GAL underexposed photoperiod and 24 h dark, respectively. This information could be useful in the commercial production of GAL as a valuable anti-AD compound through in vitro mass bulblet proliferation of *N. tazetta*.

## Key message

The regenerated mass bulblets of *Narcissus tazetta* (Amaryllidaceae) on MS medium containing 2 mg l<sup>-1</sup> NAA and BAP kept in 16/8 h light/dark are recommended to produce galantamine and lycorine.

**Keywords** Amaryllidaceae · *In vitro* culture · Propagation · Alkaloid · Galantamine · Alzheimer's disease

## Abbreviations

GAL	Galantamine	AAs	Amaryllidaceae alkaloids
AChE	Acetylcholinesterase	MS	Murashige and Skoog
AD	Alzheimer's disease	BAP	6-Benzylaminopurine
SMs	Secondary metabolites	NAA	1-Naphthalene-acetic acid
PCTOC	Plant cell, tissue, and organ culture	HPLC	High-performance liquid chromatography
PGRs	Plant growth regulators	LC-MS	Liquid chromatography-mass spectrometry
		CRD	Completely randomized design
		TDZ	Thidiazuron
		FW	Fresh weight
		DW	Dry weight
		TIBA	Triiodobenzoic acid
		CCC	Chlorocholine chloride

Communicated by Ali R. Alan.

✉ Mohammad Hossein Mirjalili  
m-mirjalili@sbu.ac.ir

<sup>1</sup> Department of Agriculture, Medicinal Plants and Drugs Research Institute, Shahid Beheshti University, G.C., Evin, 1983969411 Tehran, Iran

<sup>2</sup> Department of Phytochemistry, Medicinal Plants and Drugs Research Institute, Shahid Beheshti University, G.C., Evin, 1983969411 Tehran, Iran

## Introduction

The genus of *Narcissus* L. belonging to the subfamily Amaryllidoideae is one of the fifteen genera of the Amaryllidaceae family, which contains 80 to 100 wild species of perennial bulbiferous geophytes (Berkov et al. 2009; Hanks 2002). The genus is widely distributed in the southwest of Europe, with the Iberian Peninsula diversity center (Spain and Portugal) and North Africa (Ito et al. 1999). Few species of *Narcissus* have been developed in France and Italy, and some are found in the Balkans and the Eastern Mediterranean regions (Hanks 2002). The genus members are commonly known as daffodil, daffadow-dilly, narcissus, and jonquil throughout the world. *Narcissus tazetta* L. is an indigenous ornamental and fragrant plant species that is distributed from Portugal to Turkey. The plant is also growing from central to eastern Asia (Berkov et al. 2009; Hanks 2002). Indigenous habitats of *Narcissus* species are diverse and include a range of low-lying lands to mountainous areas as well as lawns, shrubs, forests, and cliffs (Hanks 2002). The genus *Narcissus* and the other genera of Amaryllidaceae family i.e., *Ungernia* Bunge, *Leucojum* L., *Galanthus* L., *Zephyranthes* Herb., *Hymenocallis* Salisb., and *Haemanthus* L. are well-known alkaloid-containing plants, due to the presence of Amaryllidaceae alkaloids (AAs) including lycorin, crinine, hemanthamine, narciclasine, galantamine, tazettine, homolycorine, montanine, and norbelladine which form an entirely different taxonomic and chemical group (Bastida et al. 2006; Torras-Claveria et al. 2013).

Galantamine (GAL) is a well-known long-acting, reversible AAs, and a competitive inhibitor for the acetylcholinesterase (AChE) enzyme and a nicotine receptor allosteric modifier for acetylcholine (Maelicke 2000; Woodruff-Pak et al. 2002). GAL can cross from the blood-brain barrier and effect on the central nervous system (Bastida et al. 2006; Ghosal et al. 1990). Due to the ability of GAL to enhance central cholinergic function, it has been introduced in Europe to treat Alzheimer's disease (AD) since 1980 (Bores and Kosley 1996). GAL-hydrobromide as a superior pharmaceutical profile has a higher tolerance to the major AChE inhibitors, including physostigmine or tacrine, and is currently used in the United States under the name Razadine<sup>®</sup>, known as Reminyl<sup>®</sup>, to treat certain stages of AD (Grutzendler and Morris 2001; Heinrich and Teoh 2004). Pharmaceutical demand of GAL has dramatically increased due to its significant effect on memory enhancement and AD control. The cost of GAL production is determined based on the quality of plant raw materials and their alkaloid content. The phytochemical characteristics of the plant materials were also affected by the extraction efficiency, which finally led to increase the

costs of GAL production. Bulbs of *Narcissus* species and the other members of the Amaryllidaceae family including *Leucojum aestivum* L., *Ungernia victoris* Vved. ex Artjush., and *Lycorice radiata* Herb. have been reported as GAL-rich natural sources (Abou-Donia et al. 2008; Berkov et al. 2009).

Different phytochemical patterns of plants due to the environmental changes, low content of their secondary metabolites (SMs), and a limitation for their materials collection from the natural habitats show the need to pay attention to the production of SMs by biotechnological methods (Ravishankar and Ramachandra Rao 2000). In the past decades, *in vitro* culture techniques have been used to produce massive amounts of many SMs in a short time, especially for types with valuable pharmaceutical effects (Malik et al. 2011). Biotechnology, through *in vitro* plant cell, tissue, and organ culture (PCTOC) enables the production of plant bioactive compounds under stable and controlled conditions in a shorter period. Mainly, PCTOC has been assessed in the study of plant developmental processes, physiological changes, production, and extraction of SMs. Therefore, these techniques can be simultaneously used for mass propagation, cloning, and the production of SMs from cultured organs or cells to provide the high demand for pharmaceutical industries (Bonfill et al. 2013; Malik et al. 2011; Sarmadi et al. 2019).

In a variety of plant species, especially monocotyledons, callus induction and establishment of cell culture are difficult, so the multiplication of their medicinal organs through *in vitro* cultures can be alternatively used to produce a remarkable amount of target bioactive compounds. Micropropagation of various bulbous plants has been so far reported (Haque and Ghosh 2016; Juan-Vicedo et al. 2019; Trujillo-Chacón et al. 2020; Ulrich et al. 1999; Zayed et al. 2011). *In vitro* bulb regeneration, callus induction, and somatic embryogenesis in several species and varieties of *Narcissus* have also been reported (Bergoñón et al. 1996; Chow et al. 1992; Colque et al. 2004; Abdel-Rahman et al. 2017; Malik and Bach 2017; Sage et al. 2000; Santos and Salema 2000; Santos et al. 1998; Sellés et al. 1997; Sochacki 2008; Staikidou et al. 2005).


Our literature survey revealed that *in vitro* propagation of *N. tazetta* and its varieties has already been reported (Chen et al. 2005; Chen and Ziv 2005; Zhang et al. 2013; Abdel-Rahman et al. 2017). To the best of our knowledge, the effect of plant growth regulators (PGRs), photoperiod and their interaction on the *in vitro* bulblet proliferation of the plant has not been studied yet. In the following of our recent work on the quantification of GAL in the Iranian populations of *N. tazetta* (Rahimi Khonakdari et al. 2018), the present study aimed to gain insight into the effect of photoperiod and various types and concentration

of PGRs on bulb multiplication and biomass production of the plant in culture condition *in vitro*. This information can be used in the semi-industrial production of the important anti-AD compound GAL and the other AAs through controlled *N. tazetta* bulblet culture.

## Materials and methods

### Plant material

Healthy bulbs were collected from *N. tazetta* cultivated in Matan Kola village, Aliabad Rural, Ghaemshahr,



**Fig. 1** In vitro bulblet proliferation of *Narcissus tazetta* L. Map of collection site of the plant material studied (A), The plant used as explants source (B), The fresh and healthy bulbs (C), washed and

sterile bulb used for explants excision (D), Sterile two-scale explants (E), established culture (F), proliferated bulblets on the MS medium containing NAA and BAP ( $2 \text{ mg l}^{-1}$ ) kept in 16/8 h light/dark (G)

**Table 1** Geographical and edaphological characteristics of collection site of *Narcissus tazetta* used in this study

Geographical characteristics	
Latitude (N)	36° 25' 01" N
Longitude (E)	52° 51' 13" E
Altitude (m)	106
Mean Annual Temperature (°C)	17.0
Mean Annual Precipitation(mm/year)	829.6
Edaphological characteristics	
N (total)%	0.15
Pava (ppm)	25.6
Kava (ppm)	405
pH	7.5
Electrical conductivity (ds/m)	0.72
Organic carbon (%)	1.83
Organic manure (%)	3.15
Sand (%)	35
Silt (%)	35
Clay (%)	30
Texture	Clay loam

Mazandaran Province in November 2018 and used as explants source (Fig. 1A–1C). Geographical and edaphological characteristics of the collection site are presented in Table 1.

## Sterilization and culture establishment

The plant bulbs were cleaned and washed carefully. Outer and dried scales were then removed and the rest bulbs (Fig. 1D) were washed under running tap water for 48–72 h. The bulbs were then incubated for 1 h in a bath of water at 49 °C to eliminate endophytes. The bulbs were then floated in sterile distilled water containing a few drops of Tween 80 for an hour. In the next step, the bulbs were immersed in 70% ethanol for 2 min and then washed with sterile distilled water for 2 min. The bulbs were then immersed in a 2% NaOCl solution containing a few drops of Twin 20 for 25 min with slow shaking. The bulbs were finally washed three times with sterile distilled water at intervals of 2, 5, and 7 min, and placed on filter paper. Decontaminated bulbs are divided into two-scale explants (Fig. 1E) which comprise of two adjacent scales connected by a piece of basal plate tissue (1 cm<sup>2</sup>) and were then placed on the MS medium containing 30 g l<sup>-1</sup> sugar and 8 g l<sup>-1</sup> agar supplemented with different concentrations of PGRs (Fig. 1F).

### *In vitro* bulb proliferation.

Sterile two-scale explants of *N. tazetta* were transferred to the baby food jars containing prepared MS media (40 ml)

**Table 2** Different treatments used in the present study

Treatment code	Plant growth regulator and concentration (mg l <sup>-1</sup> )
t0	GA <sub>0</sub> , NAA <sub>0</sub> , BA <sub>0</sub>
t1	NAA <sub>0.5</sub> , BA <sub>0.5</sub>
t2	NAA <sub>0.5</sub> , BA <sub>1</sub>
t3	NAA <sub>0.5</sub> , BA <sub>2</sub>
t4	NAA <sub>1</sub> , BA <sub>0.5</sub>
t5	NAA <sub>1</sub> , BA <sub>1</sub>
t6	NAA <sub>1</sub> , BA <sub>2</sub>
t7	NAA <sub>2</sub> , BA <sub>0.5</sub>
t8	NAA <sub>2</sub> , BA <sub>1</sub>
t9	NAA <sub>2</sub> , BA <sub>2</sub>
t10	GA <sub>0.5</sub> , NAA <sub>0.5</sub> , BA <sub>0.5</sub>
t11	GA <sub>0.5</sub> , NAA <sub>1</sub> , BA <sub>1</sub>
t12	GA <sub>0.5</sub> , NAA <sub>2</sub> , BA <sub>2</sub>
t13	GA <sub>1</sub> , NAA <sub>0.5</sub> , BA <sub>0.5</sub>
t14	GA <sub>1</sub> , NAA <sub>1</sub> , BA <sub>1</sub>
t15	GA <sub>1</sub> , NAA <sub>2</sub> , BA <sub>2</sub>
t16	GA <sub>2</sub> , NAA <sub>0.5</sub> , BA <sub>0.5</sub>
t17	GA <sub>2</sub> , NAA <sub>1</sub> , BA <sub>1</sub>
t18	GA <sub>2</sub> , NAA <sub>2</sub> , BA <sub>2</sub>

containing different concentrations (0, 0.5, 1, and 2 mg l<sup>-1</sup>) of BAP, NAA, and GA<sub>3</sub> (Table 2). MS medium without PGRs was also considered as a control. The induction of bulblet was also studied under two light regimes including 16/8 h light/dark provided by cool-white fluorescent lamps at a photon flux density of 40 μmol m<sup>-2</sup> s<sup>-1</sup> and 24 h dark kept in a paper box. All cultures were incubated in a growth chamber at 25 ± 2 °C. Each baby food jar was considered as an experimental unit, and the experiment was done in triplicates per treatment. In total, five jars were considered as experimental units in each replicate. Four two-scale explants were cultured in each jar. The number of bulblet and leaf, the ratio of bulblet/leaf, and leaf length were measured after four weeks (Fig. 1G).

## Extraction and HPLC-DAD-MS analysis

Four-week-old proliferated bulblets comprising photosynthetically active leaves (1–2 cm) were used for the extraction of alkaloids. Extraction, high-performance liquid chromatography (HPLC)-photo-diode array detector (DAD), and HPLC-mass spectrometry (MS) analyses have been conducted as described previously (Georgieva et al. 2007; Rahimi Khonakdari et al. 2018). For instance, 15 milliliters of methanol were added to 300 mg of completely powdered freeze-dried bulblets and then sonicated for 15 min at room

temperature. This stage was repeated thrice. The mixture was then centrifuged for 15 min at 4400 rpm and passed through the filter. The extracts were dried using a rotary evaporator (Heidolph Instruments GmbH, Schwabach Germany). The dried extract was re-dissolved in 4 ml H<sub>2</sub>SO<sub>4</sub> 3% and defatted with 5 ml diethyl ether (n=03). After basification to pH 9–10 with 25% ammonia, the GAL was extracted with chloroform (3 × 5 ml). The organic solvent was dried under reduced pressure in a rotary evaporator at 40 °C. The extracts were then dissolved in HPLC grade methanol (1 ml), filtered through a Millipore filter (0.45 mm) and used for analysis.

Galantamine quantification proceeded on a KNAUER liquid chromatography apparatus consisting of a 1000 Smartline Pump, a 5000 Smartline Manager Solvent Organizer and a 2800 Smartline Photo-DAD was used for the HPLC analysis. The injection was carried out through a 3900 Smartline Autosampler injector equipped with a 100 µl loop. The temperature of the column was controlled with a Jet Stream 2 Plus oven (Knauer, advanced scientific instrument, Berlin, Germany). The separation was achieved on a 25 cm × 4.6 mm with a pre-column, Eurospher 100-5 C<sub>18</sub> analytical column provided by Knauer (Berlin, Germany). Data acquisition and integration was performed with EZchrom Elite software. The elution was carried out in a gradient system with acetonitrile as the organic phase (solvent A) and 1% (w/v) ammonium acetate buffer adjusted to pH 6.6 with acetic acid (solvent B) with the flow-rate of 1 ml min<sup>-1</sup>. Peaks were monitored at 280 nm wavelength. Injection volume was 20 µl and the temperature was maintained at 25 °C. All injections were repeated three times (n=03). Calibration graphs were plotted subsequently for linear regression analysis of the peak area with concentration 5, 10, 25, 50, 80, 120, 150, and 200 mg l<sup>-1</sup>. The standard of GAL was purchased from Sigma (Sigma-Aldrich Corporation, MO, USA). Methanol, HPLC grade methanol, acetonitrile, and ammonium acetate of analytical grade were obtained from Merck (Darmstadt, Germany). HPLC grade water was used throughout the analysis.

HPLC-MS separations were carried out on an Agilent series 1200 system equipped with degasser, binary high

pressure mixing pump, column thermostat, and variable-wavelength UV-Vis absorbance detector (Agilent Technologies; Waldbronn, Germany). Using the same chromatographic condition, all eluted peaks were transferred to a Finnigan TM LCQ TM DECA instrument, comprising an ion trap. An ionization device was used for sample analyses (sheath gas: 80 ml/min; auxiliary gas: 20 ml min; spray voltage: 5 kV; capillary temperature: 300 °C; capillary voltage: 46 kV; tube lens: -60 kV). The Xcalibur 2.0 SR2 software (copyright Thermo Electron Corporation 1998–2006) was used.

### Statistical analysis

The experiment was a factorial design in a completely randomized design (CRD) with three replications. Data analysis was performed using SAS statistical software version 1.3.1. The means were compared using the Duncan multi-domain test at the probability level of 0.05, and according to their results, the charts were plotted using Excel software. The following equations calculate the ratio of bulb production and shoot formation indices:

Bulblet formation ratio index = number of explants/number of produced bulblets.

Shoot formation ratio index = number of explants/number of produced shoots.

### Results

In the present study, the used sterilization protocol as mentioned in the experimental comprising 70% ethanol, 2% NaOCl solution, and a few drops of Twin 20 showed satisfactory results on the control of surface contamination of the cultured two-scale explants. Pre-treatment of the explants with hot water before surface sterilization with sodium hypochlorite was additionally used. The explants incubation in hot water (49 °C) for an hour gave the best result for the disinfection of endophytes contamination of *N. tazetta* bulbs.

The results of the variance analysis revealed that hormonal treatments, light conditions, and their interaction had


**Table 3** Variance analysis of the effect of photoperiod and different concentration of plant growth regulators on in vitro regeneration of *Narcissus tazetta*

Variable source	DF	Mean of square				
		Number of bulblet	Ratio of bulblet	Number of shoot	Ratio of shoot	Shoot length
Photoperiod (A)	1	1093.06*	7.58*	805.34*	5.31*	10.50*
Plant growth regulator (B)	18	254.65*	1.76*	23.46*	0.17*	0.74*
A × B	18	130.305*	0.91*	20.38*	0.14*	0.44*
Error	74	1.21	0.01	0.59	0.005	0.02
Coefficient variability (CV)		10.53	10.76	21.34	23.07	23.70

\*Significant at 0.05 level

significant effects on the number of bulblet and leaf, the ratio of bulblet/leaf, and leaf length of *N. tazetta* (Table 3). The results showed that there was a significant difference ( $P < 0.05$ ) between treatments of photoperiod (16/8 h light/dark) and 24 h dark on the number of bulblet and leaf, and the ratio of each of them, as well as leaf length, so that the maximum number of bulblet ( $13.5 \pm 1.44$ ) and number of leaf ( $6.2 \pm 0.87$ ) were obtained under 16/8 h light/dark. It can be concluded that photoperiod is necessary for leaf production. The effect of light on the leaf length was also significant, so that, in light and dark conditions, the produced leaves were  $0.86 \pm 0.46$  and  $0.26 \pm 0.22$  cm, respectively. The mean ratio of bulblet production per sample was also significantly different between dark and photoperiod conditions. The mean value for light and dark conditions was  $1.1 \pm 0.86$  and  $0.62 \pm 0.31$  bulblet/explant, respectively.

In the study of the interaction effect of PGRs and light conditions, the highest number of bulblet ( $31.0 \pm 1.6$ ) and leaf ( $13.3 \pm 1.3$ ) were observed in the two-scale explants cultured on MS medium containing  $2 \text{ mg l}^{-1}$  of both BAP and NAA (treatment no. 9) under photoperiod condition (Fig. 2), but the highest leaf length ( $2.1 \pm 0.52$  cm) was observed on the treated explants with  $2 \text{ mg l}^{-1}$  BAP and  $0.5 \text{ mg l}^{-1}$  NAA (treatment no. 3) under light condition. The highest number of bulblet ( $28.6 \pm 1.2$ ) in continuous dark (24 h dark) was obtained from treatment no. 2 ( $0.5 \text{ mg l}^{-1}$  NAA and  $1 \text{ mg l}^{-1}$  BA). The number of the produced leaf under 24 h dark was much less than the exposed photoperiod so that the highest number of the leaf ( $2.6 \pm 0.92$ ) was measured in treatments no. 2, 6, and 7 under continuous dark (Fig. 2). No bulblet was produced on the hormone-free medium in both photoperiod (16/8 h light/dark) and 24 h dark conditions.



**Fig. 2** The effect of photoperiod and plant growth regulators on the average number of bulblet (**A**) and shoot (**B**) proliferated from two-scale explants of *Narcissus tazetta*. Different letters showed sig-


nificant differences among treatments at the level ( $P < 0.05$ ). For a detailed description of treatment code cf. Table 2

The average number of proliferated bulblet among all studied treatments in the light condition (16/8 h light/dark) was approximately twice than 24 h dark conditions. Among treatments containing GA<sub>3</sub>, treatment no. 13 (1 mg l<sup>-1</sup> GA<sub>3</sub>, 0.5 mg l<sup>-1</sup> BAP, and 0.5 mg l<sup>-1</sup> NAA) produced the highest number of the leaf (5.3 ± 0.86).

The average ratio of bulblet production per explant under 24 h dark and light conditions with different hormonal treatments is demonstrated in Fig. 3. In exposed photoperiod (16/8 h light/dark) and 24 h dark, the highest ratio of bulblet production was observed in treatments nos. 9 and 2, respectively. As can be seen, the highest percentage of leaf production ratio per explant (1.1) was in hormonal treatment no. 9 under 16/8 h light/dark, while the ratio in the treatments nos. 2, 6, and 7 under 24 h dark was up to 0.26 leaf per explant. The maximum mean of shoot length (2.1 ± 0.76 cm) was


obtained from treatment no. 3 underexposed photoperiod, while this treatment did not produce leaf in 24 h dark. The highest leaf length (0.83 ± 0.42 cm) in the dark condition was obtained from treatment no. 2. Treatment no. 9 showed a significant difference with other tested treatments and produced the highest leaf ratio in each explant. Among studied treatments containing GA<sub>3</sub>, the maximum leaf length (0.86 ± 0.38 cm) was obtained in treatment no. 10 (Fig. 3).

HPLC-UV-MS profiling of the methanolic extract of the regenerated bulblets of *N. tazetta* is shown in Fig. 4. Characteristically, the standard GAL (C<sub>17</sub>H<sub>21</sub>NO<sub>3</sub>) yielded an [M+H]<sup>+</sup> peak at a molecular weight of 288.6 Da (the corresponding exact mass for GA is 288.1594 Da). The peaks corresponding to GAL exhibited mass-to-charge (m/z) ratios corresponding to the molecular ions [M+H]<sup>+</sup> of authentic GAL (at 9.77 min), confirming the presence of GAL in the



**Fig. 3** The effect of photoperiod and plant growth regulators on the ratio of bulblet (A) and shoot (B) proliferated from two-scale explants of *Narcissus tazetta*. Different letters showed significant differences

among treatments at the level ( $P < 0.05$ ). For a detailed description of treatment code cf. Table 2



**Fig. 4** HPLC-UV-MS profiling of the methanolic extract of the regenerated bulblets of *Narcissus tazetta* L. HPLC-UV chromatogram recorded at 288 nm (**A**), corresponding total ion current chromatogram (TIC) (**B**), mass spectrum ( $[M+H]^+$ ) related to the of lycorine (**C**), and mass spectrum ( $[M+H]^+$ ) related to galantamine (**D**)

methanolic extract of the proliferated bulblets. Extracted ion chromatogram for 288.1594 Da (the exact mass related to GAL) led to find an isobaric compound of GAL. According to the LC-MS data, the presence of lycorine ( $C_{16}H_{17}NO_4$ ) in the methanolic extract of the studied *in vitro* proliferated bulblets was also confirmed. The exact mass for this compound and its corresponding retention time are 288.1230 Da and 5.6 min, respectively (Fig. 4). Quantitative HPLC determination showed that the regenerated bulblets on the best proliferation treatment underexposed photoperiod and 24 h dark produced 40 and 20  $\mu\text{g g}^{-1}$  DW GAL, respectively, while GAL has not been detected in the produced leaf.

## Discussion

The greatest problem in the *in vitro* culture techniques such as micropropagation, somatic embryogenesis, embryo rescue, somatic hybridization, somaclonal variation, and

PCTOC for the production of SMs is the contamination. A wide range of microorganisms, including fungi, yeasts, bacteria, and viruses are the main contributing factors in the contamination of *in vitro* cultures mentioned above (Langens-Gerrits and De Klerk 1999). Therefore, sterilization of the plant tissues along with other materials used in the *in vitro* techniques is one of the most basic and the most important step in the establishment of an *in vitro* culture.

The main challenge in this regard is the disinfection of explants because the intrinsic contaminants of plant tissues (endophytes) cannot be eliminated using common surface disinfectants and methods. Although this problem cannot be observed in meristem culture, the use of an effective method for the disinfection of the other materials, especially underground explants including bulbs, corms, tubers, and rhizomes is very necessary. Similar to the other bulbiferous plants such as *Lilium* and *Tulips*, bulbs of *Narcissus* contain a wide range of endophytes which can be contaminated the plant *in vitro* cultures.



In this study, the results of the disinfection and establishment of explants showed that sodium hypochlorite even at high concentrations and longtime did not affect the elimination of endophytes contamination. Although the use of mercuric chloride (0.1%) has been reported for the explant sterilization of some bulbous plants as *Zephyranthes grandiflora* (Gangopadhyay et al. 2010), and *Lilium polyphyllum* D. Don ex Royle (Taha et al. 2018), but it has not been considered in the present study due to the toxicity and health risk of its using. Abdel-Rahman et al. (2017) observed the best disinfection results for the bulbs of *N. tazetta* “Chinensis” through soaking the bulbs in 70% ethanol for 10 min followed by soaking bulbs in 2%  $\text{HgCl}_2$  for 10 min. The bulb disinfection of *Leucojum* with 70% ethanol (1 min) and domestos bleach (5% chlorine) 50% and 30% at intervals of 30 and 10 min has been reported (Stanilova et al. 2010). Treatments of the explants with hot water (40–50 °C for 1 to 4 h) followed by drying before sterilizing them with hypochlorite solutions have also been suggested (Teixeira da Silva et al. 2016). The use of hot water at 40 and 54 °C has been reported to eliminate the internal contamination of *Lilium* and *Narcissus* bulbous explants, respectively (Langens-Gerrits et al. 1998; Lagnes-Gerrits and De Klerk 1999).

*In vitro* organogenesis can be affected by several factors such as plant genotype, storage conditions, explant type, polar orientation of the explants, photoperiod, temperature, composition of culture medium, and PGRs (Chen and Ziv 2005; Abdel-Rahman et al. 2017; Begum et al. 2019; Cheesman et al. 2010; García-Fortea et al. 2020; Nasircilar et al. 2011; Sairam et al. 2003). Photoperiod and concentration of PGRs are probably the most important influencing factors on *in vitro* organogenesis in the bulbous plants. *In vitro* propagation of Amaryllidaceae members through scale explants comprising a piece of basal plate has also been reported (Ault 1995; Diop et al. 2006; Fennell et al. 2004; Rice et al. 2011). It has been claimed that adventitious shoots were successfully proliferated from tissue at the base of bulb scales in many bulbous species (Ozel et al. 2015; Rice et al. 2011). So, it is necessary to include the basal plate as part of the scale explant for further bulblet proliferation. Ozel et al. (2007) reported that the combination of BAP-NAA has a significant effect on the proliferation of *Muscari* bulbs from two-scale explants, which is consistent with the results obtained in this study (Ozel et al. 2007). An efficient *in vitro* regeneration of *N. tazetta* “Chinensis” through twin-scale explants with basal plate cultured on MS medium supplemented with 1 mg  $\text{l}^{-1}$  IBA + 1 mg  $\text{l}^{-1}$  triiodobenzoic acid (TIBA) and activated charcoal has been reported (Abdel-Rahman et al. 2017). The highest proliferation bulbs of *Muscari mirum* Speta has also been reported from four-scale explants cultured on MS medium containing 4 mg  $\text{l}^{-1}$  BAP and 0.25 mg  $\text{l}^{-1}$  NAA (Nasircilar et al. 2011). Rice et al. (2011) reported the highest bulblet induction from twin-scale explants of *Brunsvigia*

*undulata* F.M. Leight cultured on hormone-free medium. In some reports, both increasing and decreasing effects of TDZ concentration on bulblet induction of bulbous plants have been reported (Huetteman and Preece 1993; Thomas 2007), which has an increasing effect when used in combination with NAA. So, the synergistic effect of NAA in combination with cytokinins has been claimed in the bulblet formation of bulbous plants.

The significant effects of previous bulb storage on the performance of twin-scales for *in vitro* regeneration of *N. tazetta* have been reported by Chen and Ziv (2005). They revealed that the twin-scale explants isolated from the bulbs after three months in storage at 30 °C and a six week cold treatment at 15 °C produced high numbers of uniform adventitious buds. In their results, the elongated buds regenerated from inflorescence stem discs had a wide range of length variation when compared to adventitious shoots regenerated from twin-scale explants. They have also been studied the effect of polar orientation on regeneration of the inflorescence stem of *N. tazetta* which may be due to the presence of growth substances, especially the ratio of cytokinin and auxin within the explants. It has been claimed that the interactions between growth substances and the polar orientation of the explants, and their effects on the regeneration can be a complex phenomenon (Chen and Ziv 2005).

The bulblet induction on the PGRs-free medium has also been reported in many other Amaryllidaceae members including *Amaryllis belladonna* L. (De Bruyn et al. 1992), *Pancreatum maritimum* L. (Dragassaki et al. 2001), *Crinum macowanii* Baker. (Slabbert et al. 1993), and *B. undulata* (Rice et al. 2011). In this case, it can be concluded that the explant contains a high level of endogenous PGRs to induce bulblet formation. Although the effect of the genotype and the size of the explant on the bulb regeneration is confirmed, but the effect of NAA-cytokinin composition on the efficient production of bulbs in some other bulbous plants such as *Iris sanguinea* (Wang et al. 2018), *Lachenalia montana* (Aremu et al. 2015), *Fritillaria ruthenica* Wikstr. (Muraseva and Novikova 2018), *Muscari muscarimi* (Ozel et al. 2015), and *L. aestivum* (Ptak et al. 2013) have also been reported. The increasing effect of paclobutrazol and sucrose levels on the bulblet induction of *L. aestivum* in culture condition *in vitro* has been reported (Ptak et al. 2013). Sultana et al. (2010) studied the effect of BAP and chlorocholine chloride (CCC) concentrations in addition different levels of sucrose on the bulb proliferation of *Hippeastrum hybridum* and showed that the maximum bulb production was obtained on the medium culture supplemented with 6 mg  $\text{l}^{-1}$  BAP, 500 mg  $\text{l}^{-1}$  CCC, and 90 g  $\text{l}^{-1}$  sucrose (Sutlana et al. 2010). The increasing effect of polyamines and methyl jasmonate on the bulb formation of *Tulipa* has been reported by Podwyszynska et al. (2015).

The effect of photoperiod on the regeneration of bulbs in *A. belladonna* (De Bruyn et al. 1992), *Lilium longiflorum* (Niimi and Onozawa 1979), *B. undulate* (Rice et al. 2011), and *Eucomis zambesiaca* (Cheesman et al. 2010) has also been reported. The growth and developmental process of plants in *in vitro* cultures are significantly affected by the quality of light (color and wavelength), intensity, and photoperiod. Light is also often a key factor in the accumulation of SMs (Yang et al. 2018). The effect of different photoperiods on the regeneration rate of Paulownia under *in vitro* culture conditions has been reported (Yang et al. 2013). They approved that different light conditions can be affected by the leaf and root induction of the plant. In the contrary, Houllou et al. (2015) showed that photoperiod did not effect on shoot and root development of *Capraria biflora* L. while affected on the plant callus induction. The effect of photoperiod (16/8 h light/dark) on the bulb production from two-scale explants of *B. undulata* in *in vitro* condition has been reported (Rice et al. 2011). Cheesman et al. (2010) showed that 24 h dark completely inhibited bulblet formation in *Eucomis zambesiaca*. They have been found that the best bulblet induction occurred in the 8 h light cycle. Conversely, the highest bulblet regeneration in *Hyacinthus orientalis* (Kim et al. 1981) and *L. longiflorum* (Kumar et al. 2006) has been reported in continuous darkness which is in the opposite of our obtained results. The difference in results observed for *N. tazetta* could be due to the plant genotype and the type of explant used. Further studies on light conditions (quality and intensity) on the plant need to be investigated to optimize the real effect of light.

It has been revealed that environmental parameters such as light and day length interact with PGRs to cause developmental responses in plants (Cheesman et al. 2010; Kumar et al. 2006; Rice et al. 2011) which are in agreement with obtained results in the present study. Notably, the bulblet production decreased in the treatments containing GA<sub>3</sub> (No. 10–18). Thus, it can be concluded that this plant hormone has a negative effect on bulb production in *N. tazetta*. This is in agreement with obtained results by Pierik and Steegmans (1975) in the bulb proliferation of hyacinth and *E. zambesiaca* (Pierik and Steegmans 1975).

Generally, it can be concluded that low concentration of auxin and cytokinin is more suitable and efficient for bulb production, while increasing concentrations of these hormones, especially cytokinins, increased the production of the leaf. Santos et al. (2002) reported similar results in *in vitro* culture of *N. bulbocodium* L. using MS medium with two different hormonal compositions (BAP 4 mg l<sup>-1</sup> + NAA 12 mg l<sup>-1</sup> and BAP 2 mg l<sup>-1</sup> + IBA 1 mg l<sup>-1</sup>) which resulted in leaf induction on explants. In the second hormonal composition, a few small bulblets appeared after 70 days. Growth of proliferated bulblet was carried out with two subcultures on the same culture medium

with an increase of 9% sucrose and with or without NAA. They revealed that NAA has not a significant effect on the bulblet size, but led to a better formation of the root. In *in vitro* propagation of Narcissus, the effect of BAP and NAA on the leaf induction has also been reported by Hussey (1982).

Bergonon et al. (1996) showed that GAL production was stimulated by prolonged photoperiod in the shoots culture of *N. confuses* which has been reached to 2.5 mg g<sup>-1</sup> DW in the plant tissue culture Berkov et al. (2009) also claimed that light is an important factor in the biosynthesis of GAL in the shoots of *L. aestivum* cultured *in vitro*. They found that GAL production in the light conditions is nearly twice that of dark conditions, which is in agreement with obtained results in the present study. The concentration of GAL in different plant organs of *L. aestivum* has been reported between 0.02 and 0.60 mg g<sup>-1</sup> DW. As the other monocotyledons, callus induction and cell suspension culture establishment in Narcissus is relatively difficult. So, optimization of mass bulb proliferation of the plant can be considered as an efficient and alternative method for the production of medicinally important compound GAL. So far, several attempts for the production of GAL through *in vitro* cultures of the Amaryllidaceae family have been done. The effect of elicitation on the production of GAL in the shoot culture of *N. confuses* has been reported (Colque et al. 2004). Quantification of GAL in bulbs and *in vitro* cultures of *N. papyraceus* and four varieties of *N. tazetta* grown in Iran using GC-MS analysis has been recently reported (Tarakemeh et al. 2019). It has been shown that the content of GAL ranged from 6 to 105 µg g<sup>-1</sup> DW among the studied species and varieties. They also showed that GAL accumulation was significantly stimulated by 4'-O-methylnorbelladine precursor in *in vitro* culture samples of *N. tazetta* which has been reached to 1.7 and 82 µg g<sup>-1</sup> DW in the calli and bulblet cultures, respectively. Production of GAL through *in vitro* tissue culture of *N. pseudonarcissus* cv. Carlton has been currently reported by Ferdausi et al. (2020). According to the obtained LC-MS results comparing with the previous report (Karakoyun and Unver-Somer 2019), lycorine, an Amaryllidaceae alkaloid, was also found in the proliferated *N. tazetta* bulblets. Interestingly, lycorine is a well-known natural alkaloid with a wide range of pharmaceutical properties (Roy et al. 2018). Antiviral activity of lycorine against severe acute respiratory syndrome-associated coronavirus (SARS-CoV), measles virus, coxsackie virus, poliovirus, human immunodeficiency virus (HIV-1), and herpes simplex virus type 1 has been reported (Szlávik et al. 2004; Li et al. 2005). Therefore, further studies including optimization of medium culture conditions, elicitation, precursor feeding, and the other enhancement strategies can be recommended to produce this valuable alkaloid through developed *in vitro* mass bulblets culture of *N. tazetta*.

## Conclusions

The chemical synthesis of GAL has been successfully reported, but the main natural source for its commercial production is the plant materials of Amaryllidaceae members. The biomass production rate in these plants is very slow and time-consuming, so *in vitro* culture establishment of the plants like *N. tazetta* can be a good way to achieve maximum production of this medicinally important compound for further pharmaceutical purposes and commercial exploitation. Since GAL accumulation in the bulbs of *Narcissus* is higher than in the shoots, therefore, the present study focused on *in vitro* mass bulb production of the plant. This study showed that two-scale bulb explants of *N. tazetta* cultured on the MS medium supplemented with 2 mg l<sup>-1</sup> of both BAP and NAA under photoperiod condition produced the highest number of bulblet. Media culture containing GA<sub>3</sub> decreased bulblet proliferation in *N. tazetta*. It is worthy to note that more bulblets obtained from the medium supplemented with lower hormonal concentrations, which can be considered to reduce production costs in the mass bulblet culture of *N. tazetta* in the bioreactor for the commercial production of anti-AD compound GAL and medicinally important antiviral component lycorine as well.

**Acknowledgements** The authors thank the Shahid Beheshti University Research Council and the Iran National Science Foundation (INSF, Grant No. 94021603) for financial support of this project. We also thank Fatemeh Goudarzi for her kind collaboration in HPLC analysis.

**Author contributions** MRK contributed to the conception of the study, *in vitro* cultures establishment, data extraction statistical analysis, and writing of the manuscript. HR helped in HPLC-MS analysis. RH contributed in data collection, statistical analysis and revising the manuscript. MHM supervised the whole experiments and wrote the manuscript. All authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

## References

- Abdel-Rahman H, Al-Ansary AMF, Rashed KN, Rizkalla AA (2017) Micropropagation of *Narcissus tazetta* ‘Chinensis’ and its relation to secondary metabolites content. *J App Life Sci Int* 14(1):1–11
- Abou-Donia AH, Toaima SM, Hammada HM, Shawky E (2008) New rapid validated HPTLC method for the determination of galanthamine in Amaryllidaceae plant extracts. *Phytochem Anal* 19(4):353–358
- Aremu A, Dolezal K, Van Staden J (2015) New cytokinin-like compounds as a tool to improve rooting and establishment of micropropagated plantlets. In: VI international symposium on production and establishment of micropropagated plants, vol 1155, pp 497–504
- Ault JR (1995) *In vitro* propagation of *Eucomis autumnalis*, *E. comosa*, and *E. zambesiaca* by twin-scaling. *HortScience* 30(7):1441–1442
- Bastida J, Lavilla R, Viladomat F (2006) Chemical and biological aspects of *Narcissus* alkaloids. In: Cordel GA (ed) *The alkaloids: chemistry and biology*. Elsevier Inc, Amsterdam, pp 87–179
- Begum N, Zenat EA, Sarkar MK, Roy K, Munshi C, Jahan JL, Miskat A (2019) Micro propagation of soybean (*Glycine max* L.) BARI-5 variety. *Open Microbiol J* 13(1):177–187
- Bergoñón S, Codina C, Bastida J, Viladomat F, Melé E (1996) Galanthamine production in “shoot-clump” cultures of *Narcissus confusus* in liquid-shake medium. *Plant Cell Tiss Org Cult* 45(3):191–199
- Berkov S, Georgieva L, Kondakova V, Atanassov A, Viladomat F, Bastida J, Codina C (2009) Plant sources of galanthamine: phytochemical and biotechnological aspects. *Biotech Biotechnol Equip* 23(2):1170–1176
- Bonfill M, Malik S, Mirjalili MH, Goleniowski M, Cusido R, Palazon J (2013) Production and genetic engineering of terpenoids production in plant cell and organ cultures. In: Merillon KG (ed) *Natural products*. Ramawat, Berlin, pp 2761–2796
- Bores G, Kosley R (1996) Galanthamine derivatives for the treatment of Alzheimer’s disease. *Drugs Future* 21(6):621–635
- Cheesman L, Finnie J, Van Staden J (2010) *Eucomis zambesiaca* Baker: factors affecting *in vitro* bulblet induction. *S Afr J Bot* 76(3):543–549
- Chen J, Ziv M (2005) The effects of storage condition on starch metabolism and regeneration potentials of twin-scales and inflorescence stem explants of *Narcissus tazetta*. *In Vitro Cell Dev Biol—Plant* 41:816–821
- Chen L, Zhu X, Gu L, Wu J (2005) Efficient callus induction and plant regeneration from anther of Chinese narcissus (*Narcissus tazetta* L. var. *chinensis* Roem). *Plant Cell Rep* 24(7):401–407
- Chow Y, Selby C, Harvey B (1992) A simple method for maintaining high multiplication of *Narcissus* shoot cultures *in vitro*. *Plant Cell Tiss Org Cult* 30(3):227–230
- Colque R, Viladomat F, Bastida J, Codina C (2004) Improved production of galanthamine and related alkaloids by methyl jasmonate in *Narcissus confusus* shoot-clumps. *Planta Med* 70(12):1180–1188
- De Bruyn M, Ferreira D, Slabbert M, Pretorius J (1992) *In vitro* propagation of *Amaryllis belladonna*. *Plant Cell Tiss Org Cult* 31(3):179–184
- Diop MF, Ptak A, Chrétien F, Henry M, Chapleur Y, Laurain-Mattar D (2006) Galanthamine content of bulbs and *in vitro* cultures of *Leucojum aestivum* L. *Nat Prod Commun* 1(6):475–479
- Dragassaki M, Economou A, Vlahos J (2001) Bulblet formation *in vitro* and plantlet survival extra vitrum in *Pancreatum maritimum* L. In: I International symposium on acclimatization and establishment of micropropagated plants, vol 616, pp 347–352
- Fennell C, Van Staden J, Bornman C (2004) Biotechnology of southern African bulbs. *S Afr J Bot* 70(1):37–46
- Ferdausi A, Chang X, Hall A, Jones M (2020) Galanthamine production in tissue culture and metabolomic study on Amaryllidaceae alkaloids in *Narcissus pseudonarcissus* cv. *Carlton* Ind Crops Prod 144:112058
- Gangopadhyay M, Chakraborty D, Dewanjee S, Bhattacharya S (2010) Clonal propagation of *Zephyranthes grandiflora* using bulbs as explants. *Biol Plant* 54(4):793–797
- García-Forteza E, Lluch-Ruiz A, Pineda-Chaza BJ, García-Pérez A, Bracho-Gil JP, Plazas M, Gramazio P, Vilanova S, Moreno V, Prohens J (2020) A highly efficient organogenesis protocol based on zeatin riboside for *in vitro* regeneration of eggplant. *BMC Plant Biol* 20(1):6

- Georgieva L, Berkov S, Kondakova V, Bastida J, Viladomat F, Atanassov A, Codina C (2007) Alkaloid variability in *Leucojum aestivum* from wild populations. *Z Naturforsch C* 62:627–635
- Ghosal S, Singh SK, Unnikrishnan SG (1990) Effects of stress on alkaloid metabolism in *Crinum asiaticum*. *Phytochemistry* 29(3):805–811
- Grutzendler J, Morris JC (2001) Cholinesterase inhibitors for Alzheimer's disease. *Drugs* 61(1):41–52
- Hanks GR (2002) Narcissus and daffodil: the genus Narcissus. CRC Press, Boca Raton
- Haque SM, Ghosh B (2016) High-frequency somatic embryogenesis and artificial seeds for mass production of true-to-type plants in *Ledebouria revoluta*: an important cardioprotective plant. *Plant Cell Tiss Org Cult* 127(1):71–83
- Heinrich M, Teoh HL (2004) Galanthamine from snowdrop—the development of a modern drug against Alzheimer's disease from local Caucasian knowledge. *J Ethnopharmacol* 92:147–162
- Houllou LM, de Santana ERB, de Andrade Lima CS, de Souza RA, de Castro Torres GR, Alves GD, Yara R (2015) Effect of light, explant maturity and antioxidant on *in vitro* organogenesis and plant regeneration of the medicinal plant *Capraria biflora* L. *Res Biotechnol* 6(3):24–30
- Huetteman CA, Preece JE (1993) Thidiazuron: a potent cytokinin for woody plant tissue culture. *Plant Cell Tiss Org Cult* 33(2):105–119
- Hussey G (1982) *In vitro* propagation of Narcissus. *Ann Bot* 49(5):707–719
- Ito M, Kawamoto A, Kita Y, Yukawa T, Kurita S (1999) Phylogenetic relationships of Amaryllidaceae based on matK sequence data. *J Plant Res* 112(2):207–216
- Juan-Vicedo J, Pavlov A, Ríos S, Casas JL (2019) *In vitro* culture and micropropagation of the Baetic-Moroccan endemic plant *Lapiedra martinezii* Lag. (Amaryllidaceae). *In Vitro Cell Dev Biol Plant* 55(6):725–732
- Karakoyun C, Unver-Somer N (2019) Simultaneous quantitative analysis of biologically important Amaryllidaceae alkaloids in *Narcissus tazetta* L. subsp. *tazetta* by HPLC/PDA. *J Res Pharm* 23:498–505
- Kim Y, Hasegawa P, Bressan R (1981) *In vitro* propagation of hyacinth [*Hyacinthus orientalis*]. *HortScience* 16:645–647
- Kumar S, Kanwar J, Sharma D (2006) *In vitro* propagation of *Lilium*. *Adv Hortic Sci* 20(2):181–188
- Langens-Gerrits M, Alberts M, De Klerk GJ (1998) Hot-water treatment before tissue culture reduces initial contamination in *Lilium* and *Acer*. *Plant Cell Tiss Org Cult* 52:75–77
- Langens-Gerrits MM, De Klerk G-JM (1999) Micropropagation of flower bulbs: Lily and Narcissus. *Methods Mol Biol* 11:141–147
- Li SY, Chen C, Zhang HQ, Guo HY, Wang H, Wang L, Zhang X, Hua SN, Yu J, Xiao PG, Li RS, Tan X (2005) Identification of natural compounds with antiviral activities against SARS-associated coronavirus. *Antiviral Res* 67(1):18–23
- Maelicke A (2000) Allosteric modulation of nicotinic receptors as a treatment strategy for Alzheimer's disease. *Dement Geriatr Cogn Disord* 11:11–18
- Malik S, Cusidó RM, Mirjalili MH, Moyano E, Palazón J, Bonfill M (2011) Production of the anticancer drug taxol in *Taxus baccata* suspension cultures: a review. *Process Biochem* 46(1):23–34
- Malik M, Bach A (2017) High-yielding repetitive somatic embryogenesis in cultures of *Narcissus* L. 'Carlton'. *Acta Sci Pol Hortorum Cultus* 16(2):107–112
- Muraseva DS, Novikova TI (2018) Efficient protocol for *in vitro* propagation from bulb scale explants of *Fritillaria ruthenica* Wikstr. (Liliaceae), a rare ornamental species. *Rend Lincei-Sci Fis* 29(2):491–497
- Nasircilar AG, Mirici S, Karagüzel ÜÖ, Eren Ö, Baktir I (2011) *In vitro* propagation of endemic and endangered *Muscari mirum* from different explant types. *Turk J Bot* 35(1):37–43
- Niimi Y, Onozawa T (1979) *In vitro* bulblet formation from leaf segments of lilies, especially *Lilium rubellum* Baker. *Sci Hortic* 11(4):379–389
- Ozel CA, Khawar KM, Unal F (2007) *In vitro* axillary bulblet regeneration of Turkish yellow grape hyacinth (*Muscari macrocarpum* Sweet) from twin scale explants. *Res J Agric Biol Sci* 3(6):924–929
- Ozel CA, Khawar KM, Unal F (2015) Factors affecting efficient *in vitro* micropropagation of *Muscari muscarimi* Medikus using twin bulb scale. *Saudi J Biol Sci* 22(2):132–138
- Pierik R, Steegmans H (1975) Effect of auxins, cytokinins, gibberellins, abscisic acid and ethephon on regeneration and growth of bulblets on excised bulb scale segments of hyacinth. *Physiol Plant* 34(1):14–17
- Podwyszyńska M, Kosson R, Treder J (2015) Polyamines and methyl jasmonate in bulb formation of *in vitro* propagated tulips. *Plant Cell Tiss Org Cult* 123(3):591–605
- Ptak A, Simlat M, Kwiecień M, Laurain-Mattar D (2013) *Leucojum aestivum* plants propagated in *in vitro* bioreactor culture and on solid media containing cytokinins. *Eng Life Sci* 13(3):261–270
- Rahimi Khonakdari M, Mirjalili MH, Gholipour A, Rezadoost H, Moridi Farimani M (2018) Quantification of galantamine in *Narcissus tazetta* and *Galanthus nivalis* (Amaryllidaceae) populations growing wild in Iran. *Plant Genet Resour* 16(2):188–192
- Ravishankar G, Ramachandra Rao S (2000) Biotechnological production of phyto-pharmaceuticals. *J Biochem Mol Biol Bioph* 4:73–102
- Rice L, Finnie J, Van Staden J (2011) *In vitro* bulblet production of *Brunsvigia undulata* from twin-scales. *S Afr J Bot* 77(2):305–312
- Roy M, Liang L, Xiao X, Feng P, Ye M, Liu J (2018) Lycorine: a prospective natural lead for anticancer drug discovery. *Biomed Pharmacother* 107:615–624
- Sage DO, Lynn J, Hammatt N (2000) Somatic embryogenesis in *Narcissus pseudonarcissus* cvs. Golden Harvest and St. Keverne. *Plant Sci* 150(2):209–216
- Sairam R, Franklin G, Hassel R, Smith B, Meeker K, Kashikar N, Parani M, Abed DA, Ismail S, Berry K (2003) A study on the effect of genotypes, plant growth regulators and sugars in promoting plant regeneration via organogenesis from soybean cotyledonary nodal callus. *Plant Cell Tiss Org Cult* 75(1):79–85
- Santos J, Santos I, Salema R (1998) *In vitro* production of bulbs of *Narcissus bulbocodium* flowering in the first season of growth. *Sci Hortic* 76(3–4):205–217
- Santos I, Salema R (2000) Promotion by jasmonic acid of bulb formation in shoot cultures of *Narcissus triandrus* L. *Plant Growth Reg* 30(2):133–138
- Sarmadi M, Karimi N, Palazón J, Ghassempour A, Mirjalili MH (2019) Improved effects of polyethylene glycol on the growth, antioxidative enzymes activity and taxanes production in a *Taxus baccata* L. callus culture. *Plant Cell Tiss Org Cult* 137(2):319–328
- Sellés M, Bergoñón S, Viladomat F, Bastida J, Codina C (1997) Effect of sucrose on growth and galanthamine production in shoot-clump cultures of *Narcissus confusus* in liquid-shake medium. *Plant Cell Tiss Org Cult* 49(2):129–136
- Slabbert M, De Bruyn M, Ferreira D, Pretorius J (1993) Regeneration of bulblets from twin scales of *Crinum macowanii* *in vitro*. *Plant Cell Tiss Org Cult* 33(2):133–141
- Sochacki D (2008) The use of ELISA in the micropropagation of virus-free Narcissus. In: X international symposium on flower bulbs and herbaceous perennials, vol 886, pp 253–258
- Staikidou I, Watson S, Harvey BM, Selby C (2005) Narcissus bulblet formation *in vitro*: effects of carbohydrate type and osmolarity of the culture medium. *Plant Cell Tiss Org Cult* 80(3):313–320

- Stanilova MI, Molle ED, Yanev SG (2010) Galanthamine production by *Leucojum aestivum* cultures in vitro. In: Cordel GA (ed) The alkaloids: chemistry and biology, 68. Elsevier Inc, Amsterdam, pp 167–270
- Sutlana J, Sutlana N, Siddique M, Islam A, Hossain M, Hossain T (2010) In vitro bulb production in *Hippeastrum (Hippeastrum hybridum)*. J Cent Eur Agric 11(4):469–474
- Szlávik L, Gyuris Á, Minárovits J, Forgo P, Molnár J, Hohmann J (2004) Alkaloids from *Leucojum vernum* and antiretroviral activity of Amaryllidaceae alkaloids. Planta Med 70(9):871–873
- Taha LS, Sayed SS, Farahat MM, El-Sayed IM (2018) In vitro culture and bulblets induction of Asiatic hybrid lily 'red alert'. J Biol Sci 18:84–91
- Tarakemeh A, Azizi M, Rowshan V, Salehi H, Spina R, Dupire F, Arouie H, Laurain-Mattar D (2019) Screening of Amaryllidaceae alkaloids in bulbs and tissue cultures of *Narcissus papyraceus* and four varieties of *N. tazetta*. J Pharm Biomed Anal 172:230–237
- Teixeira da Silva JA, Kulus D, Zhang X, Zeng S, Ma G, Piqueras A (2016) Disinfection of explants for saffron (*Crocus sativus*) tissue culture. Environ Exp Biol 14:183–198
- Thomas TD (2007) Pretreatment in thidiazuron improves the in vitro shoot induction from leaves in *Curculigo orchioides* Gaertn., an endangered medicinal plant. Acta Physiol Plant 29(5):455–461
- Torras-Claveria L, Berkov S, Codina C, Viladomat F, Bastida J (2013) Daffodils as potential crops of galanthamine. Assessment of more than 100 ornamental varieties for their alkaloid content and acetylcholinesterase inhibitory activity. Ind Crops Prod 43:237–244
- Trujillo-Chacón LM, Pastene-Navarrete ER, Bustamante L, Baeza M, Alarcón-Enos JE, Céspedes-Acuña CL (2020) In vitro micropropagation and alkaloids analysis by GC–MS of Chilean Amaryllidaceae plants: *Rhodophiala pratensis*. Phytochem Anal 31(1):46–56
- Ulrich MR, Davies FT Jr, Koh YC, Duray SA, Egilla JN (1999) Micropropagation of *Crinum* 'Ellen Bosanquet' by tri-scales. Sci Hortic 82:95–102
- Wang L, Du Y, Rahman MM, Tang B, Fan L-J, Kilaru A (2018) Establishment of an efficient in vitro propagation system for *Iris sanguinea*. Sci Rep 8(1):1–10
- Woodruff-Pak DS, Lander C, Geerts H (2002) Nicotinic cholinergic modulation: galanthamine as a prototype. CNS Drug Rev 8(4):405–426
- Yang X, Huang Y, Fan G (2013) Effects of different photoperiods on in vitro plantlet regeneration of *Paulownia* plants. J Chem Pharm Res 5(12):1446–1450
- Yang L, Wen K-S, Ruan X, Zhao Y-X, Wei F, Wang Q (2018) Response of plant secondary metabolites to environmental factors. Molecules 23(4):762
- Zayed R, El-Shamy H, Berkov S, Bastida J, Codina C (2011) In vitro micropropagation and alkaloids of *Hippeastrum vittatum*. In Vitro Cell Dev Biol Plant 47(6):695–701
- Zhang X, Gao J, Peng Z (2013) Callus induction, differentiation and histological observation of *Narcissus tazetta* var. *chinensis* floral organs. Forest Res 26(3):320–325

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.