Correction to: contrasting effects of urbanization on arboreal and ground-dwelling land snails: role of trophic interactions and habitat fragmentation

Ikuyo Saeki¹ • Shigeru Niwa² • Noriyuki Osada³ • Wakana Azuma⁴ • Tsutomu Hiura⁵

Published online: 3 March 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

Correction to: Urban Ecosyst

<https://doi.org/10.1007/s11252-020-00930-6>

The original version of the article unfortunately contained mistakes in the legends of Fig. 3.

The legends (1) Blue square: *Euhadra brandtii sapporo*, ground is changed to Blue square: *Euhadra brandtii sapporo*, canopy (2) Red circle: *Euhadra brandtii sapporo*, canopy is changed to Red circle: *Euhadra brandtii sapporo*, ground.

The authors apologize for this mistake and the updated figure is shown below.

The online version of the original article can be found at <https://doi.org/10.1007/s11252-020-00930-6>

✉ Ikuyo Saeki
saeki.ikuyo.ge@u.tsukuba.ac.jp

¹ Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8571, Japan

² Japan Wildlife Research Center, 3-3-7 Kotobashi, Sumida-ku, Tokyo 130-8606, Japan

³ Laboratory of Plant Conservation Science, Faculty of Agriculture, Meijo University, Nagoya, Aichi 468-8502, Japan

⁴ Present address: Graduate School of Agricultural Science, Kobe University, Kobe, Hyogo 657-8501, Japan

⁵ Field Science Center, Hokkaido University, Kita 9 Nishi 9, Sapporo, Hokkaido 060-0809, Japan

Fig. 3 Comparison of survival curves of *Euhadra brandtii sapporo* and *Ezohelix gainesi* for canopy and ground treatments in the 2017-fall (left) and 2018-summer (right) experiments. Error bars indicate 1 SE based on results of 8 samples $\times$ 3 sites (i.e., $n = 24$) for one forest type. See Table 2 for results of pairwise comparisons by log-rank tests