ERRATUM

Erratum to: Angiomyolipoma being surgically excised for presumed kidney carcinoma

Weibin Hou¹ · He Xiao¹ · Guanghua Liu¹ · Zhigang Ji¹

Published online: 17 December 2015
© Springer Science+Business Media Dordrecht 2015

Erratum to: Int Urol Nephrol (2015) 47:1037–1043 DOI 10.1007/s11255-015-0996-y

Unfortunately, some errors occurred in the original publication of the article. The authors apologize to the editors and readers for an error in the second sentence in the Materials and methods section of Abstract and Table 1. The corrected Abstract and the new version of Table 1 are given below.

Abstract

Materials and methods From 2008 to 2014, AML was pathologically confirmed in 38 patients who underwent

radical or partial nephrectomy for presumed malignant renal tumor. Control group 1 were patients with renal cell carcinoma (RCC) matched for sex and tumor size; control group 2 were patients with typical AML matched for age and sex. Pertinent data of the studied group and its matched control groups were recorded and analyzed.

The online version of the original article can be found under doi:[10.1007/s11255-015-0996-y](https://doi.org/10.1007/s11255-015-0996-y).

✉ Zhigang Ji
jizg1129@163.com

¹ Department of Urology, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, 1#, Shuaifuyuan, Dongcheng District, Beijing 100730, People's Republic of China

Table 1 Comparison of misdiagnosed AML group and its matched RCC control group

Parameter	All patients (<i>n</i> = 76)	AMLs (<i>n</i> = 38)	RCCs (<i>n</i> = 38)	<i>P</i>
Sex				1
Male	34	17	17	
Female	42	21	21	
Tumor size (cm)				0.930
Mean $\pm$ SD	3.14 $\pm$ 1.68	3.12 $\pm$ 1.68	3.16 $\pm$ 1.70	
Range	0.9–9.6	0.9–9.4	0.9–9.6	
Age (years)				0.001
Mean $\pm$ SD	52.54 $\pm$ 12.41	48.11 $\pm$ 12.92	56.97 $\pm$ 10.23	
Range	20–80	20–74	31–80	
Laterality				0.818
Left	39	19	20	
Right	37	19	18	
Presentation				1
Incidental	6	3	3	
Symptomatic	70	35	35	
Ultrasonography				0.526
Hypoechoic	50	24	26	
Isoechoic	10	4	6	
Hyperechoic	11	7	4	
Fat content on CT				0.077
Yes	3	3	0	
No	73	35	38	
Nephrectomy type				1
Radical	16	8	8	
Partial	60	30	30	
Surgical approach				0.238
Open	29	17	12	
Laparoscopy	47	21	26	