CORRECTION



Correction to: Radiomics analysis of [18F]-fluoro-2-deoxyglucose positron emission tomography for the prediction of cervical lymph node metastasis in tongue squamous cell carcinoma

Published online: 11 June 2022

© The Author(s) under exclusive licence to Japanese Society for Oral and Maxillofacial Radiology 2022

Correction to: Oral Radiology

https://doi.org/10.1007/s11282-022-00600-7

In the original publication of the article, the tables 4 and 6 were published incorrectly. The correct Tables 4 and 6 are given in this correction.

The original article can be found online at https://doi.org/10.1007/s11282-022-00600-7.

- ☐ Takaharu Kudoh kudoh@tokushima-u.ac.jp
- Department of Oral Surgery, Tokushima University Graduate School of Biomedical Sciences, Kuramoto-cho, Tokushima, Japan
- Department of Medical Image Informatics, Tokushima University Graduate School of Biomedical Sciences, Kuramoto-cho, Tokushima, Japan
- Department of Therapeutic Radiology, Tokushima University Graduate School of Biomedical Sciences, Kuramoto-cho, Tokushima, Japan
- Department of Oral Bioscience, Tokushima University Graduate School of Biomedical Sciences, Kuramoto-cho, Tokushima, Japan



52 Oral Radiology (2023) 39:51–52

Table 4 Clinicopathological factors

Factors	Total
Differentiation	
Well	20
Moderate	19
Poorly	1
Y-K classification	
2	2
3	14
4c	17
4d	7

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Table 6 Clinicopathological characteristics of patients stratified by neck status

Characteristics	Neck status	
	Positive	Negative
Sex		
Male	13	14
Female	6	7
Age (years)*	68 ± 13	64 ± 16
Clinical T classification		
1	2	5
2	5	7
3	7	6
4a	5	2
4b	1	0
Differentiation		
Well	11	9
Moderate	8	11
Poorly	1	0
Y-K classification		
2	1	1
3	5	9
4c	12	5
4d	2	5
DOI, mm	10 ± 7	8 ± 5

DOI depth of invasion



 $[*]Mean \pm standard deviation$