

Erratum to: Serially coupling hydrophilic interaction and reversed-phase chromatography with simultaneous gradients provides greater coverage of the metabolome

Jennifer Haggarty · Madalina Oppermann ·
Matthew J. Dalby · Richard J. Burchmore ·
Ken Cook · Stefan Weidt · Karl E. V. Burgess

Published online: 3 February 2015
© Springer Science+Business Media New York 2015

Erratum to: Metabolomics
DOI: 10.1007/s11306-014-0770-7

This paper was originally published under the title ‘Serially coupling hydrophobic interaction and reversed-phase chromatography with simultaneous gradients provides greater coverage of the metabolome’. The word “hydrophobic” is incorrect. The title should read ‘Serially coupling hydrophilic interaction and reversed-phase chromatography with simultaneous gradients provides greater coverage of the metabolome’.

The online version of the original article can be found under
doi:[10.1007/s11306-014-0770-7](https://doi.org/10.1007/s11306-014-0770-7).

J. Haggarty · R. J. Burchmore · S. Weidt ·
K. E. V. Burgess (✉)
Polyomics, University of Glasgow, 211 Wolfson Wohl
Translational Cancer Research Centre, Garscube Campus,
Glasgow G61 1QH, UK
e-mail: karl.burgess@glasgow.ac.uk

M. Oppermann · K. Cook
Thermo Fisher Scientific, Hemel Hempstead, UK

M. J. Dalby
Centre for Cell Engineering, University of Glasgow, Glasgow,
UK