Correction to: biocide emissions from building materials during wet weather: identification of substances, mechanism of release and transfer to the aquatic environment

Claudia Paijens^{1,2} • Adèle Bressy¹ • Bertrand Frère² • Régis Moilleron³

Published online: 26 December 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: Environmental Science and Pollution Research
<https://doi.org/10.1007/s11356-019-06608-7>

The original publication of this paper contains a mistake.
The text in page 17 belongs to the footnote of Table 6.
The original article was corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s11356-019-06608-7>

✉ Adèle Bressy
adele.bressy@enpc.fr

¹ Leesu, UMR-MA-102, Ecole des Ponts ParisTech, Université Paris-Est Créteil, AgroParisTech, 6 et 8 avenue Blaise Pascal – Cité Descartes, 77455 Champs-sur-Marne Cedex 2, France

² Laboratoire Central de la Préfecture de Police, 39 bis rue de Dantzig, 75015 Paris, France

³ Leesu, UMR-MA-102, Université Paris-Est Créteil, Ecole des Ponts ParisTech, AgroParisTech, 61 avenue du Général de Gaulle, 94010 Créteil Cedex, France

Table 6 Surface water concentrations (ng/L) of some biocides used either in building materials or as PPP

Country	Concentrations in surface water (ng/L)									
	France	Germany	Spain	Switzerland	Other European countries	North America	Brazil	China	Australia	Tanzania
References	a	b	c	d	e	f	g	h	i	j
Carbendazim	1	< 2.5–116	0.04–11.6	2.9–206	< 0.33–269	21–360	< 8	6.6–610		
IPBC		< 5		< 30						
Diuron	20–5600	2–41	7.5–8550	1.1–310	< 0.51–290	< 1–910	< 8	26–51		
Isoproturon	10–61	< 1–115	2.4–9.1	1.1–5500	1–340					
Cybutryn		0.9–12		< 4.5–6	0.1	< 7	7.7–15.9			
Terbutryn	10	5.3–5600	0.9–30.5	< 4.5–50	1–22	< 10			1–22	
Terbuthylazine	10	2.3–372	0.1–12600	5–40	< 3–254.4		102.2			
Propiconazole	12	4.2–7		1.9–65	< 4.5–44	< 3–291		10	2–22	
Tebuconazole	14	2.2–12	1.7–15.4	1.9–86	< 2.7–120	< 3–53	5.6–60.6		21	
Thiabendazole		0.1–21	0.01–1		< 10–53	< 3.9–153.2		< 0.01		170
Permethrin	14				< 0.5–97	< 0.003–18	< 375–1650	370		
Mecoprop	5–30	3–147		4–1750	1.7–30	5.1–94				

^a Blanchoud et al. (2004); Botta et al. (2012); Direction régionale et interdépartementale de l'environnement et de l'énergie d'Ile-de-France (DRIEE) (2016), ^b Quednow and Püttmann (2007); Van Pinxteren et al. (2009); Loos et al. (2010); Wick et al. (2010); Reemtsma et al. (2013); Wluka et al. (2016), ^c Claver et al. (2006); Benvenuto et al. (2010); Gros et al. (2012); Köck-Schulmeyer et al. (2012); Campo et al. (2013); Herrero-Hernández et al. (2013); Masiá et al. (2013); Ccanccapa et al. (2016), ^d Öllers et al. (2001); Singer et al. (2010); Bonvin et al. (2011); Wittmer et al. (2011a); Moschet et al. (2015), ^e Bonwick et al. (1995); Benvenuto et al. (2010); Stamatis et al. (2010); Matamoros et al. (2012); Emelogu et al. (2013); Kalogridi et al. (2014); Palma et al. (2014); Vorkamp et al. (2014); Antić et al. (2015); Chitescu et al. (2015); Stipaničev et al. (2017); ^f Bartelt-Hunt et al. (2009); Murray et al. (2010); Battaglin et al. (2011); Struger et al. (2011); Weston and Lydy (2012); Sengupta et al. (2013); Metcalfe et al. (2016), ^g Pinheiro et al. (2011); Caldas et al. (2013); Robles-Molina et al. (2014), ^h Chen et al. (2012); Heeb et al. (2012), ⁱ Allinson et al. (2015), ^j Kishimba (2004)