Correction to: Hemotoxic effects of some herbicides on juvenile of Nile tilapia *Oreochromis niloticus*

Mohamed Fathy¹ · Ibrahim A. Mohamed¹ · Ahmed I. A. Farghal¹ · Sobhy A. H. Temerak¹ · Alaa El-Din H. Sayed²

Published online: 21 December 2019
 © Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: Environmental Science and Pollution Research (2019) 26:30857–30865
<https://doi.org/10.1007/s11356-019-06280-x>

The correct presentation of Table 2 is shown in this paper.

Table 2: Biochemical variables in *O. niloticus* exposed to sbu-lethal concentrations of certain herbicides for 96 h (Mean ± S.D).

Parameters	Treatments						
	Control	Acetochlor	Bentazon	Bispyribac-sodium	Halosulfuron-methyl	Bensulfuron-methyl	Quinclorac
AST (μ/l)	(54.25± 1.24) ^b	(66.42 ± 2.60) ^a	(56.3± 4.08) ^b	(55.72 ± 2.05) ^b	(53.75 ± 0.98) ^b	(54.62± 2.00) ^b	(57.17±4.25) ^b
ALT (μ/l)	(27.05± 1.84) ^c	(31.1 ± 1.53) ^a	(29.29± 1.49) ^{abc}	(28.82 ± 1.66) ^{abc}	(28.5± 0.75) ^{abc}	(30.25 ± 0.95) ^{ab}	(27.92 ± 2.83) ^{bc}
ALP (μ/l)	(24.37±0.34) ^b	(29.4±3.42) ^a	(31.04 ± 1.39) ^a	(29±0.67) ^a	(30.42 ± 0.97) ^a	(30 ± 1.41) ^a	(26.55±0.81) ^b
Glucose (mg/dl)	(96.25 ± 1.33) ^a	(97.92 ± 9.05) ^a	(101.2 ± 6.49) ^a	(106.95±11.00) ^a	(101 ± 6.61) ^a	(104.35±6.94) ^a	(106.61 ± 5.93) ^a
Cholesterol (mg/l)	(169.52±27.88) ^b	(201 ± 4.96) ^a	(185.75±10.21) ^{ab}	(198.5 ± 10.40) ^a	(189±6.78) ^{ab}	(198± 8.52) ^a	(195.5±7.04) ^a
Total protein (g/dl)	(5.05±0.86) ^b	(6.45±0.86) ^a	(5.42±0.34) ^b	(5.67 ± 0.18) ^{ab}	(5.5±0.83) ^b	(5.05 ± 0.17) ^b	(5.22 ± 0.23) ^b
Uric acid (mg/dl)	(12.32 ± 0.15) ^c	(17.22 ± 2.05) ^a	(14 ± 0.29) ^b	(15.2 ± 0.86) ^b	(13.90 ± 0.39) ^b	(13.92 ± 0.55) ^b	(13.67 ± 1.09) ^{bc}
Creatinine (mg/dl)	(0.58 ± 0.18) ^a	(0.74 ± 0.02) ^a	(0.23 ± 0.07) ^b	(0.61 ± 0.06) ^a	(0.30 ± 0.12) ^b	(0.58 ± 0.06) ^a	(0.60 ± 0.15) ^a
Albumin (g/dl)	(0.82 ± 0.47) ^c	(1.50 ± 0.06) ^a	(1.12 ± 0.02) ^b	(1.27 ± 0.05) ^{ab}	(1.18 ± 0.05) ^{ab}	(1.22 ± 0.08) ^{ab}	(1.32 ± 0.22) ^{ab}
Globulin (g/dl)	(2.09 ± 0.03) ^d	(2.37 ± 0.06) ^a	(2.14 ± 0.05) ^{cd}	(2.24 ± 0.06) ^{bc}	(2.14 ± 0.05) ^{cd}	(2.19 ± 0.04) ^{bcd}	(2.28 ± 0.13) ^{ab}
A/G ratio	(0.38 ± 0.22) ^b	(0.63 ± 0.02) ^a	(0.52 ± 0.01) ^a	(0.56 ± 0.01) ^a	(0.55 ± 0.02) ^a	(0.55 ± 0.03) ^a	(0.57 ± 0.06) ^a

Means with the same superscript letters in the same row for each biochemical parameters are not significantly different ($P < 0.05$)

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s11356-019-06280-x>

✉ Alaa El-Din H. Sayed
 alaasayed@aun.edu.eg; alaa_h254@yahoo.com

¹ Plant Protection Department, Faculty of Agriculture, Assiut University, Assiut 71526, Egypt

² Zoology Department, Faculty of Sciences, Assiut University, Assiut 71516, Egypt