Correction to: Recalibration and validation of the Swiss lichen bioindication methods for air quality assessment

Rolf Herzig¹ · Christian Schindler^{2,3} · Martin Urech⁴ · Beat Rihm⁵ · Hanspeter Lötscher⁶ · Georg Thomann⁶

Published online: 22 June 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Environmental Science and Pollution Research

<https://doi.org/10.1007/s11356-020-09001-x>

The correct fig. 2 is presented in this paper.

The original article is corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s11356-020-09001-x>

✉ Rolf Herzig
rolf.herzig.agb@bluewin.ch

¹ AGB, Arbeitsgemeinschaft für Bioindikation, Umweltbeobachtung und ökologische Planung, Quartiergasse 12, 3013 Bern, Switzerland

² Swiss Tropical and Public Health Institute, Basel, Switzerland

³ University of Basel, Basel, Switzerland

⁴ puls, Umweltberatung, Bern, Switzerland

⁵ Meteotest AG, Bern, Switzerland

⁶ Amt für Natur und Umwelt, Chur, Switzerland

Fig. 2 Correlations (R^2_{adj}) of new lichen-indices for primary pollutants, NH_3 and AOT40f and of the traditional index IAP18 with logtransformed air pollutant and meteo variables for entire Switzerland. a The new primary pollutant lichen index (PP-LI) shows a high multivariable correlation ($\text{adj } R^2 = 0.86$) with primary pollutant variables $\ln(\text{PM}_{10})$, $\ln(\text{SO}_2)$, $\ln(\text{NO}_2)$ (left blue bar), followed by bivariate correlations with individual primary pollutant variables (other blue bars). The bivariate correlations with the other pollutant variables ($\ln(\text{NH}_3)$ and $\ln(\text{AOT40f})$, dark gray bars) and the log-transformed meteo variables (light gray bars) follow on the right side of the bar graphs. b AOT40f

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.