Correction to: Trace elements exposure influences proximate body composition and antioxidant enzyme activities of the species tilapia and catfish in Burullus Lake-Egypt: human risk assessment for the consumers

Heba H. Abdel-Kader¹ • Mohamed H. Mourad¹

Published online: 17 August 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Environmental Science and Pollution Research
<https://doi.org/10.1007/s11356-020-10207-2>

The correct images of figs. 2 and 3 are presented in this paper.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s11356-020-10207-2>

✉ Heba H. Abdel-Kader
hebaelalkamy3232@gmail.com

¹ National Institute of Oceanography and Fisheries, NIOF,
Alexandria, Egypt

Fig. 2 Proximate body composition in muscle tissue of tilapia species and *C. gariepinus*. Data are expressed as mean values and standard errors of the mean (SEM) ($n = 6$). Different letters are significantly different $P < 0.05$ in the same graph

Fig. 3 Antioxidant enzyme activity in the liver tissue of tilapia species and *C. gariepinus*. Data are expressed as mean values and standard errors of the mean (SEM) ($n = 6$). Different letters are significantly different $P < 0.05$ in the same graph