Biowastes of slaughterhouses and wet markets: an overview of waste management for disease prevention

Adel Al-Gheethi¹ · Nyuk Ling Ma² · Parveen Fatemeh Rupani³ · Naznin Sultana⁴ · Maizatul Azrina Yaakob⁵ · Radin Maya Saphira Radin Mohamed¹ · Chin Fhong Soon⁶

Received: 16 December 2020 / Accepted: 16 September 2021

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2021

Abstract

Slaughterhouse and wet market wastes are pollutants that have been always neglected by society. According to the Food and Agriculture Organization of the United Nations, more than three billion and nineteen million livestock were consumed worldwide in 2018, which reflects the vast amount and the broad spectrum of the biowastes generated. Slaughterhouse biowastes are a significant volume of biohazards that poses a high risk of contamination to the environment, an outbreak of diseases, and insecure food safety. This work comprehensively reviewed existing biowaste disposal practices and revealed the limitations of technological advancements to eradicate the threat of possible harmful infectious agents from these wastes. Policies, including strict supervision and uniform minimum hygienic regulations at all raw food processing factories, should therefore be tightened to ensure the protection of the food supply. The vast quantity of biowastes also offers a zero-waste potential for a circular economy, but the incorporation of biowaste recycling, including composting, anaerobic digestion, and thermal treatment, nevertheless remains challenging.

Keywords Bacteria · Disease transmission · Sustainable management · Biological treatment · Hazards' risk · Recycle

Introduction

Inedible animal body parts that represent 45% of slaughtered animals, which include the tissues, blood, blood vessels, feathers, and bones, are major constituents of slaughterhouse and wet market wastes that are ultimately sent for disposal or recycling (Franke-Whittle and Insam 2013). The disposal of these substantial amounts of wastes is challenging for the meat, aquatic, and poultry industries due to several legal

restraints and ever-rising processing costs associated with disposal. Unsafe and improper disposal of such wastes can lead to tremendous ecological troubles. Table 1 shows the physicochemical properties of biowastes generated in slaughterhouses and wet markets globally. The contents of slaughterhouse and wet market biowastes, such as moisture, organic carbon solids, and elements such as potassium, are potentially reservoirs for microbial diseases (Mensah-Attipoe and Toyinbo 2019).

Adel Al-Gheethi and Nyuk Ling Ma are both first authors and contribute the same in the manuscript.

Responsible Editor: Ta Yeong Wu

✉ Parveen Fatemeh Rupani
parveen@ujs.edu.cn

✉ Chin Fhong Soon
soon@uthm.edu.my

¹ Micropollutant Research Centre (MPRC), Faculty of Civil Engineering & Built Environment, Universiti Tun Hussein Onn Malaysia, 86400 Parit Raja, Batu Pahat, Johor, Malaysia

² Faculty of Science and Marine Environment, University Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu, Malaysia

³ School of Energy and Power Engineering, Jiangsu University, Zhenjiang 212013, Jiangsu, China

⁴ Medical Academy, Prairie View A&M University, Prairie View, TX 77446, USA

⁵ Institute for Integrated Engineering, Universiti Tun Hussein Onn Malaysia, 86400 Parit Raja, Batu Pahat, Johor, Malaysia

⁶ Microelectronics and Nanotechnology-Shamsuddin Research Centre, Institute for Integrated Engineering, Universiti Tun Hussein Onn Malaysia, 86400 Parit Raja, Batu Pahat, Johor, Malaysia

Table 1 Physicochemical properties of slaughterhouse and wet market biowastes (NR indicates non-reported)

Physicochemical properties	Slaughterhouse		Wet market		
	Sengupta (2004)	Gendebien et al. (2001)	Tweib et al. (2011)	Lohri et al. (2016)	Lou et al. (2015)
Moisture	69.45%	NR	60%	22.3%	75.5 ± 6.4%
Total solids	30.55%	10–180 g/m ³	510.48 kg/ m ³	25.9%	34.7 ± 2.3 kg/m ³
Volatile solids	87.95%	10–107 g/m ³	NR	87.2%	NR
Fixed solids	12.05%	NR	NR	NR	NR
Organic carbon	23.32%	NR	45.16%	12.8%	35.2 ± 0.7%
Total nitrogen	2.71%	NR	1.4%	1.5%	2.0 ± 0.1%
Phosphorus	4.19 mg/g	NR	110970.826 ppb	NR	NR
Potassium	6.9 mg/g	NR	NR	NR	NR
Organic nitrogen	NR	2–18 g/m ³	NR	NR	NR
Ammonia nitrogen	NR	0.6–2.2 g/m ³	NR	NR	NR
pH	NR	NR	6.2	NR	5.69 ± 0.01

In addition, the improper disposal of slaughterhouse biowastes (SH-biowastes) has caused harmful algal blooms in surface water (Arvanitoyannis and Kassaveti 2008) and the contamination of groundwater from antibiotics from the poultry production (Alam et al. 2019), therefore raising safety concerns. However, the implementation of proper waste management has been limited by legal restraints, cost affordability, and available technologies. To date, several regulations for SH-biowaste treatment in minimizing pathogen transmission have been enforced, such as manure composting regulations by the Environmental Protection Agency (EPA)(Erickson et al. 2014), SH-biowaste thermal treatment regulations by the Canadian Food Inspection Agency (CFIA)(Adhikari et al. 2018), and pathogen disease transmission and ethical regulation (Regulation (EC) No 1774/2002) by the European Union (Russ and Meyer-Pittroff 2004). With the recent SARS-associated coronavirus (SARS-CoV) outbreak from a wet market, the proper management of wet markets, including most slaughterhouses, is being put under the spotlight. Hence, more transparent management is demanded. Although burial and burning are the most common disposal methods of animal wastes in underdeveloped countries (Anon 2002), the developed countries do not allow similar approaches. For example, the European Union controls the dumping by incineration through licensed waste collectors (NABC 2004). The use of modern incineration plants as the standard practice to manage SH-biowastes in most developed countries is favorable, as this method effectively kills all microbes (Alnahas et al. 2020). The biowaste composition from slaughterhouses and wet markets might be different from one region to another due to the population's location, lifestyle, and food consumption (Lohri et al. 2016). Differences in biowaste composition should be considered when

implementing efficient treatment techniques. Hence, integrated recycling technologies for slaughterhouse and wet market waste management are essential for the safe disposal of wastes and the removal of various infectious agents.

This review therefore highlights the current technology gaps in waste management practices based on the sources of SH-biowaste components and the implications on the ecosystem and human health. Proper implementation of waste management can reduce environmental and health issues, while introducing new business opportunities that will ultimately benefit economic growth.

Slaughterhouse and wet market wastes

Biowastes generated from the food industry (slaughterhouses and wet markets) are from a broad spectrum and are derived from different stages of food processing, and the wastes are classified into solid wastes and wastewater (Ho and Chu 2019; Ratnawati and Trihadiningrum 2014). SH-biowastes contain a high proportion of organic materials, but these are not suitable for human consumption. The typical body composition of different edible animals is presented in Table 2. Hot carcass is the body of an animal after the removal of internal organs. Generally, 49–76% of hot carcass is generated from cattle, 38–47% from sheep and goats, 44–48% from pigs, and 37–40% from broilers (Shija et al. 2013).

In small slaughterhouses in developing countries, the dressing of carcasses, the removal of hides, and evisceration are performed on the floor, which is also the location for cutting the edible portion of the slaughtered animals (Fearon et al. 2014). Slaughter and evisceration are performed in a simple process. The abattoirs' wastewater and solid wastes are

Table 2 Typical body compositions of calf, sheep, goat, pig, broiler chicken and duck (NR indicates non-reported)

Animal	Component	Weight (kg) Mean $\pm$ SE	Percentage (%)	Reference
Calf	Body (18 months)	414.50 $\pm$ 20.5	100	Oh et al. (2008)
	Hot carcass	250.70 $\pm$ 8.62	60.38	
	Boneless meat	173.74 $\pm$ 3.53	41.90	
	Trimmed carcass fat	24.76 $\pm$ 4.67	5.98	
Sheep	Body (14 to 24 months)	22.29 $\pm$ 0.55	100	Shija et al. (2013)
	Hot carcass	9.43 $\pm$ 0.03	42.30	
	Gastrointestinal tract	3.53 $\pm$ 0.10	15.84	
	Head	1.59 $\pm$ 0.04	7.90	
	Skin	1.66 $\pm$ 0.05	8.23	
	Feet	0.52 $\pm$ 0.02	2.61	
	Internal organs	0.94 $\pm$ 0.03	4.70	
	Kidneys	0.07 $\pm$ 0.01	0.35	
	Internal fats	0.07 $\pm$ 0.03	0.36	
	Testicles	0.23 $\pm$ 0.02	1.16	
	Tail	0.59 $\pm$ 0.04	2.91	
	Total non-carcass	7.76 $\pm$ 0.17	38.58	
Goat	Body (14 to 24 months)	20.50 $\pm$ 0.55	100	Diarra et al. (2015)
	Hot carcass	9.68 $\pm$ 0.03	47.20	
	Gastrointestinal tract	3.09 $\pm$ 0.10	15.10	
	Head	1.27 $\pm$ 0.04	6.83	
	Skin	1.39 $\pm$ 0.05	7.44	
	Feet	0.56 $\pm$ 0.02	3.02	
	Internal organs	0.97 $\pm$ 0.03	5.24	
	Kidneys	0.08 $\pm$ 0.01	0.44	
	Internal fats	0.11 $\pm$ 0.03	0.59	
	Testicles	0.09 $\pm$ 0.02	1.02	
	Tail	0.02 $\pm$ 0.04	0.10	
	Total non-carcass	6.68 $\pm$ 0.17	35.84	
Pig	Body	NR	100	Goldstrand (1992)
	Carcass meat	NR	52	
	Bones	NR	17	
	Organ	NR	7	
	Skin and attached fats	NR	6	
	Blood	NR	3	
	Fatty tissues	NR	3	
	Horns, huff, feet, and skull	NR	6	
	Abdominal and intestinal contents	NR	6	
Broiler chicken	Body (42 days)	2.19	100	Diarra et al. (2015)
	Dressing	1.79	81.52	
	Carcass cut-up parts	0.894	40.83	
	Liver weight	0.046	2.12	
	Pancreas	0.004	0.18	
	Gizzard weight	0.027	1.25	
	Heart	0.011	0.52	
	Small intestine	0.012	0.55	
	Abdominal fat pad	0.0007	0.30	
Duck	Body (7 weeks)	3.42 $\pm$ 1.02	100	Wawro et al. (2004)
	Carcass	2.43 $\pm$ 0.76	71	
	Giblets	0.16 $\pm$ 0.05	4.6	

Table 2 (continued)

Animal	Component	Weight (kg) Mean $\pm$ SE	Percentage (%)	Reference
	Abdominal fat	0.07 $\pm$ 0.03	2	
	Total meat	1.189 $\pm$ 0.382	35	

directly disposed into water bodies and lands adjacent to the abattoirs, creating pollution problems (Fearon et al. 2014). This is due to inadequate waste management and treatment facilities.

In contrast, modern abattoirs in developed countries are found in enclosed buildings with specific process lines for different types of slaughtered animals. The abattoirs are well-established with waste management processes for separating blood, screening of solids, and trapping of grease, which are aided with a conveyor line system (Woodard, Curran, Inc 2006). Blood is collected for stockfeed or fertilizer production. In waste treatment plants, bulk animal carcass incinerators are used to render animal wastes into ashes that can be used as fertilizers. Solid wastes, such as hides, hairs, intestines, and hooves, are screened and sorted for animal feed or disposal. Large amounts of water are used for the cleaning process, which results in effluents that contain solidified lipids. In advanced wastewater processing plants, grease, oil, fat, floating and suspended solids, and settling materials that rise to the surface of the water are removed regularly using mechanical pre-treatment systems, such as the micro strainer, rotary drum fine screener, and screw conveyor (for example, the Huber technology from Germany). Wastewater from the abattoirs is processed using a full automation system and further treated before being discharged into municipal sewers. In modern abattoirs, all animal wastes are recycled, while ensuring minimum effects to the environment.

The wastewater produced by the meat processing industry has increased tremendously in recent years (Aleksic et al. 2020). According to the Food and Agriculture Organization (FAO) of the United Nations (FAO 2021), the demand for broiler chicken and meat is growing faster than any other meat type. Other types of poultry, such as duck, turkey, guinea fowl, pigeon, and geese, are far less consumed, which is 1.2 billion per year (FAO 2021). The number of chickens slaughtered, at approximately 24.5 billion per year, surpasses those of pigs and cattle (Fig. 1). Moreover, the amount of water required by processing poultry is an average of 15 L per bird, which could add up to a few hundred billion liters of wastewater each year (Matsumura and Mierzwa 2008). Effluents from meat processing are recognized as harmful and threatening due to the complex composition of fibers, proteins, fats, high organic contents, pathogens, and veterinary pharmaceuticals (Bustillo-Lecompte and Mehrvar

2017; Chang et al. 2010). The chemical contents of wastewater from slaughterhouses are almost similar to municipal sewage (Damaceno et al. 2019). These induce algal blooms that limit oxygen supply for the aquatic life and cause microbiological contaminations of water bodies. Hence, the implications of the astonishing volume and wide range of SH-biowastes disposed into the ecological system from slaughterhouses and wet markets should not be disregarded (Fig. 2).

The major biowastes from marine products are from fish, representing 76 million tons, mostly from wholesalers and the seafood processing industry that processed canned and frozen fish (Ching-Velasquez et al. 2020). Majority of seafood are processed in shore-based processing facilities near the ports. During the processing of demersal fish, for instance, the viscera, frame, skin, fin, and head are usually discarded as waste (Ravanipour et al. 2021). The waste products of shellfish are shells and viscera. The inedible portions of some demersal species and shellfish, as outlined in Table 3, are usually discarded. However, aquatic wastes are considered valuable by-products that are recyclable as fertilizers, animal feed, dietetic products, natural colorings, biodiesel, and enzymes (Arvanitoyannis and Kassaveti 2008). In addition, every thousand kg of fish production during skinning and canning generated 15 to 17 liters of wastewater (Arvanitoyannis and Kassaveti 2008), and these effluents consist of feces, antibiotics, fish feed, and components that are rich in protein and can be post-processed to be used as animal feed or manure. However, effluents from the fish processing industry are often released directly into the shallow coastal water, leaving impacts on the aquatic environment (Arvanitoyannis and Ladas 2008).

Ecological and health implications

Wet market and slaughterhouse facilities are usually located near water bodies, and therefore, untreated or partially treated effluents of these facilities commonly ended up in the water bodies (Fig. 2). SH-biowastes generated from slaughterhouses contain a high level of organic compounds, which could lead to rapid oxygen consumption by microorganisms, resulting in an anoxia condition in water bodies (Adeyemo 2002). Hence, the anaerobic stage causes an insufficient amount of dissolved oxygen for aquatic life and plants. Consequently, the high

Fig. 1 Animal production by country, 2018 (Food and Agriculture Organization United Nation, <http://www.fao.org/faostat/en/>, 2018)

Fig. 2 SH-biowaste contaminations from slaughterhouses and wet markets and the effects on ecological system and human food chain

level of biochemical oxygen demand (BOD) tends to increase the concentrations of nitrates and ammonia in water bodies, which causes an increase in uncontrollable algae blooms.

In addition to organic wastes, the extensive use of arsenic in the poultry industry has raised ecological concerns (Hileman 2007; Lee et al. 2010). The application of arsenic is to increase the weight of poultry, prevent germs or parasites, and enhance the skin color of the poultry. But it has been reported that arsenic found in water bodies gives adverse effects on plants' growth, causes accumulation of microbe, and is closely related to cancer and skin diseases (Adesiyun et al. 2018).

Another biohazard from slaughterhouses and wet markets is the discharged antibiotics, resulting from the massive usage for diseases control and animal growth promoter (Maizatun et al. 2017). The most common antibiotics associated with slaughterhouse and wet market wastes are doxycycline, chlor-tetracycline, and sulfadiazine (Savin et al. 2020). Discharging antibiotic compounds directly into rivers or leachate directly impacts the living organisms in the water bodies and eventually harm aquatic life, leading to increased distribution of antibiotic-resistant bacteria in the environment (Maizatun et al. 2017).

Table 3 Inedible portions of fish and shellfish as wastes (Michaela Archer 2001)

Type	Species	Inedible portion (%)	Inedible parts
Fish	Catfish	65%	Viscera, frame, skin, fin, and head
	Demersal (general)	57%	
	Haddock	57%	
	Hake	50%	
	Lemon sole	58%	
	Ling	52%	
	Plaice	65%	
	Redfish	70%	
	Whiting	62%	
Shellfish	Crab	68%	Shell and viscera
	Lobster	56%	
	Nephrops	76%	
	Shrimp	65%	
	Crustacea	60%	
	Oyster	61%	
	Cockle	86%	
	Winkle	77%	
	Scallop	86%	
	Mussel	86%	
	Whelk	58%	
	Mollusc	80%	

The wastewater from slaughterhouses and wet markets also contains several toxic materials, such as nitrate, detergents, surfactants, and chloric anions, which are classified as emerging contaminants (Latiffi et al. 2021). Both cationic and anionic surfactants are used in slaughterhouses for cleaning and disinfection purposes but are associated with increased total nitrogen and phosphorus concentrations in wastewater (Apandi 2020). All these factors cause the wastewater to become susceptible to the risk of infectious pathogens. Based on the bibliometric analysis for 500 published papers downloaded from the Scopus database using the keywords of “transmission” AND “slaughterhouse” OR “wet market” AND “pathogens” AND “water” OR “food”, it was noted that the most common pathogenic bacteria associated with slaughterhouse and wet market wastewater include *Salmonella* spp., *Listeria* sp., *E. coli*, *Shigella* sp., and *Staphylococcus* sp. (Fig. 3).

Some common pathogenic bacteria in slaughterhouse and wet market wastes with infection potential have been identified and classified by the World Health Organization. These bacteria can be appropriately eliminated using sanitary methods, such as the filtration method and garbage pails. In some tropical countries, high cases of cholera, typhoid, malaria, dengue fever, and dysentery have been reported due to the increasing number of flies and mosquitoes found in the wastewater of slaughterhouses (Maizatul et al. 2017).

Disease transmission and control

Zoonotic transmission in slaughterhouses or wet markets is a threat to both the food industry and human health (Garcia et al. 2019; Guo et al. 2019). *Campylobacter* spp., *Salmonella* spp., *Yersinia* spp., Shiga toxin-producing *Escherichia coli* (STEC), Hepatitis E virus (HEV) from pigs, avian influenza virus, and other viruses are common pathogens detected in slaughterhouses (Maria 2014). With the recent outbreak of zoonotic diseases emerging from wet markets, such as SARS-associated coronavirus (SARS-CoV) (Mizumoto et al. 2020), this puts slaughterhouses and wet markets under the utmost attention. Hence, the public is concerned with the appropriate waste management and safe disposal regulations for disease prevention. Zoonotic transmission represents 60% of human pathogens from wildlife and livestock (Rahman et al. 2020). Workers in slaughterhouses are a high-risk group to become infected but have less knowledge of the threats of disease transmission (Fredriksson-Ahomaa 2014). A polluted environment at slaughterhouses due to the growth of microbes in the drainage is a big issue (Ketchell 2017). Investments in basic protective equipment, such as masks, gloves, and goggles to prevent direct contact with any discharges from animals, are lacking in many slaughterhouses (Department of Veterinary Services Malaysia 2018). Many abattoirs in rural areas have deplorable infrastructures, such as without proper rooftops and suitable water supply facilities, and animal flesh

Fig. 3 Interactions between SH-biowastes from slaughterhouses and wet markets, and most common bacterial strains in wastes as well as most common antibiotics used

being processed on the floor (Cook et al. 2017). Additionally, even sick animals are processed by some slaughterhouses to increase profit. Adonu et al. (2017) reported that several diseases could spread through sick animals and cause severe health issues for workers who are in direct contact with the animals in the slaughterhouses. This occurred because 70% of slaughterhouses are not properly controlled and inspected (Ketchell 2017).

Some slaughterhouses are combined with wet markets to reduce cost and offer fresh meat sources directly to the customers. These wet markets host vast types of wildlife, poultry, and even seafood, some of them together in stacked cages. Different types of animals are slaughtered at the same place, which escalates the chances of disease transmission (Steven 2020). SAR-CoV2 is an example of novel coronavirus transmission detected in the sewage pipeline of such joint wet markets (Medema et al. 2020). Therefore, in order to increase the knowledge to avoid another pandemic outbreak from meat reservoirs, refining the Meat Control Act (2012) integrated with the “One Health” policy demands immediate attention (Zheng et al. 2019). “One Health” is a program under the World Health Organization (WHO) to design and implement policies, programs, and legislation involving human and animal health in relation to public health, such as preventing a disease outbreak (Martin et al. 2017). The arrangement of

mixed-animal slaughtering in the same slaughterhouse should be banned to avoid cross-contamination of microbes from different livestock to cut the potential of zoonotic transmission.

In addition, enhanced multi-sectional collaboration and data sharing between WHO partners and the Centers for Disease Control and Prevention (CDC) on public health are envisaged to ensure refinements in the prediction of viral emergence, disease severity, and reduction of disease transmission (Eisenstein 2018).

Techniques for effective slaughterhouse SH-biowaste recycling

The current biowaste segregation processes associated with the segregation of liquid and solid wastes are known as feed-stock technologies. Segregation processes are key for reducing the health risk of slaughterhouse and wet market wastes and minimizing the survivability of infectious agents, where the solids in wastewater contribute effectively to the colonization of microbes. After segregation, the solid wastes from slaughterhouses and wet markets are collected and stored in a refuse room not exceeding 24 h to minimize the growth of fungi and bacteria (Noman et al. 2016). The solid wastes

separated during different processes are skin trimmings, meat trimmings, dung, and rumens. Biowaste management technologies have evolved greatly from simple processes to the most efficient and high-end processes, which can be classified into four (4) categories, which are direct use of the biowastes generated, biological processes, physical-chemical processes, and thermo-chemical processes (Lohri et al. 2016). Currently, biological and thermo-chemical processes have gained tremendous attention from industries because these processes could recycle large amounts of biowastes in a short period of time. This paper will further discuss how biological and thermo-chemical processes are used as biowaste recycling technologies.

Biological processes are the conversion of biowastes by living organisms and require less energy consumption. Anaerobic digestion and composting are among the well-known biowaste recycling methods for waste management. In well-developed and modernized slaughterhouses, high-loading anaerobic reactors are used to treat effluents from slaughterhouses using high-rate anaerobic systems (Baker et al. 2020). There are two typical types of anaerobic reactors' feeding modes, which are batch reactor and continuous reactor. The simplest type of reactor is the batch reactor with agitation and heating, or a cooling process under a closed system. On the other hand, a continuous reactor is an open system that allows adding and removing materials or by-products. This type of reactor is well equipped with agitation to provide proper mixing of reactants.

These anaerobic reactors can be operated according to the reactors' temperature conditions, which are mesophilic and thermophilic anaerobic digestion conditions. Mesophilic anaerobic digestion involves bacteria/microbes primarily present at ambient temperatures ranging between 20 and 45 °C. On the other hand, thermophilic anaerobic digestion involves bacteria/microbes that can tolerate high temperatures in the range of 49 to 70°C. The optimum performance of the system depends on the biological aspects of the anaerobic digestion ecosystem (Franke-Whittle and Insam 2013).

The anaerobic digestion technology is being used to utilize the biowastes and simultaneously produce methane as an alternative energy source (Yoon et al. 2014; Sari et al. 2016). The technology has several advantages, such as reducing carbon dioxide emissions, odors, and pathogenic disease transmission. This technology can save land spaces and are applicable to different types of wastes including solid or liquid wastes. Budiyo et al. (2011) reported that 48.89% of the biogas yielded is methane, 47.87% is CO₂, and 2.43% is ammonia during the anaerobic digestion of lipid, carbohydrate, and fatty acid contents at certain temperatures.

The composting technique, on the other hand, is a common method to treat different wastes, such as manure, litter, and feathers from slaughtering activities, and recycle them into valuable bio-products (Salminen and Rintala 2002) (Fig. 4).

Table 4 presents information on the composting processes of wastes from slaughterhouses and wet markets along with their end product's (compost) characteristics. Large-scale poultry slaughterhouse wastes are mixed in different ratios with bulking agents, such as cardboards, agricultural waste, or sewage waste (Asses et al. 2019). Although there are concerns regarding the pathogens available in the compost and its possible infectious disease transmission, Asses et al. (2019) confirmed that the resultant compost is rich in effective microbial communities, while harmful pathogens such as *Streptococci* and *Escherichia coli* are not detected at the end of the process. Lou et al. (2015) also reported the reduction of pathogenic microorganisms in SH-biowaste compost, while the nutrients converted out of the compost can enhance soil fertility. The main products of composting processes are fully mineralized materials, such as CO₂, H₂O, NH₄⁺, stabilized organic matters heavily populated with competitive microbial biomasses, and ash. Based on the facility's requirements and energy consumption, the composting treatment can have a higher impact as compared with other treatments because composting may reduce the weight and volume of slaughterhouse and wet market biowastes dumped into landfills and reduce pathogenic microorganisms in the soil, and the nutrient conversion from the wastes can enhance soil properties.

Table 4 shows that separated biowastes, such as cow manure, have lower physical impurities compared with mixed biowastes (animal feedstock). Thus, mixed biowastes are subject to present higher environmental problems than separated biowastes depending on their characteristics. The physico-chemical biowaste characteristics can be assessed through the American Public Health Association's (APHA) standard method for determining total nitrogen (TN), total phosphorus (TP), total organic carbon (TOC), total dissolved solids (TDS), and carbon-to-nitrogen ratio (C/N) (Table 1). Different characteristics of biowaste require different types of treatment necessary to deactivate pathogens and avoid disease outbreaks and environmental problems that are harmful to both humans and animals. The potential risks of infectivity associated with pathogen-contaminated biowaste tissues and prions have been comprehensively discussed in a previous study (Adhikari et al. 2018).

Thermo-chemical processes for biowaste recycling are widely used and include pyrolysis, liquefaction, incineration, and gasification. Thermal-chemical treatments are processes in which heat is applied to sanitize SH-biowastes, and the primary function of thermal treatments is to convert SH-biowastes into usable end products with a stable state and to reduce the final disposal of SH-biowastes in landfills (Staron et al. 2017). Thermal-chemical treatments are highly recommended by the Canadian Food Inspection Agency (CFIA) for the deactivation of available pathogens in SH-biowastes. Pyrolysis is conducted at high temperatures in low oxygen, while gasification is a thermal treatment technique that uses

Table 4 The composting efficiency for different types of slaughterhouses biowastes

SH-biowaste types	Composting efficiency	Reference
Cattle slaughter specified risk materials (SRM) and mortalities with manures (COWC)	<p>Final compost properties:</p> <ul style="list-style-type: none"> • Temperature: 47°C (SRMC), 44.1°C (COWC) • Moisture: 0.483 ± 0.006 kg/kg (SRMC), 0.460 ± 0.019 kg/kg (COWC) • pH: 8.0 (SRMC and COWC) • Total nitrogen: 15.4 ± 1.6 g/kg (SRMC), 18.3 ± 0.6 g/kg (COWC) <p>A bin composter was used in this research and the temperature increase in the mesophilic phase. Final composting process were recorded</p>	Hao et al. (2008)
Wet market waste	<p>as follows:</p> <ul style="list-style-type: none"> • Moisture content: 53.66% • Density: 510.48 kgm⁻³ • pH: 8.45 • Phosphorus: 189372.7 mg/L • Copper: 51.50 ppb • Zinc: 233.75 ppb • Total organic carbon: 44.75% • Total nitrogen: 3.19 mg/L 	Tweib et al. (2011)
Abattoir waste	<ul style="list-style-type: none"> • Carried out in two windrows comprising mixed organic waste (MOW) and cow intestinal waste (CIW) • Total nitrogen, C:N ratio, total phosphorus, potassium, iron, lead, chromium, nickel, zinc, and cadmium contents were found higher in the MOW pile rather than in the CIW pile due to nutrient contents in abattoir waste 	Hammed et al. (2011)
Broiler slaughter waste and coconut coir pith waste	<ul style="list-style-type: none"> • The composting process was done in two stages in thermophilic temperatures for 52 days • pH, EC, TDS, total organic carbon and total nitrogen, weight, and volume of the waste were reduced significantly ($P < 0.01$) • Calcium, phosphorus, and potassium contents increased, while total bacterial count and coliform were reduced 	Bharathy et al. (2012)
Slaughterhouse solid waste (SSW)	<ul style="list-style-type: none"> • Treatment was carried out in a 60L reactor. The aerobic composting efficiency is as follows: • SSW mass reduction: 55.75% • Average total nitrogen content: 5.17% • Carbon to nitrogen (C:N) ratio: 11.99 • Ammonium sulfate can inactivate <i>Salmonella</i> spp. 	Trihadiningrum et al. (2015)
Cow manure-based mixtures	<ul style="list-style-type: none"> • The initial C:N ratio of 20:1, 30:1 and 40:1 required 4, 5, and 7 days, respectively, for <i>Salmonella</i> inactivation 	Erickson et al. (2009)
Manure	<ul style="list-style-type: none"> • Temperature (>55°C) sanitized the animal wastes efficiently, with a complete destruction of <i>E. coli</i> (1.4×10^3 CFU) and <i>Salmonella</i> sp. (2.6×10^4 CFU) and a significant reduction in the viable counts of <i>Listeria</i> sp. (1.7×10^4 CFU) 	Goberna et al. (2011) Erickson et al. (2015)
Dairy manure	<ul style="list-style-type: none"> • Nitrogen was shown efficient for the inactivation of <i>Salmonella</i>, <i>Listeria</i> spp., and <i>Escherichia coli</i> in animal manure 	Erickson et al. (2014)
Animal waste feedstock	<ul style="list-style-type: none"> • Ammonia inactivated pathogens (<i>Salmonella</i> spp., <i>L. monocytogenes</i>, and <i>E. coli</i>) faster in fresh manure compared with aged manure • Pathogens can be inactivated through heat and microbial competition during the composting process 	Gurtler et al. (2018)

partial air. The emergence of pyrolysis, which works at high temperatures ($>500^{\circ}\text{C}$) in low or absence of oxygen, proposes a great potential in energy recovery. Incineration is the most effective method among all treatments, as it will kill all infectious agents that may present risks to human health and the environment (Tweib et al. 2011). Nevertheless, the downsides of the incineration method are the relatively slow process, higher cost, and stringent waste code criteria and capacity that the process is conducted in one facility (Sengupta 2004). The pyrolysis technique often produces greenhouse gases, such as carbon monoxide and nitrogen. Comparatively, incineration produces carbon dioxide (CO_2), oxygen (O_2), nitrogen (N_2), water, slag, and ash (Igalavithana et al. 2020).

Gasification is one of the thermal treatment techniques for partial oxidation at operating temperatures of 800 to 900°C . Gasification produces composite gas or syngas primarily comprising carbon monoxide, hydrogen, methane, and carbon dioxide, including catalysts. The technology is widely used in slaughterhouse and wet market biowaste treatments, and it releases minimum gases into the atmosphere (Gusiatin et al. 2020). The gases generated from gasification are utilized as sources of energy for heating, transportation, and electrical power (Fig. 4).

The gasification process may release or absorb heat to the surroundings, depending on the system's design, the characteristics of the gas, and the calorific values and production of the by-products. The major benefit of the gasification of slaughterhouse and wet market biowastes is that the gas generated from gasification can be used directly to fuel a gas turbine system, thus improving the gasification thermal

efficiency of the plant. Moreover, Gusiatin et al. (2020) reported that poultry ash, as the by-product of the gasification process, can remove toxic heavy metals, such as cadmium (Cd), from the wastewater. However, the main disadvantage of the gasification process is the higher initial capital investment for the construction and assembly of the thermal treatment equipment (Sari et al. 2016). This technique is associated with high upfront capital for the construction of complex equipment and infrastructure and with high plant maintenance and operation costs. In addition, this technique is restricted to small-scale waste loads, and thus large-scale applications require more complex treatment plants. Competent operators are needed to operate the plant continuously at an optimum condition, and additional energy is required (fuel and electricity).

Advanced biowaste recycling techniques are ideal for waste management, as these will reduce the threats to the public (spread of diseases, foul odors, and olfactory nuisance) and also the environment (air pollution and water and ground-water pollution). Additionally, biowaste recycling processes have evolved greatly due to the increasing demand for biowaste by-products and also due to limited landfills. The development of biowaste by-products can shift the economic paradigm toward a circular economy, which focuses on the recovery of resources and turning them back into valuable by-products for commercialization. Thus, biowaste recycling technologies provide future opportunities for economic growth, as the technologies can act as a driving force for biowaste management business development. The recovery of nutrient sources, such as phosphorus and nitrogen, from biowaste recycling is a good opportunity for waste-to-wealth

Fig. 4 Bio-products and renewable energy generated from SH-biowastes from slaughterhouses and wet markets using different recycling techniques

product development and thus provides financial stability for biowaste businesses (Lohri et al. 2016). The market demand for biowaste recycling technologies will rise, as demand increases for waste products, such as stockfeed and fertilizers from animal biowastes, and also bio-energy as an alternative source of energy (Lohri et al. 2016).

Conclusion

With the rapid increase of the food production industry, health risk aspects associated with slaughterhouses and wet markets should not be neglected. Biowastes produced from slaughterhouses and wet markets are of a broad spectrum and these are the growing grounds for infectious disease. Different variances in slaughterhouse biowaste composition require proper sorting and segregation for efficient waste management and selection of efficient treatment methods. Developed countries have been applying automation in processing and recycling large-scale biowastes from slaughterhouses. This is in huge contrast with the processes in the slaughterhouses of underdeveloped countries, in which the slaughtering is performed in an under-privileged setup, with direct disposal of animal wastes into the land and water bodies. Therefore, the rotting of biowastes on land promotes environmental depletion, along with the growth of infectious diseases. Moreover, the accumulation of these biowastes in water bodies increases the content of nitrates and ammonia, inducing uncontrollable ecological problems, such as algae bloom. Land and river contaminations become more dreadful when pathogens and veterinary pharmaceutical chemicals are disposed through biowastes from slaughterhouses into water bodies.

Among various biowaste management techniques, biological and thermo-chemical processes have gained great consideration from industries due to the large recycling capacity and the short processing time. Moreover, the thermo-chemical technique produces gases that are sources for regenerating energy for heating, transportation, and electrical power. Additionally, the high organic contents of wastes from slaughterhouses and wet markets offer a lot of promising benefits for commercial purposes in the future, such as being used as fertilizers. Hence, biowaste recycling technologies provide future opportunities for biowaste management business development.

Author contribution AAG collected the data and drafted the manuscript. NLM analyzed the data and drafted the manuscript. PFR edited the manuscript. NS collected the biowaste data. MAY collected the data on advanced techniques for recycling and drafted the manuscript. RMSRM collected the data on current technology for waste management. CFS supervised, drafted, and edited the manuscript. All authors read and approved the final manuscript

Funding This research was funded by the Ministry of Higher Education Malaysia under the Research Excellence Consortium Grant Scheme (KKP) or KPM-Special Grant RMK-10 with reference number of JPT(BPKI)1000/016/018/25(54) or KKP Vot. No. K343.

Data availability Not applicable.

Declarations

Ethical approval Not applicable.

Consent for publication Not applicable.

Competing interests The authors declare no competing interests.

References

- Adesiyun IM, Bisi-Johnson M, Aladesanmi OT, Okoh AI, Ogunfowokan AO (2018) Concentrations and human health risk of heavy metals in rivers in Southwest Nigeria. *J Health Pollut* 8:180907
- Adeyemo OK (2002) Unhygienic operation of a city abattoir in South Western Nigeria: environmental implication. *African J Environ Assess Ment Manag* 4:23–28
- Adhikari BB, Chae M, Bressler DC (2018) Utilization of slaughterhouse waste in value-added applications: Recent advances in the development of wood adhesives. *Polymers* 10:176
- Adonu RE, Dzokoto L, Salifu SI (2017) Sanitary and hygiene conditions of slaughterhouses and its effect on the health of residents: a case study of Amasaman slaughterhouse in the Ga west municipality, Ghana. *Food Sci Qual Manage* 65:11–15
- Alam MU, Rahman M, Islam MA, Asaduzzaman M, Sarker S, Rousham E, Unicomb L (2019) Human exposure to antimicrobial resistance from poultry production: assessing hygiene and waste-disposal practices in Bangladesh. *Int J Hyg Environ Health* 222:1068–1076
- Aleksic N, Nesovic A, Sustersic V, Gordic D, Milovanovic D (2020) Slaughterhouse water consumption and wastewater characteristics in the meat processing industry in Serbia. *Desalin Water Treat* 190: 98–112
- Alnahas F, Yeboah P, Flidell L, Abidin AY, Alhareth K (2020) Expired Medication: societal, regulatory and ethical aspects of a wasted opportunity. *Int J Environ Res Public Health* 17:787
- Anon (2002) The Animal By-Products Regulations (EC) no. 1774/2002. European Commission, Brussels
- Apandi N (2020) Integrated growth potential of *Scenedesmus* sp. using public market wastewater via phytoremediation process. *Int J Integr Eng* 12:290–299
- Arvanitoyannis IS, Kassaveti A (2008) Fish industry waste: treatments, environmental impacts, current and potential uses. *Int J Food Sci Technol* 43:726–745
- Arvanitoyannis IS, Ladas D (2008) Meat waste treatment methods and potential uses. *Int J Food Sci Technol* 43:543–559
- Asses N, Farhat W, Hamdi M, Bouallagui H (2019) Large scale composting of poultry slaughterhouse processing waste: microbial removal and agricultural biofertilizer application. *Process Saf Environ Prot* 124:128–136
- Baker BR, Mohamed R, Al-Gheethi A, Aziz HA (2020) Advanced technologies for poultry slaughterhouse wastewater treatment: a systematic review. *J Dispers Sci Technol* 42:880–899
- Bharathy N, Sakthivadivu R, Sivakumar K, Saravanakumar VR (2012) Disposal and utilization of broiler slaughter waste by composting. *Vet World* 5:359–361

- Budiyo IN, Widiya S, Johari S (2011) Study on slaughterhouse wastes potency and characteristic for biogas production. *Int J Waste Resour* 1:4–7
- Bustillo-Lecompte C, Mehrvar M (2017) Slaughterhouse wastewater: treatment, management and resource recovery. *Physico-chemical wastewater treatment and resource recovery*:153–174
- Chang X, Meyer MT, Liu X, Zhao Q, Chen H, Chen J, Qiu Z, Yang L, Cao J, Shu W (2010) Determination of antibiotics in sewage from hospitals, nursery and slaughter house, wastewater treatment plant and source water in Chongqing region of Three Gorge Reservoir in China. *Environ Pollut* 158:1444–1450
- Ching-Velasquez J, Fernández-Lafuente R, Rodrigues RC, Plata V, Rosales-Quintero A, Torrestiana-Sánchez B, Tacias-Pascacio VG (2020) Production and characterization of biodiesel from oil of fish waste by enzymatic catalysis. *Renew Energy* 153:1346–1354
- Cook EAJ, de Glanville WA, Thomas LF, Kariuki S, de Clare Bronsvoot BM, Fèvre EM (2017) Working conditions and public health risks in slaughterhouses in western Kenya. *BMC Public Health* 17:14
- Damaceno FM, Buligon EL, Restrepo JCPS, Chiarello M, Niedzialkoski RK, de Mendonça Costa LA, de Lucas JJ, de Mendonça Costa MSS (2019) Semi-continuous anaerobic co-digestion of flotation sludge from broiler chicken slaughter and sweet potato: nutrients and energy recovery. *Sci Total Environ* 683:773–781
- Department of Veterinary Services Malaysia (2018) Peninsular Malaysia: recorded slaughter of livestock. Department of Veterinary Services Malaysia (DVS). Retrieve on 30 March 2020 from: http://www.dvs.gov.my/dvs/resources/user_1/2019/BP/Perangkaan%20Ternakan%2020182019/_Sem.Malaysia_Sembelihan_Ternakan_pdf
- Diarra S, Sandakabatu D, Perera D, Tabuaciri P, Mohammed U (2015) Growth performance and carcass yield of broiler chickens fed commercial finisher and cassava copra meal-based diets. *Appl Anim Res* 43:352–356
- Eisenstein M (2018) Infection forecasts powered by big data. *Nature* 555(7695):S2–S4
- Erickson MC, Liao J, Maa L, Jiang X, Doyle MP (2009) Inactivation of *Salmonella* spp. in cow manure composts formulated to different initial C: N ratios. *Bioresour Technol* 100:5898–5903
- Erickson MC, Liao J, Jiang X, Doyle MP (2014) Inactivation of pathogens during aerobic composting of fresh and aged dairy manure and different carbon amendments. *J Food Prot* 77:1911–1918
- Erickson MC, Smith C, Jiang X, Flitcroft ID, Doyle MP (2015) Manure source and age affect survival of zoonotic pathogens during aerobic composting at sublethal temperatures. *J Food Prot* 78:302–310
- FAO (2021) Food and Agriculture Organization United Nation, <http://www.fao.org/faostat/en/>. 2018. Accessed 29 Jun 2021
- Fearon J, Mensah SB, Boateng V (2014) Abattoir operations, waste generation and management in the Tamale metropolis: case study of the Tamale. *J Public Health Epidemiol* 6:14–19
- Franke-Whittle IH, Insam H (2013) Treatment alternatives of slaughterhouse wastes, and their effect on the inactivation of different pathogens: a review. *Crit Rev Microbiol* 39:139–151
- Fredriksson-Ahomaa M (2014) Public Health Hazards: A. Biological Hazards. *Meat Inspection and Control in the Slaughterhouse*, 277–333
- Garcia SN, Osburn BI, Cullor JS (2019) A one health perspective on dairy production and dairy food safety. *One Health* 7:100086
- Gendebien A, Ferguson R, Brink J, Horth H, Sullivan M, Davis R, Brunet H, Dalimier F, Landrea B, Krack D, Perot J, Orsi C (2001) Survey of wastes spread on land – Final Report, Study Contract B4-3040/99/110194/MAR/E3, European Commission
- Goberna M, Podmirseg SM, Waldhuber S, Knapp BA, García C, Insama H (2011) Pathogenic bacteria and mineral N in soils following the land spreading of biogas digestates and fresh manure. *Appl Soil Ecol* 49:18–25
- Goldstrand RE (1992) An overview of inedible meat, poultry and fishery by-products. In *Inedible Meat By-Products*, pp 1–17. Springer, Dordrecht
- Guo Z, Li B, Shunlong G (2019) Government regulations and voluntary certifications in food safety in China: a review. *Trends Food Sci Technol* 90:160–165
- Gurtler JB, Doyle MP, Erickson MC, Jiang X, Millner P, Sharma M (2018) Composting to inactivate foodborne pathogens for crop soil application: a review. *J Food Prot* 81:1821–1837
- Gusiatin ZM, Kulikowska D, Klik B (2020) New-generation washing agents in remediation of metal-polluted soils and methods for washing effluent treatment: a review. *Int J Environ Res Public Health* 17: 6220
- Hammed TB, Soyingbe AA, Adewole DO (2011) An Abattoir wastewater management through composting: a case study of Alesinloye waste recycling complex. *Int J Interdiscip Soc Sci* 6:1–19
- Hao X, Stanford K, McAllister TA, Larney FJ, Xu S (2008) Greenhouse gas emissions and final compost properties from co-composting bovine specified risk material and mortalities with manure. *Nutr Cycl Agroecosyst* 83:289–299
- Hileman B (2007) Arsenic in chicken production: a common feed additive adds arsenic to human food and endangers water supplies. *Government and Policy* 85:34–35
- Ho K, Chu LM (2019) Characterization of food waste from different sources in Hong Kong. *J Air Waste Manag Assoc* 69:277–288
- Igalavithana AD, Choi SW, Dissanayake PD, Shang J, Wang CH, Yang X, Kim S, Tsang DC, Lee KB, Ok YS (2020) Gasification biochar from biowaste (food waste and wood waste) for effective CO₂ adsorption. *J Hazard Mater* 391:121147
- Ketchell M (2017) From farm to table: poor hygiene in slaughterhouses in rural Kenya. <https://theconversation.com/from-farm-to-table-poor-hygiene-in-slaughterhouses-in-rural-kenya-71820>
- Latiffi N, Mohamed R, Al-Gheethi A, Tajuddin R (2021) Harvesting of *Scenedesmus* sp. after phycoremediation of meat processing wastewater: optimization of flocculation and chemical analysis of biomass. *J Chem Technol Biotechnol* 96:254–261
- Lee CS, Chang CH, Wen CG, Chang SP (2010) Comprehensive nonpoint source pollution models for a free-range chicken farm in a rural watershed in Taiwan. *Agric Ecosyst Environ* 139:23–32
- Lohri CR, Rajabu HM, Sweeney DJ, Zurbrugg C (2016) Char fuel production in developing countries—a review of urban biowaste carbonization. *Renew Sustain Energy Rev* 59:1514–1530
- Lou Z, Wang M, Zhao Y, Huang R (2015) The contribution of biowaste disposal to odor emission from landfills. *J Air Waste Manag Assoc* 65:479–484
- Maizatul A, Mohamed R, Al-Gheethi A, Hashim M (2017) An overview of the utilisation of microalgae biomass derived from nutrient recycling of wet market wastewater and slaughterhouse wastewater. *Int Aquat Res* 9:177–193
- Maria FA (2014) Public Health Hazards: A. Biological Hazards. *Meat Inspection and Control in the Slaughterhouse*, Thimjos Ninios (ed), pp 277–333
- Martin H, Roberto E, Massimo C, Maurizio A, Rüegg B (2017) Knowledge integration in one health policy formulation, implementation and evaluation, (ed) W.H.O. (WHO)
- Matsumura EM, Mierzwa JC (2008) Water conservation and reuse in poultry processing plant—a case study. *Resour Conserv Recycl* 52:835–842
- Medema G, Heijnen L, Elsinga G, Italiaander R, Brouwer A (2020) Presence of SARS-Coronavirus-2 in sewage. *MedRxiv*
- Mensah-Attipoe J, Toyinbo O (2019) Fungal growth and aerosolization from various conditions and materials. *Fungal Infect*:1–10
- Michaela Archer (2001) Fish waste production in the United Kingdom. *Seafish report number SR537*, The Sea Fish Industry Authority, United Kingdom, November 2001

- Mizumoto K, Kagaya K, Chowell G (2020) Effect of the wet market on the coronavirus disease (COVID-19) transmission dynamics in China, 2019-2020. *Int J Infect Dis* 97:96–101
- NABC (2004) Carcass disposal: a comprehensive review. Report written for the USDA Animal and Plant Health Inspection Service. National Agricultural Biosecurity Centre, Kansas State University, USA., 2004
- Noman EA, Rahman N, Shahadat M, Nagao H, Al-Karkhi A, Al-Gheethi A, Lah T, Omar AK (2016) Supercritical fluid CO₂ technique for destruction of pathogenic fungal spores in solid clinical wastes. *Clean-Soil Air Water* 44:1700–1708
- Oh WY, Lee WS, Lee SS, Khan MA, Ko MS, Yang SH, Kim H, Ha JK (2008) Feed consumption, body weight gain and carcass characteristics of Jeju Native Cattle and its crossbreds fed for short fattening period. *Asian-Australas J Anim Sci* 21:1745–1752
- Rahman M, Sobur M, Islam M, Ievy S, Hossain M, El Zowalaty ME, Rahman A, Ashour HM (2020) Zoonotic diseases: etiology, impact, and control. *Microorganisms* 8:1405
- Ratnawati R, Trihadiningrum Y (2014) Slaughter house solid waste management in Indonesia. *J Biol Res* 19:69–73
- Ravanipour M, Bagherzadeh R, Mahvi AH (2021) Fish and shrimp waste management at household and market in Bushehr. *Iran J Mater Cycles Waste Manag* 23:1394–1403. <https://doi.org/10.1007/s10163-021-01219-2>
- Russ W, Meyer-Pittroff R (2004) Utilizing waste products from the food production and processing industries. *Crit Rev Food Sci Nutr* 44: 57–62
- Salminen E, Rintala J (2002) Anaerobic digestion of organic solid poultry slaughterhouse waste—a review. *Bioresour Technol* 83:13–26
- Sari OF, Ozdemir S, Celebi A (2016) May. Utilization and management of poultry slaughterhouses waste with new methods. In *Eurasia 2016 Waste Management Symposium*
- Savin M, Bierbaum G, Hammerl JA, Heinemann C, Parcina M, Sib E, Voigt A, Kreyenschmidt J (2020) Antibiotic-resistant bacteria and antimicrobial residues in wastewater and process water from German pig slaughterhouses and their receiving municipal wastewater treatment plants. *Sci Total Environ* 727:138788
- Sengupta B (2004) Solid waste management in slaughter house, published by Central Pollution Control Board (CPCB). Ministry of Environment and Forests, Delhi
- Shija DS, Mtenga L, Kimambo A, Laswai G, Mushi D, Mgheni D, Mwilawa A, Shirima E, Safari J (2013) Preliminary evaluation of slaughter value and carcass composition of indigenous sheep and goats from traditional production system in Tanzania. *Asian-Australas J Anim Sci* 26:143–150
- Staron P, Kowalski Z, Staron A, Banach M (2017) Thermal treatment of waste from the meat industry in high scale rotary kiln. *Int J Environ Sci Technol* 14:1157–1168
- Steven LM (2020) China's omnivorous markets are in the eye of a lethal outbreak once again. *The New York Times*. <https://www.nytimes.com/2020/01/25/world/asia/china-markets-coronavirus-sars.html>
- Trihadiningrum Y, Ratnawati R, Wulandari R, Radita DR, Juliastuti S (2015) Comparison of slaughterhouse solid waste treatment using anaerobic-anoxic-oxic and aerobic composting methods. In 5th Environmental Technology and Management Conference (ETMC). Bandung, Indonesia
- Tweib SA, Rahman RA, Khalil MS (2011) composting of solid waste from wet market of Bandar Baru Bangi Malaysia. *Aust J Basic Appl Sci* 5:975–983
- Woodard, Curran, Inc (2006) Chapter 10 - Wastes from Industries (Case Studies), Editor(s): Woodard and Curran. Inc, Industrial Waste Treatment Handbook (Second Edition), Butterworth-Heinemann, pp 409–496. <https://doi.org/10.1016/B978-075067963-3/50012-6>
- Yoon YM, Kim SH, Oh SY, Kim CH (2014) Potential of anaerobic digestion for material recovery and energy production in waste biomass from a poultry slaughterhouse. *J Waste Manag* 34:204–209
- Zheng Z, Lu Y, Short KR, Lu J (2019) One health insights to prevent the next HxNy viral outbreak: learning from the epidemiology of H7N9. *BMC Infect Dis* 19:138

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.