Correction to: Hydrogen sulfide ameliorates aging-associated changes in the kidney

Hak Joo Lee · Denis Feliers · Jeffrey L. Barnes · Sae Oh · Goutam Ghosh Choudhury · Vivian Diaz · Veronica Galvan · Randy Strong · James Nelson · Adam Salmon · Christopher G. Kevil · Balakuntalam S. Kasinath

Published online: 28 January 2021
© American Aging Association 2021

Correction to: GeroScience (2018) 40:163–176
<https://doi.org/10.1007/s11357-018-0018-y>

The original version of this article unfortunately contained an error in Fig. 7a image due to an error in

The online version of the original article can be found at
<https://doi.org/10.1007/s11357-018-0018-y>

H. J. Lee · D. Feliers · J. L. Barnes · S. Oh ·
G. G. Choudhury · B. S. Kasinath (✉)
Department of Medicine, University of Texas Health San Antonio,
7703, Floyd Curl Drive, MC7882, San Antonio, TX 78229, USA
e-mail: kasinath@uthscsa.edu

J. L. Barnes · G. G. Choudhury · V. Galvan · R. Strong ·
A. Salmon · B. S. Kasinath
South Texas Veterans Health Care System, San Antonio, TX,
USA

V. Diaz · V. Galvan · R. Strong · J. Nelson · A. Salmon ·
B. S. Kasinath
Barshop Institute for Longevity and Aging Studies, University of
Texas Health, San Antonio, TX, USA

A. Salmon
Department of Molecular Medicine, University of Texas Health
San Antonio, San Antonio, TX, USA

C. G. Kevil
Louisiana State University Health Science Center, Shreveport,
LA, USA

urinary creatinine estimation. The corrected Fig. 7a and caption are shown below.

Fig. 7 Hydrogen sulfide inhibits age-related increase in albuminuria. Mice, 18–19 months of age, were randomized to receive NaHS in drinking water or water alone (control) and followed for 5 months. (A) Urinary albumin to creatinine ratio (ACR) increased in control mice during the study period which was prevented by NaHS (a) $p < 0.05$, (b) $p < 0.01$, and (c) $p < 0.001$

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.