



Handgrip strength is associated with learning and verbal fluency in older men without dementia: insights from the NHANES

Konstantinos Prokopidis · Panagiotis Giannos · Theocharis Ispoglou · Ben Kirk · Oliver C. Witard · Yannis Dionysiotis · David Scott · Helen Macpherson · Gustavo Duque · Masoud Isanejad

Received: 31 October 2022 / Accepted: 24 November 2022 / Published online: 30 November 2022
© The Author(s) 2022

Abstract Low handgrip strength, a hallmark measure of whole-body strength, has been linked with greater odds of cognitive decline and dementia; however, conflicting findings, which could be due to population characteristics and choice of tools, such for the assessment of handgrip strength and cognitive function domains, also exist. Therefore, we examined the relationship of handgrip strength with a comprehensive list of tests to assess domains of cognitive function using a representative sample of US older

men and women without neurodegenerative disorders such as dementia. We analyzed cross-sectional data from the US National Health and Nutrition Examination Survey (NHANES) between 2011 and 2014, with a study cohort of 777 older adults (380 men and 397 women) above 60 years of age. Handgrip strength was assessed using a handgrip dynamometer, while cognitive function was assessed through the Consortium to Establish a Registry for Alzheimer’s Disease (CERAD) Word List Learning Test (WLLT), Word List Recall Test (WLRT), Intrusion Word Count Test (WLLT-IC and WLRT-IC), the Animal Fluency Test (AFT), and the Digit Symbol Substitution Test (DSST). Sex-stratified multiple linear regression analyses were performed upon covariate adjustment for age, ethnicity, socio-economic status, education, medical history, body mass index, physical activity,

Konstantinos Prokopidis and Panagiotis Giannos have contributed equally to this work and share first authorship.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11357-022-00703-3>.

K. Prokopidis (✉) · M. Isanejad
Department of Musculoskeletal Biology, Institute of Life Course and Medical Sciences, University of Liverpool, Liverpool, UK
e-mail: k.prokopidis@liverpool.ac.uk

K. Prokopidis · P. Giannos
Society of Meta-Research and Biomedical Innovation, London, UK

P. Giannos
Department of Life Sciences, Faculty of Natural Sciences, Imperial College London, London, UK

T. Ispoglou
Carnegie School of Sport, Leeds Beckett University, Leeds, UK

B. Kirk · G. Duque
Australian Institute for Musculoskeletal Science (AIMSS), The University of Melbourne and Western Health, St Albans, VIC 3021, Australia

B. Kirk
Department of Medicine-Western Health, Melbourne Medical School, The University of Melbourne, St Albans, VIC 3021, Australia

O. C. Witard
Faculty of Life Sciences and Medicine, Centre for Human and Applied Physiological Sciences, King’s College London, London, UK

energy, protein, and alcohol intake. Maximal handgrip strength was positively associated with cognitive function scores, including CERAD WLLT ($P=0.009$, $R^2=0.146$) and AFT ($P=0.022$, $R^2=0.024$) in older men, but not in women (CERAD WLLT: $P=0.253$, AFT: $P=0.370$). No significant associations with CERAD WLLRT (men: $P=0.057$, women: $P=0.976$), WLLT-IC (men: $P=0.671$, women: $P=0.869$), WLLRT-IC (men: $P=0.111$, women: $P=0.861$), and DSST (men: $P=0.108$, women: $P=0.091$) were observed. Dose–response curves exhibited a prominent linear relationship between all significant associations after covariate adjustment, with no indication of a plateau in these relationships. In conclusion, higher handgrip strength was independently associated with better learning ability for novel verbal information and verbal fluency in US men over the age of 60 and without dementia. Longitudinal studies are required to confirm whether muscle strength independently predicts cognitive function changes in older adults in a sex-specific manner, and whether this connection is affirmed to the possibility of reverse causation due to declines in physical activity levels in the preclinical phase of dementia.

Keywords Handgrip strength · Cognitive function · Older adults · CERAD · AFT · DSST

Introduction

Aging is associated with cognitive impairments that lead to deficits in attention, executive function, processing speed, and reaction time [1]. These age-related changes in cognitive function are characterized by alternations in brain structure including gray and white matter shrinkage, decreased dendritic

synapses, and neuronal cell death [2]. Interestingly, a concomitant decline in cognitive and physical performance has been observed in older populations as emerging data link low muscle strength with poor cognitive outcomes, including cognitive impairment and dementia [3]. In addition, age-related declines in muscle strength reduce independence and result in poor health outcomes among older adults that are linked to malnutrition, depression, recurrence of falls and fractures, and poor balance [4].

Women appear to be twice as likely to develop dementia than men [5], which is partly due to sex hormone changes after menopause, since hormone therapy is associated with reduced risk of neurodegenerative diseases including dementia [6]. Interestingly, a 12-year longitudinal study revealed a greater number of modifiable risk factors for dementia during middle-to-older age in men compared with women [7]. Loss of physical function may co-occur with losses in cognitive performance as a part of the aging process, but it is not known whether alterations in physical performance correspond with a sex-specific magnitude of cognitive decline. Handgrip strength constitutes a hallmark measure of physical performance and function [8] and it has recently been shown, by using pooled data from general population-based prospective cohort studies, that increased handgrip strength is associated with reduced risk of cognitive impairments such as dementia [3]. Nevertheless, associations between handgrip strength and cognitive outcomes in men and women with good cognitive function tend to be non-significant [3]. Considering that the association between handgrip strength and all-cause mortality is stronger in women than men [9], although the age-related decline in handgrip strength has been speculated to be greater in men compared to women from the age of 30 onwards [10],

Y. Dionyssiotis
Laboratory for Research of the Musculoskeletal System,
National and Kapodistrian University of Athens, Kifissia,
Greece

Y. Dionyssiotis
1st Physical Medicine and Rehabilitation Department,
National Rehabilitation Center EKA, Athens, Greece

D. Scott
Institute for Physical Activity and Nutrition (IPAN),
School of Exercise and Nutrition Sciences, Deakin
University, Burwood, VIC, Australia

D. Scott
Department of Medicine, School of Clinical Sciences
at Monash Health, Monash University, Clayton, VIC,
Australia

H. Macpherson
Institute for Physical Activity and Nutrition, Deakin
University, Geelong, Australia

G. Duque
Research Institute of the McGill University Health Centre,
Department of Medicine, McGill University, Montreal,
QC, Canada

it is important to investigate sex-specific associations between handgrip strength and health-related and/or functional outcomes.

At present, there is a large body of literature reporting that handgrip strength is linked with multiple aspects of cognition across healthy [11–18], community-dwelling [19], and diseased aging populations [20, 21]. Indeed, lower handgrip strength has been linked with increased white matter hyperintensity volume [22, 23] and lower whole-brain volume [24], both of which are associated with cognitive decline and dementia risk [19, 25, 26].

Sex differences in cognitive function are well documented with women typically achieving higher scores in verbal processing tasks while men in visuospatial processing tasks [27]. Similarly, Ling et al. demonstrated significant cross-sectional associations between handgrip strength and mini-mental state examination (MMSE) scores in older women but not men [28]. By contrast, Kim et al. showed that higher handgrip strength at baseline was positively associated with MMSE scores both in older men and women, but handgrip strength could positively predict MMSE scores over time only in women [29]. Research has also shown a link between handgrip strength and MMSE [30], and Clock Drawing Test performance in both older men and women, but language, short-term, and delayed memory scores only in men [31]. However, considering a potential risk of vascular dementia due to hypertension that is linked to micro- and macro-cerebrovascular lesions, these results may not be applicable to those who are normotensive. Moreover, a previous longitudinal study revealed greater odds of cognitive impairment in older women as opposed to men [32], although that was based on prediction risk rather than the evaluation of a linear response between handgrip strength and aspects of cognition. Finally, handgrip strength has been associated with multiple aspects of cognitive performance such as verbal fluency, sustained attention, processing speed, and working memory in people with history or ongoing incidence of cancer [33]. However, whether handgrip strength is linked with these aspects of cognitive function in a sex-specific manner in community-dwelling older individuals without cognitive impairment remains to be elucidated. Further support for potential sex differences is provided by studies which have shown that testosterone is related to greater handgrip strength in

men [34], and free testosterone is associated with significant positive association with processing speed, sustained attention, and working memory in older men above 60 years of age [35]. In addition, previous studies have not controlled for important confounding variables that can influence cognitive function in community-dwelling older adults, such as history of stroke incidence, and arthritis diagnosis [31, 36] or, in some cases, physical activity and nutritional intake [30]. Therefore, there is a need to address these major limitations in order to make concrete conclusions regarding sex-specific differences.

Using a representative national sample of older US adults (≥ 60 years of age) with normal cognitive function from the National Health and Nutrition Examination Survey (NHANES), we comprehensively examined the sex-specific associations between handgrip strength and cognitive function in terms of immediate and delayed learning ability and inhibition for novel verbal information, verbal fluency, sustained attention, processing speed, and working memory.

Methods

Study design and participants

Cross-sectional data was retrieved from participants aged ≥ 60 years with normal cognitive function using the NHANES between 2011 and 2014. NHANES is a project conducted by the Centers for Disease Control and Prevention (CDC) and the National Center for Health Statistics (NCHS) that is comprised of interviews (demographic, socioeconomic, dietary, and medical history-related questions), physical examinations (anthropometrical, medical, and physiological measurements), and biochemical analyses. The NHANES protocol is approved by the NCHS Research Ethics Review Board and all participants provided written informed consent.

Handgrip strength assessment

Handgrip strength was assessed using a handgrip dynamometer (Takei Digital 5401) by a trained examiner that was monitored by field supervisors. A practice trial was utilized to determine if participants understood the procedure, which was followed by a repeated dynamometer squeeze from each hand with

the participant in a seated position. Each hand was tested three times with a 60-s rest between the same hand measurements. Handgrip strength was measured in kilograms (kg) and expressed as the maximum value out of the three handgrip attempts in each hand.

Cognitive assessment

Cognitive function was assessed in terms of immediate and delayed learning ability and inhibition for novel verbal information—Consortium to Establish a Registry for Alzheimer’s Disease (CERAD) Word List Learning Test (WLLT), Word List Recall Test (WLRT), and Intrusion Word Count Test (WLLT-IC and WLRT-IC)—, verbal fluency [Animal Fluency Test (AFT)], and sustained attention, processing speed, and working memory [Digit Symbol Substitution Test (DSST)]. Scoring in the CERAD WLLT, WLLT-IC, WLRT, and WLRT-IC ranged between 0 and 10, in AFT from 1 to 40, and in DSST from 0 to 100. Higher test scores represented better cognitive performance. Participants that did not provide complete data for any of the aforementioned cognitive function tests were excluded from the study. These tests have been previously validated for use in research practice.

Covariates

Age (years), ethnicity (race), socioeconomic status (family income to poverty ratio (FIPR)), education level (school qualification), medical history [stroke incidence and arthritis (osteoarthritis, rheumatoid arthritis, psoriatic arthritis) diagnosis], body mass index (BMI) (kg/m^2), physical activity (minutes), daily energy intake (kcal), protein (g), and alcohol intake (g) were used as covariates. All covariates were potential confounders in the relationship between handgrip strength and cognitive performance.

Age groups consisted of participants aged ≥ 60 years and classified per decade into 60–69, 70–79, and ≥ 80 years of age to distinguish potential changes that may occur in each decade. Ethnic groups were comprised of Mexican American, other Hispanic, non-Hispanic White, non-Hispanic Black, non-Hispanic Asian, and other (multi) races. Social economic status was categorized as low-middle [family income to poverty ratio (FIPR) < 1] and middle-high (FIPR ≥ 1). Education level was defined as no high school degree, at most a high school degree, or a

college degree at minimum. Medical history in terms of loss of cognitive-memory function, stroke incidence, and arthritis diagnosis was categorized as Yes/No responses based on past incidence reported by a doctor or other health professional. Physical activity was expressed as minutes of moderate-intensity recreational activities on a typical week. A physical activity of < 150 min per week was considered as low-moderate and ≥ 150 min as moderate-high. A BMI $< 18 \text{ kg}/\text{m}^2$ was considered as low, $18\text{--}24.9 \text{ kg}/\text{m}^2$ as moderate, and $\geq 25 \text{ kg}/\text{m}^2$ as high. Energy, protein, and alcohol intake were calculated as averages of the 24-h recall and categorized into low, moderate, and high. In men, energy intake < 2000 kcal was considered as low, $2000\text{--}3000$ kcal as moderate, and > 3000 kcal as high. In women, energy intake of < 1600 kcal was considered as low, $1600\text{--}2400$ kcal as moderate, and > 2400 kcal as high. Protein intake $\leq 0.8 \text{ g}/\text{kg}/\text{body}$ weight was considered low, and $> 0.8 \text{ g}/\text{kg}/\text{body}$ as high. In men, alcohol intake $< 15 \text{ g}$ was considered as low, $15\text{--}30 \text{ g}$ as moderate, and $> 30 \text{ g}$ as high. In women, alcohol intake $< 10 \text{ g}$ was considered as low, $10\text{--}20 \text{ g}$ as moderate, and $> 20 \text{ g}$ as high.

Statistical analysis

Multiple linear regression analyses were performed to examine the association between handgrip strength and cognitive function in terms of test-specific cognitive performance by sex with adjustment of all covariates. A restricted cubic spline was employed to model the non-linear and dose–response relationship between handgrip strength and cognitive function using three knots after covariate adjustments. Handgrip strength and cognitive performance were considered as continuous variables, and all covariates were categorical in type. Statistical significance was established as $P < 0.05$ and statistical analysis was performed using IBM SPSS Statistics v28.

Results

Characteristics of study participants

Handgrip strength and cognitive performance data were available for 777 participants (380 men and 397 women). Baseline sociodemographic, anthropometric, nutritional, and medical history characteristics stratified by sex are summarized in Tables S1 and S2.

Most men were between 60 and 69 years of age (55.8%). These were mostly non-Hispanic White (52.1%) and Black (21.7%), of middle-high socioeconomic status (91.6%) with a college degree at minimum (65.8%). Energy intake was mostly lower than the recommended value (50.3%) or just within this range (42.6%). For most of the participants, protein intake was higher than the recommended value (66.8%). Alcohol intake was low (78.4%) and BMI was high (70.0%). Most participants had moderate-high physical activity (85%). Twenty (5.3%) participants had a history of stroke and 134 (35.3%) had an arthritis diagnosis. The average handgrip strength in men was 40.1 (± 0.4) kg.

Most women were between 60 and 69 years of age (60.2%). These were mostly non-Hispanic White (52.4%) and Black (22.2%), of middle-high socioeconomic status (88.9%) with a college degree at minimum (63.7%). Energy intake was mostly lower than the recommended value (48.4%) or just within this range (44.6%). For most of the participants, protein intake was higher than the recommended value (58.7%). Alcohol intake was low (83.8%) and BMI was high (71.5%). Most participants had moderate-high physical activity (84.1%). Seventeen (4.3%) participants had a history of stroke and 210 (52.9%) had an arthritis diagnosis. The average handgrip strength in women was 25.4 (± 0.2) kg.

Handgrip strength and cognitive function

Handgrip strength was positively associated with CERAD WLLT ($P=0.009$) and AFT ($P=0.022$)

in older men but not in women (CERAD WLLT: $P=0.253$, AFT: $P=0.370$). No associations were found between handgrip strength and CERAD WLLRT (men: $P=0.057$, women: $P=0.976$), WLLT-IC (men: $P=0.671$, women: $P=0.869$), WLLRT-IC (men: $P=0.111$, women: $P=0.861$), and DSST (men: $P=0.108$, women: $P=0.091$) (Table 1). Dose–response curves displayed a linear relationship between all significant associations after adjustment for covariates, with no indication of a plateau in these relationships (Fig. 1). Similar analysis of the total cohort showed significant associations with CERAD WLLT (unadjusted P : 0.008; adjusted P : 0.003) and WLLRT (unadjusted P : 0.033), AFT (unadjusted P : 0.004; adjusted P : 0.010), and DSST (adjusted P : 0.025) (Table S3).

Discussion

In older adults aged 60 years and above living in the USA, handgrip strength was significantly positively associated with learning ability for novel verbal information and verbal fluency in older men but not in women, suggesting that future research is needed to unravel the neurobiological and physiological mechanisms that underpin the differences in both handgrip strength and cognitive function declines during aging in a sex-specific manner.

Our findings are in line with previous research showing sex-specific discrepancies in associations between handgrip strength and domains of cognitive function. Indeed, in a recent study in a

Table 1 Sex-stratified multiple linear regression analysis of the association between maximum handgrip strength and cognitive function by sex, upon covariate adjustment for age,

Cognitive tests	Men			Women		
	β	P	R^2	β	P	R^2
CERAD WLLT	0.084	0.009	0.146	0.050	0.253	0.178
CERAD WLLRT	0.030	0.057	0.133	−0.001	0.976	0.146
CERAD WLLT-IC	0.004	0.671	0.047	0.002	0.869	0.052
CERAD WLLRT-IC	−0.009	0.111	0.041	−0.001	0.861	0.056
AFT	0.087	0.022	0.024	0.049	0.370	0.234
DSST	0.157	0.108	0.334	0.273	0.091	0.349

AFT Animal Fluency Test, BMI body mass index, CERAD Consortium to Establish a Registry for Alzheimer’s Disease, DSST Digit Symbol Substitution Test, WLLT Word List Learning Test, WLLRT Word List Recall Test. WLLT-IC Word List Learning Test—Intrusion Word Count, WLLRT-IC Word List Recall Test—Intrusion Word Count




Fig. 1 Dose–response relationship between handgrip strength and cognitive function in male participants. Significant associations with CERAD WLLT (A) and AFT (B) scores were observed. Linear (blue) and spline (red) models were adjusted for age, gender, ethnicity, socio-economic status, educational level, and medical history (stroke incidence and arthritis diagnosis, body mass index, physical activity, alcohol, protein, and energy intake). Abbreviations: AFT Animal Fluency Test, CERAD Consortium to Establish a Registry for Alzheimer’s Disease, WLLT Word List Learning Test

cohort of Chinese hypertensive older participants (~60 years), higher handgrip strength was linked with increased language, short-term, and delayed memory scores in older men (assessed via Digit Span Forward and Backward Test, Verbal Fluency Test, and Verbal Recall Test), but no associations were observed in older women [31]. However, the authors did not control for important variables that may alter sex-specific responses to indices of cognitive function, including physical activity, energy, and nutrient intake. On the other hand, in cancer survivors (breast, prostate, colon, and cervix

cancer) above 60 years of age, handgrip strength was associated with verbal fluency (AFT), sustained attention, processing speed, and working memory (DSST) in both men and women [33]. It is possible that disease states and the degree to which a patient is exposed may overcome the potential sex-specific associations between handgrip strength and cognitive function at an older age, since in our relatively physically active population, associations were observed with a verbal fluency test only in men.

Our results also conflict with prior longitudinal studies on the topic. For instance, in a 7-year cohort of Mexican Americans aged over 65 years, lower handgrip strength was correlated with reductions in MMSE scores, a test which assesses multiple cognitive domains, in both sexes [30]. Likewise, a 4.4-year follow-up also failed to identify a sex-specific relationship between handgrip strength and MMSE scores in both older men and women 72 years of age, while in another 8-year longitudinal cohort of Korean adults aged 65 years or older, correlations between handgrip strength and cognitive function (via K-MMSE score) were observed in both sexes [37]. Moreover, no sex-specific differences have been found between men and women in the oldest old. Specifically, in a 4-year longitudinal cohort of older Dutch adults aged 89 years, reduced handgrip strength was associated with lower MMSE scores in both men and women [28], while another 1-year longitudinal sample of older Italian men and women 80 years of age showed a significant association between handgrip strength and Clock Drawing Test (CDT) in both sexes [38]. The aforementioned findings from longitudinal studies reveal that studying the relationship of handgrip strength with cognitive domains such as MMSE and CDT infers identical results in both older men and women. However, considering that MMSE is a widely used instrument relying on socioeconomic and educational characteristics as well as a mixture of cognitive aspects [39], the potential sex-specific differences may be more relevant to specific cognitive domains rather than composite scores of multiple tests [36]. Our findings also suggest that strength may be more relevant to cognitive domains (i.e., verbal processing tasks) that men typically perform poorer in than women [27].

Based on the current scarcity of the literature, the mechanisms contributing to differential sex-specific

associations between handgrip strength and distinct domains of cognitive function among observational studies are unclear. In a 16-year longitudinal British cohort of adults aged 69–71 years, a stronger association between handgrip strength and higher whole-brain volume at follow-up was observed in women compared to men [12]. Interestingly, bioavailable testosterone may be a link mediating this relationship in men, considering its associations with indices of cognitive function highlighted in this study [35]. Nevertheless, further insights on how handgrip strength may alter brain physiology under conditions of sexual dimorphism and how that translates to sex-specific alterations in cognitive function are warranted. However, sampling bias should also be taken into account in relation to the aforementioned findings pertaining to sex-specific differences. Considering the cross-sectional data collection and the lack of gold-standard assessment tools for overall cognitive function, it is worth stating that one-time measurement of exposure and outcome does not support causal relationships [40]. Ignoring sampling variance may lead to misinterpretation of underlying biological pathways that we have previously described in the above sections [41]. For example, sampling error due to convenience sampling referring to variations in participant willingness to take part in some of the assessments tests in such studies with large sample sizes may partially explain our results [42, 43]. Given the limited mechanistic evidence that could explain the sex-specific differences found in handgrip strength and its association with indices of cognitive function, these are plausible factors that should be considered.

Strengths and limitations

Our study employed nationally representative data that has undergone rigorous quality control, and multiple confounders were adjusted to accurately estimate the association between handgrip strength and several domains of cognitive function in older men and women without dementia. However, this study was also prone to limitations. Observational studies using data from cross-sectional surveys are unable to exhibit a causal relationship between dependent and independent variables. In addition, NHANES uses specific tests of cognitive performance that may not fully represent overall cognitive function. Cognitive function is comprised of multiple interconnected mental processes and hence, more refined assessment

tools are required. For instance, backward number recall involves greater attention and executive function reserves compared to forward recall testing [44]. In addition, data availability regarding handgrip strength and cognitive performance from NHANES was limited to only two annual cycles, which could amplify sampling bias. Finally, poor physical capacity may rather be a consequence of cognitive decline during aging rather than its cause. This may be of particular importance in longitudinal studies that have not appropriately controlled for physical activity throughout the longitudinal period. The likelihood of reverse causality should also not be dismissed since physical activity levels are reduced before the clinical diagnosis of dementia [45] while previous research has also linked higher white matter hyperintensities with lower physical performance in older populations [46].

Conclusions

Handgrip strength is positively associated with learning ability for novel verbal information and verbal fluency in US older men aged 60 years and above, but not in older women. There is also scope for future studies to confirm the role of strength as a risk factor for decline in specific domains of cognitive function, and specifically in domains where men typically perform worse than women; this can then follow with promotion of appropriate interventions for maintenance and enhancement of strength as a potential means for attenuation of age-related cognitive decline. However, our findings should be interpreted with caution due to their cross-sectional nature. Future studies are warranted to uncover the biological mechanisms to confirm the directionality and explore the potential sex-specific relationship between handgrip strength and components of cognitive function in older men and women, while exercise interventions could unravel potential links involving these factors.

Author contribution Study concept and design: KP and PG. Acquisition of data: KP and PG. Analysis and interpretation of data: KP and PG. Drafting of the manuscript: KP and PG. Statistical analysis: KP and PG. Obtained funding: NA. Administrative, technical, or material support: KP and PG. Study supervision: TI, BK, OW, YD, DS, HM, GD, MI. Critical revision of the manuscript for important intellectual content: TI, BK, OW, YD, DS, HM, GD, MI.

Declarations

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Peters R. Ageing and the brain. *Postgrad Med J*. 2006;82:84–8. <https://doi.org/10.1136/pgmj.2005.036665>.
- Morrison JH, Baxter MG. The ageing cortical synapse: hallmarks and implications for cognitive decline. *Nat Rev Neurosci*. 2012;13:240–50. <https://doi.org/10.1038/nrn3200>.
- Kunutsor SK, Isozoz NM, Voutilainen A, Laukkanen JA. Handgrip strength and risk of cognitive outcomes: new prospective study and meta-analysis of 16 observational cohort studies. *Geroscience*. 2022;44:2007–24. <https://doi.org/10.1007/s11357-022-00514-6>.
- Kirk B, Zanker J, Bani Hassan E, Bird S, Brennan-Olsen S, Duque G. Sarcopenia Definitions and Outcomes Consortium (SDOC) criteria are strongly associated with malnutrition, depression, falls, and fractures in high-risk older persons. *J Am Med Dir Assoc*. 2021;22:741–5. <https://doi.org/10.1016/j.jamda.2020.06.050>.
- Niu H, Alvarez-Alvarez I, Guillen-Grima F, Aguinaga-Ontoso I. Prevalence and incidence of Alzheimer's disease in Europe: a meta-analysis. *Neurologia*. 2017;32:523–32. <https://doi.org/10.1016/j.nrl.2016.02.016>.
- Kim YJ, Soto M, Branigan GL, Rodgers K, Brinton RD. Association between menopausal hormone therapy and risk of neurodegenerative diseases: implications for precision hormone therapy. *Alzheimers Dement: Transl Res Clin Interv*. 2021;7:e12174. <https://doi.org/10.1002/trc2.12174>.
- Anstey KJ, Peters R, Mortby ME, Kiely KM, Eramudugolla R, Cherbuin N, et al. Association of sex differences in dementia risk factors with sex differences in memory decline in a population-based cohort spanning 20–76 years. *Sci Rep*. 2021;11:7710. <https://doi.org/10.1038/s41598-021-86397-7>.
- Adamo DE, Anderson T, Koochaki M, Fritz NE. Declines in grip strength may indicate early changes in cognition in healthy middle-aged adults. *Plos One*. 2020;15:e0232021. <https://doi.org/10.1371/journal.pone.0232021> (ARTN e0232021).
- Arvandi M, Strasser B, Meisinger C, Volaklis K, Gothe RM, Siebert U, et al. Gender differences in the association between grip strength and mortality in older adults: results from the KORA-age study. *Bmc Geriatrics*. 2016;16:1–8. <https://doi.org/10.1186/s12877-016-0381-4> (ARTN 201).
- Vianna LC, Oliveira RB, Araujo CG. Age-related decline in handgrip strength differs according to gender. *J Strength Cond Res*. 2007;21:1310–4. <https://doi.org/10.1519/R-23156.1>.
- Stijntjes M, Aartsen MJ, Taekema DG, Gussekloo J, Huisman M, Meskers CGM, et al. Temporal relationship between cognitive and physical performance in middle-aged to oldest old people. *J Gerontol A Biol Sci Med Sci*. 2017;72:662–8. <https://doi.org/10.1093/gerona/glw133>.
- Dercon Q, Nicholas JM, James SN, Schott JM, Richards M. Grip strength from midlife as an indicator of later-life brain health and cognition: evidence from a British birth cohort. *BMC Geriatr*. 2021;21:475. <https://doi.org/10.1186/s12877-021-02411-7>.
- Chou MY, Nishita Y, Nakagawa T, Tange C, Tomida M, Shimokata H, et al. Role of gait speed and grip strength in predicting 10-year cognitive decline among community-dwelling older people. *BMC Geriatr*. 2019;19:186. <https://doi.org/10.1186/s12877-019-1199-7>.
- Björk MP, Johansson B, Hassing LB. I forgot when I lost my grip—strong associations between cognition and grip strength in level of performance and change across time in relation to impending death. *Neurobiol Aging*. 2016;38:68–72.
- Sternang O, Reynolds CA, Finkel D, Ernsth-Bravell M, Pedersen NL, Dahl Aslan AK. Grip strength and cognitive abilities: associations in old age. *J Gerontol B Psychol Sci Soc Sci*. 2016;71:841–8. <https://doi.org/10.1093/geronb/gbv017>.
- Szlejf C, Suemoto CK, Lotufo PA, Bensenor IM. Association of sarcopenia with performance on multiple cognitive domains: results from the ELSA-Brasil Study. *J Gerontol A Biol Sci Med Sci*. 2019;74:1805–11. <https://doi.org/10.1093/gerona/glz118>.
- Huang CY, Hwang AC, Liu LK, Lee WJ, Chen LY, Peng LN, et al. Association of dynapenia, sarcopenia, and cognitive impairment among community-dwelling older Taiwanese. *Rejuvenation Res*. 2016;19:71–8. <https://doi.org/10.1089/rej.2015.1710>.
- Vaz-Patto M, Bueno B, Ribeiro O, Teixeira L, Afonso RM. Association between handgrip strength, walking, age-related illnesses and cognitive status in a sample of Portuguese centenarians. *Eur Rev Aging Phys Act*. 2017;14:9. <https://doi.org/10.1186/s11556-017-0178-2>.
- Chen KK, Lee SY, Pang BWJ, Lau LK, Jabbar KA, Seah WT, et al. Associations of low handgrip strength and hand laterality with cognitive function and functional mobility - the Yishun Study. *BMC Geriatr*. 2022;22:677. <https://doi.org/10.1186/s12877-022-03363-2>.

20. Huang HH, Chang JC, Liu HC, Yang ZY, Yang YJ, Chen LK, et al. Handgrip strength, tumor necrosis factor- α , interleukin-6, and visfatin levels in oldest elderly patients with cognitive impairment. *Exp Gerontol*. 2020;142:111138. <https://doi.org/10.1016/j.exger.2020.111138>.
21. Su H, Sun X, Li F, Guo Q. Association between handgrip strength and cognition in a Chinese population with Alzheimer's disease and mild cognitive impairment. *BMC Geriatr*. 2021;21:459. <https://doi.org/10.1186/s12877-021-02383-8>.
22. Firth JA, Smith L, Sarris J, Vancampfort D, Schuch F, Carvalho AF, et al. Handgrip strength is associated with hippocampal volume and white matter hyperintensities in major depression and healthy controls: a UK Biobank study. *Psychosom Med*. 2020;82:39–46. <https://doi.org/10.1097/Psy.0000000000000753>.
23. Kilgour AHM, Todd OM, Starr JM. A systematic review of the evidence that brain structure is related to muscle structure and their relationship to brain and muscle function in humans over the lifecourse. *Bmc Geriatrics*. 2014;14:1–35. <https://doi.org/10.1186/1471-2318-14-85> (Artn 85).
24. Camargo EC, Weinstein G, Beiser AS, Tan ZS, DeCarli C, Kelly-Hayes M, et al. Association of physical function with clinical and subclinical brain disease: the Framingham Offspring Study. *J Alzheimers Dis*. 2016;53:1597–608. <https://doi.org/10.3233/JAD-160229>.
25. Vermeer SE, Prins ND, den Heijer T, Hofman A, Koudstaal PJ, Breteler MM. Silent brain infarcts and the risk of dementia and cognitive decline. *N Engl J Med*. 2003;348:1215–22. <https://doi.org/10.1056/NEJMoa022066>.
26. Debette S, Markus HS. The clinical importance of white matter hyperintensities on brain magnetic resonance imaging: systematic review and meta-analysis. *BMJ*. 2010;341:c3666. <https://doi.org/10.1136/bmj.c3666>.
27. Hyde JS. Sex and cognition: gender and cognitive functions. *Curr Opin Neurobiol*. 2016;38:53–6. <https://doi.org/10.1016/j.conb.2016.02.007>.
28. Ling CHY, Gussekloo J, Trompet S, Meskers CGM, Maier AB. Clinical determinants of low handgrip strength and its decline in the oldest old: the Leiden 85-plus Study. *Aging Clin Exp Res*. 2021;33:1307–13. <https://doi.org/10.1007/s40520-020-01639-4>.
29. Kim KH, Park SK, Lee DR, Lee J. The relationship between handgrip strength and cognitive function in elderly Koreans over 8 years: a prospective population-based study using Korean Longitudinal Study of Ageing. *Korean J Fam Med*. 2019;40:9–15. <https://doi.org/10.4082/kjfm.17.0074>.
30. Alfaro-Acha A, Al Snih S, Raji MA, Kuo YF, Markides KS, Ottenbacher KJ. Handgrip strength and cognitive decline in older Mexican Americans. *J Gerontol A Biol Sci Med Sci*. 2006;61:859–65. <https://doi.org/10.1093/gerona/61.8.859>.
31. Lu S, Herold F, Zhang Y, Lei Y, Kramer AF, Jiao C, et al. Higher handgrip strength is linked to better cognitive performance in Chinese adults with hypertension. *Brain Sci*. 2021;11. <https://doi.org/10.3390/brainsci11080985>.
32. McGrath R, Robinson-Lane SG, Cook S, Clark BC, Herrmann S, O'Connor ML, et al. Handgrip strength is associated with poorer cognitive functioning in aging Americans. *J Alzheimers Dis*. 2019;70:1187–96. <https://doi.org/10.3233/JAD-190042>.
33. Yang L, Koyanagi A, Smith L, Hu L, Colditz GA, Toriola AT, et al. Hand grip strength and cognitive function among elderly cancer survivors. *PLoS ONE*. 2018;13:e0197909. <https://doi.org/10.1371/journal.pone.0197909>.
34. Chiu HT, Shih MT, Chen WL. Examining the association between grip strength and testosterone. *Aging Male*. 2020;23:915–22. <https://doi.org/10.1080/13685538.2019.1632282>.
35. Giannos P, Prokopidis K, Church DD, Kirk B, Morgan PT, Ni Lochlainn M, et al. Associations of bioavailable serum testosterone with cognitive function in older men: results from the National Health and Nutrition Examination Survey. *J Gerontol A Biol Sci Med Sci*. 2022. <https://doi.org/10.1093/gerona/glac162>.
36. Lee S, Oh JW, Son NH, Chung W. Association between handgrip strength and cognitive function in older adults: Korean Longitudinal Study of Aging (2006–2018). *Int J Environ Res Public Health*. 2022;19. <https://doi.org/10.3390/ijerph19031048>.
37. Kim H, Kim SH, Jeong W, Jang SI, Park EC, Kim Y. Association between change in handgrip strength and cognitive function in Korean adults: a longitudinal panel study. *BMC Geriatr*. 2021;21:671. <https://doi.org/10.1186/s12877-021-02610-2>.
38. Viscogliosi G, Di Bernardo MG, Ettorre E, Chiriack IM. Handgrip strength predicts longitudinal changes in clock drawing test performance. An observational study in a sample of older non-demented adults. *J Nutr Health Aging*. 2017;21:593–6. <https://doi.org/10.1007/s12603-016-0816-9>.
39. Sczufca M, Almeida OP, Vallada HP, Tasse WA, Menezes PR. Limitations of the Mini-Mental State Examination for screening dementia in a community with low socioeconomic status: results from the Sao Paulo Ageing & Health Study. *Eur Arch Psychiatry Clin Neurosci*. 2009;259:8–15. <https://doi.org/10.1007/s00406-008-0827-6>.
40. Setia MS. Methodology series module 3: cross-sectional studies. *Indian J Dermatol*. 2016;61:261–4. <https://doi.org/10.4103/0019-5154.182410>.
41. Santin-Janin H, Huguely B, Aubry P, Fouchet D, Gimenez O, Pontier D. Accounting for sampling error when inferring population synchrony from time-series data: a Bayesian state-space modelling approach with applications. *PLoS ONE*. 2014;9:e87084. <https://doi.org/10.1371/journal.pone.0087084>.
42. Wang X, Cheng Z. Cross-sectional studies: strengths, weaknesses, and recommendations. *Chest*. 2020;158:S65–71. <https://doi.org/10.1016/j.chest.2020.03.012>.
43. Tyrer S, Heyman B. Sampling in epidemiological research: issues, hazards and pitfalls. *BJPsych Bull*. 2016;40:57–60. <https://doi.org/10.1192/pb.bp.114.050203>.
44. Sun X, Zhang X, Chen X, Zhang P, Bao M, Zhang D, et al. Age-dependent brain activation during forward and backward digit recall revealed by fMRI. *Neuroimage*. 2005;26:36–47. <https://doi.org/10.1016/j.neuroimage.2005.01.022>.
45. Sabia S, Dugravot A, Dartigues J-F, Abell J, Elbaz A, Kivimäki M, et al. Physical activity, cognitive decline, and

risk of dementia: 28 year follow-up of Whitehall II cohort study. *BMJ*. 2017;357:j2709. <https://doi.org/10.1136/bmj.j2709>.

46. Nadkarni NK, Studenski SA, Perera S, Rosano C, Aizenstein HJ, Brach JS, et al. White matter hyperintensities, exercise, and improvement in gait speed: does type of gait rehabilitation matter? *J Am Geriatr Soc*. 2013;61:686–93. <https://doi.org/10.1111/jgs.12211>.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.