



Correction to: Cannabimimetic activities of cumyl carboxamide-type synthetic cannabinoids

Akiko Asada¹  · Takahiro Doi¹ · Takaomi Tagami¹ · Akihiro Takeda¹ · Yuka Satsuki¹ · Masami Kawaguchi¹ · Akihiko Nakamura¹ · Yoshiyuki Sawabe¹

Published online: 21 July 2020
© Japanese Association of Forensic Toxicology 2020

Correction to: Forensic Toxicol (2018) 36:170–177
<https://doi.org/10.1007/s11419-017-0374-9>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

There were some mistakes in Table 1 (the values of EC₅₀ of compounds 5, 6, and 7). The correct Table 1 is placed below.

Table 1 Half maximal (50%) effective concentration (EC₅₀) values of cumyl carboxamide-type cannabinoid compounds

Entry	Compound	EC ₅₀ (mol/L)			
		CB ₁	CB ₂	CB ₁ /CB ₂	CB ₂ /CB ₁
1	CUMYL-PINACA	5.12×10^{-9}	4.75×10^{-8}	0.11	9.28
2	CUMYL-5F-PINACA	1.51×10^{-8}	3.48×10^{-8}	0.43	2.30
3	CUMYL-PICA	7.77×10^{-7}	4.44×10^{-7}	1.75	0.57
4	CUMYL-5F-PICA	2.09×10^{-8}	9.09×10^{-6}	<0.01	435
5	CUMYL-THPINACA	8.14×10^{-10}	1.33×10^{-9}	0.61	1.63
6	CUMYL-BICA	1.24×10^{-9}	3.95×10^{-10}	3.14	0.32
7	CUMYL-5F-P7AICA	1.21×10^{-9}	2.48×10^{-10}	4.88	0.20
8	CP 55940 (positive control)	7.35×10^{-10}	4.69×10^{-10}	1.57	0.64

The original article can be found online at <https://doi.org/10.1007/s11419-017-0374-9>.

✉ Akiko Asada
asada@iph.osaka.jp

¹ Osaka Institute of Public Health, 1-3-69 Nakamichi, Higashinari-ku, Osaka 537-0025, Japan