

The hypoxia signaling pathway and hypoxic adaptation in fishes

XIAO Wuhan

Key Laboratory of Aquatic Biodiversity and Conservation, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430072, China

Received September 18, 2014; accepted November 6, 2014; published online January 15, 2015

The hypoxia signaling pathway is an evolutionarily conserved cellular signaling pathway present in animals ranging from *Caenorhabditis elegans* to mammals. The pathway is crucial for oxygen homeostasis maintenance. Hypoxia-inducible factors (HIF-1 α and HIF-2 α) are master regulators in the hypoxia signaling pathway. Oxygen concentrations vary a lot in the aquatic environment. To deal with this, fishes have adapted and developed varying strategies for living in hypoxic conditions. Investigations into the strategies and mechanisms of hypoxia adaptation in fishes will allow us to understand fish speciation and breed hypoxia-tolerant fish species/strains. This review summarizes the process of the hypoxia signaling pathway and its regulation, as well as the mechanism of hypoxia adaptation in fishes.

hypoxia, hypoxia-inducible factors, PHDs, pVHL, FIH, fish

Citation: Xiao WH. The hypoxia signaling pathway and hypoxic adaptation in fishes. *Sci China Life Sci*, 2015, 58: 148–155, doi: 10.1007/s11427-015-4801-z

Approximately 2.5 billion years ago, photosynthesis led to the accumulation of oxygen to levels that were likely toxic to many obligate anaerobes. However, increased availability of atmospheric O₂ led to the evolution of an extraordinarily efficient system of oxidative phosphorylation. In this system, chemical energy stored in the carbon bonds of organic molecules is transferred to the high-energy phosphate bond in ATP, which is then used to power physicochemical reactions in living cells [1]. Additionally, O₂ serves as the final electron acceptor in oxidative phosphorylation, which is not only required for energy production, but is also the direct substrate of many enzymes. Thus, it is critical for the growth, development, and reproduction of organisms. Consequently, metazoans have evolved complicated systems of cellular metabolism and physiology to maintain oxygen homeostasis and have developed a biochemical response to low oxygen levels [2]. There are a number of oxygen-sensing pathways that promote hypoxia tolerance by activating transcription and inhibiting translation: the energy and nutrient sensor mTOR, the unfolded protein re-

sponse that activates the endoplasmic stress response, and the nuclear factor (NF)- κ B transcriptional response [3]. However, hypoxia-inducible factors (HIFs) are recognized as master regulators of the cellular response to hypoxic stress [4,5].

The hypoxia signaling pathway is evolutionarily conserved from *Caenorhabditis elegans* to human beings and it activates similar or homogenous gene expression, resulting in similar physical and biochemical responses. Compared with the terrestrial environment, oxygen concentrations vary greatly in the aquatic environment [6]. Thus, compared with most birds and mammals, fishes are tolerant of this varying oxygen availability. Natural selection by oxygen concentration has facilitated the evolution of fishes with a range of adaptations to variable oxygen concentration. Even in waters at the same latitude, closely related species or different strains within a species exhibit varied adaptations to oxygen concentration. Additionally, closely related fishes distributed in waters at different latitudes exhibit extensive variation in their tolerance of hypoxia. Determining the mechanisms of hypoxia adaptation in fishes will not only help us to un-

email: w-xiao@ihb.ac.cn

derstand fish speciation and the evolution of the hypoxia signaling pathway, but will also guide us in the breeding of hypoxia-tolerant fish species/strains.

1 HIF and the regulation of the hypoxia signaling pathway

HIF is a master regulator in the hypoxia signaling pathway and is expressed by all extant metazoan species analyzed to date. HIF is a heterodimer comprising an oxygen-labile α -subunit (HIF- α) and a constitutively expressed β -subunit (HIF- β or ARNT). Each unit contains the basic helix-loop-helix-PAS (bHLH-PAS) domains, which mediate the formation of heterodimers and DNA binding. HIF- β dimerizes with other bHLH-PAS proteins and is stably expressed, but HIF- α determines HIF-1 transcriptional activity [1,4,7].

Under normoxia (normal oxygen tension), either HIF-1 α or HIF-2 α is hydroxylated on specific conserved proline residues by prolyl-hydroxylase domains (PHDs), which contain enzymes (including PHD1, PHD2, and PHD3; PHD2 is thought to be essential), using molecular oxygen as a substrate [8]. In the reaction, one oxygen atom is inserted into the prolyl residue; a second atom is inserted into the co-substrate α -ketoglutarate, splitting it into CO₂ and succinate. Hydroxy-HIF- α is recognized by von Hippel-Lindau tumor suppressor protein (pVHL) and is subsequently ubiquitinated by the VBC ubiquitin-ligase complex, marking HIF- α for degradation by the 26S proteasome [9]. However, under hypoxic conditions, PHD catalytic activity is inhibited by the lack of oxygen and HIF- α is not hydroxylated, thus, HIF- α is not recognized by the VBC complex, allowing HIF- α to stabilize. Stable HIF- α is transferred into the nucleus and heterodimerizes with stable HIF-1 β . HIF heterodimers recognize and bind to hypoxia response elements (HREs) in the genome with the consensus sequence G/ACGTG to regulate the transcription of genes together with co-activators such as CBP/p300, resulting in a series of physical and biochemical responses (Figure 1) [1,4].

Because HIF- α plays such an important role in the hypoxia signaling pathway, HIF- α modulation is assumed to be a major mechanism for the regulation of the hypoxia signaling pathway. HIF- α modulation includes transcription regulation and post translation modulation (PTM). At present, there are few reports on HIF- α transcription regulation. Those available mainly focus on post translation HIF- α modulation. NF- κ B activates HIF-1 α expression, linking innate immunity to the hypoxic response [10]. HIF- α PTM is exhibited in many pathways. Apart from VHL-mediated HIF- α proteasomal degradation, HIF- α is modulated by acetylation/deacetylation, phosphorylation/de-phosphorylation [11], sumoylation, and neddylation [12]. Histone acetyltransferase p300/CBP enhances HIF-1 α transcriptional activity by interacting with the HIF-1 α C-terminus [13].




Figure 1 Hypoxia signaling pathway.

NAD⁺ dependent-deacetylase, Sirt1, Sirt3, Sirt6, and Sirt7 regulate HIF- α activity either positively or negatively, playing important roles in cell metabolism, life span, tumorigenesis, and cardiovascular disease [14–21]. As reported by Shao et al. [22] hypoxia stimulates an increase in mRNA and SUMO-1 protein levels. SUMO-1 co-localize with HIF-1 α in the nucleus to induce HIF-1 α sumoylation, resulting in the stabilization and enhancement of HIF-1 α transcriptional activity [22–24].

Furthermore, even though some HIF-1 α and HIF-2 α interacting proteins cannot modify HIF-1 α and HIF-2 α , they can regulate their stability and transcriptional activity either positively or negatively via specific mechanisms. PKM2 enhances HIF-1 α transcriptional activity through recruiting p300; PHD3 enhances PKM2 activity as a co-activator by hydroxylating PKM2 [25]. SHARP1 promotes HIF- α degradation by serving as the HIF-presenting factor to the proteasome [26]. EAF2 is a specific HIF-1 α downstream target, which interacts with HIF-1 α but not HIF-2 α , resulting in the inhibition of HIF-1 α transcriptional activity via preventing p300 from binding to HIF-1 α [27]. Other proteins do not interact with HIF- α directly, but interact with other factors that regulate HIF- α , e.g., pVHL and PHDs among others. These proteins indirectly regulate HIF- α activity. E3 ligase Siah2 affects cellular HIF-1 α protein levels and mediates the hypoxic response through regulating PHD protein stability [28]. E2-EPF UCP targets pVHL for degradation to control the HIF-1 α stability [29]. PML inhibits HIF-1 α translation through mTOR repression [30]. SENP3 is responsible for the enhancement of HIF-1 transactivation under mild oxidative stress via de-sumoylation [31]. The LIMD1 protein bridges an association between the PHDs and VHL to inhibit HIF-1 activity [32].

In summary, as a master regulator of the hypoxia signaling pathway, the direct or indirect regulation of HIF- α plays

key roles in the pathway. Moreover, HIF- α regulation is currently an area of intense study [7].

2 Hypoxic adaptation strategies in fish

Oxygen concentration varies greatly in aquatic environments depending on the geographic location, season, composition, and water flow. Fishes have evolved different adaptive strategies with the 20,000 extant species varying greatly in hypoxia tolerance. Fishes that depend on aerobic metabolism for rapid swimming, such as salmon and tuna, are moderately to extremely sensitive to hypoxia, whereas carp, eels, and hagfish can live comfortably in hypoxic conditions [6]. Hypoxic adaptation in fishes can be attributed to long-term adaptation and the acute response.

2.1 Fish adaptation strategies to long-term hypoxia

To adapt to varying oxygen levels in the long-term, fishes usually undergo gene mutations that induce changes in physical and biochemical responses, and tissue and organ structure, resulting in three basic capabilities: (a) to decrease metabolic rate under hypoxia, (b) tolerance to increased levels of metabolic by-products, and (c) to avoid and/or repair cellular injury during re-oxygenation [6].

The first long-term hypoxia adaptation strategy in fishes is either to modify respiratory organs or develop accessory organs in addition to gills. For example, loaches have developed the ability to breathe through the skin and intestine, and eels through the mouth and epithelia of cavum larynges.

The second strategy is to modify the metabolic pathway. The crucian carp (*Carassius carassius*) is the greatest extreme hypoxia tolerant species, enduring months of hypoxia at low temperatures [33]. Their cousins, common goldfish, have half-lethal times of 45 h under anoxia at 5°C and 22 h at 20°C. Under hypoxia, strong metabolic depression is key to the survival of these *Carassius* species [34–36]. Under anoxia, the crucian carp can metabolize anaerobically and convert lactate to ethanol and CO₂. These end products can be excreted into the water through the gills. Unlike the dormant turtles, the crucian carp remains active during both hypoxia and anoxia [35]. The main reason is that crucian carp can take advantage of large muscle and liver glycogen reserves, reduced metabolism, and avoidance of lactic acidosis by converting lactate to ethanol and CO₂ [34,37].

Yet another strategy is to increase the number of red blood cells and promote the oxygen-carrying capabilities of hemoglobin [38]. During their evolution, fishes have been undergone genetic segregation many times, resulting in huge diversity in the structure and function of genes related to hypoxia adaptation. In carp, HIF- α is duplicated. Even though one copy retains the ancient hypoxia response, a second copy increases hypoxia-sensitivity. This phenomenon indicates that HIF- α in carp has differentiated function-

ally during speciation [39]. Furthermore, Chi et al. [40] found different evolutionary patterns of HIF- α isoforms in the basal branches of the Actinopterygii and Sarcopterygii.

2.2 The acute hypoxia stress response in fishes

Acute hypoxia stress occurs with temperature, seasonal, and compositional changes in water, particularly in high-density aquaculture situations. Fishes respond to acute hypoxia strongly. First, they try to get more oxygen by breathing directly through the mouth, this is usually referred to as floating heads.

Second, they alter the shape and structure of the gill to enhance oxygen exchange [41–46]. Some fishes alter the cardiac K(ATP) channel, metabolic rate, and increase the number of red blood cells [47,48].

Even though fishes can have an acute reaction to acute hypoxia to maintain normal physical activity, a sudden lack of oxygen will result in mortality. In aquaculture this is referred to as turning pond. The brain and cardiac cells of fishes go through apoptosis under hypoxia, this is the main cause of turning pond. Compared with other vertebrates, some of the key factors in the hypoxia signaling pathway control this physical process [49].

Elucidating the reasons and mechanisms of fish mortality under acute hypoxia will help to prevent or diminish yield losses in aquaculture.

2.3 The regulatory network of fish genes in response to hypoxia

Because of their specific habits, investigating the hypoxia signaling pathway and the key factors in its regulatory network in different fish species is necessary. To date, some key factors related to fish hypoxia adaptation and the hypoxia signaling pathway have been identified (Table 1).

2.3.1 Hypoxia-inducible factors

Fishes possess homologs of HIF- α and HIF- β , which play similar roles in mammals. The fish HIF-1 α sequence, which encodes a protein of 766 amino acids and includes basic-helix-loop-helix, PAS, and ODD domains of HIF α -subunits, was first reported from rainbow trout [50]. To date, HIF-1 α sequences have been reported in several fish species, e.g., Atlantic croaker (*Micropogonias undulatus*) [51], zebrafish (*Danio rerio*) [52], Wuchang bream (*Megalobrama amblycephala*) [53], scale-less carp (*Gymnocypris przewalskii*) [54], ruff (*Gymnocephalus cernuus*), flounder (*Platichthys flesus*), perch (*Perca fluviatilis*), stickleback (*Gasterosteus aculeatus*), Russian sturgeon (*Acipenser gueldenstaedtii*) [55], sea bass (*Dicentrarchus labrax*) [56], Indian catfish (*Clarias batrachus*) [57], and channel catfish (*Ictalurus punctatus*) [58]. Additionally, HIF-2 α has been identified in zebrafish [52], Atlantic croaker [51], grass carp (*Ctenopharyngodon idella*) [59], Indian catfish [57], and

Table 1 Hypoxia signaling proteins in fishes

Gene name	Fishes
HIF-1 α	<i>Oncorhynchus mykiss</i>
	<i>Micropogonias undulates</i>
	<i>Danio rerio</i>
	<i>Megalobrama amblycephala</i>
	<i>Gymnocypris przewalskii</i>
	<i>Gymnocephalus cernuus</i>
	<i>Platichthys flesus</i>
	<i>Perca fluviatilis</i>
	<i>Gasterosteus aculeatus</i>
	<i>Acipenser gueldenstaedtii</i>
	<i>Dicentrarchus labrax</i>
<i>Clarias batrachus</i>	
<i>Ictalurus punctatus</i>	
HIF-2 α	<i>Danio rerio</i>
	<i>Micropogonias undulates</i>
	<i>Ctenopharyngodon idellus</i>
	<i>Clarias batrachus</i>
PHD	<i>Ictalurus punctatus</i>
	<i>Salmo salar</i>
	<i>Oreochromis niloticus</i>
FIH	<i>Danio rerio</i>
	<i>Ictalurus punctatus</i>
VHL	<i>Danio rerio</i>

channel catfish [58].

Because of HIF- α 's important role in the hypoxia signaling pathway, the relationship between HIF expression and hypoxia has been investigated in several fishes. Eurasian perch (*Perca fluviatilis*) is a hypoxia-sensitive fresh water fish in which acute hypoxia can induce HIF-1 α expression in the brain and liver, whereas chronic hypoxia leads to a significant change in HIF-1 α expression in muscle [60]. The Indian catfish is a hypoxia-tolerant species, in which short-term exposure to hypoxia induces HIF-1 α and HIF-2 α expression in the brain, liver, and kidney, whereas the long-term exposure induces HIF-1 α expression in the spleen and HIF-2 α expression in muscle [57]. Furthermore, in channel catfish, another hypoxia-tolerant fish, HIF- α mRNA expression decreases under hypoxia for the first 1.5 h and then increases after 5 h [42].

The prolyl hydroxylase (PHD) proteins belong to the Fe(II) and 2-oxoglutarate-dependent oxygenase superfamily, whose activities are closely associated with oxygen levels. Thus, they are considered oxygen sensors [61]. To date, PHD sequences have been reported in some fish species, such as Atlantic salmon, channel catfish, Nile tilapia, and zebrafish [62]. Even though little is known about the function of fish PHDs, they are thought to act similarly to those of other organisms, which play important roles in HIF- α stability.

2.3.2 Hypoxia-inducible factor asparaginyl hydroxylase (FIH-1)

FIH-1, similar to PHD family members, is also a hydroxylase. However, FIH-1 cannot cause HIF- α proteasomal degradation after hydroxylating HIF- α . FIH-1 can inhibit the recruitment of the transcriptional coactivators p300/CBP. To date, FIH-1 sequences have been identified in zebrafish [63] and channel catfish [58].

2.3.3 VHL

VHL belongs to a component of the VBC E3 ligase complex, which acts as an E3 ligase. Under normoxia, pVHL recognizes hydroxylated HIF- α and causes HIF- α proteasomal degradation. Therefore, pVHL plays an important role in the regulation of the hypoxia signaling pathway. To date, zebrafish VHL is well studied. VHL-mutant zebrafish have been identified through mutant screening. Similar to that of pVHL-null mice, VHL-mutant zebrafish are embryonically lethal and develop erythrocytosis [64].

3 Hypoxia adaptation mechanisms in fishes

The key adaptation to long-term hypoxia is a reduction in metabolic rate and demands. However, whether or not anoxic survival is associated with reductions in both ATP production and use remains unclear. Protein synthesis consumes a lot of ATP. For hypoxia-tolerant crucian carp, the rate of protein synthesis decreases rapidly under both hypoxia and anoxia; however, in hypoxia-sensitive fishes, the rate of protein synthesis remains constant. In this process, AMPK might function as a sensor that regulates protein synthesis under hypoxia. When intracellular AMP is increased, AMPK is activated; whereas when intracellular AMP is decreased, AMPK is inhibited. Moreover, AMPK is also regulated by phosphorylation [65].

Adenosine is a low-oxygen signal. In the hypoxia-tolerant fishes, such as the crucian carp, adenosine has a powerful depressant effect on protein synthesis and Na⁺-K⁺ ATPase activity [66]. Adenosine receptor blockade in the crucian carp results in a threefold increase in ethanol release [67]. Additionally, adenosine and adenosine receptor blockers have surprising impacts on cerebral blood flow. In the crucian carp, cerebral blood flow increases 2.2-fold during anoxia, and superfusing the brain with aminophylline abolishes this increase. Thus, adenosine seems to be an important anoxic signal mediating vasodilation and cerebral blood flow [68].

Tolerance of acidic metabolic end products is another mechanism for fish hypoxia tolerance. The crucian carp can use ethanol as an anaerobic end product. Under anoxia, they maintain ion gradients in hepatocytes and convert lactate to ethanol. Ethanol can be eliminated by diffusion across the gills, avoiding lactic acid accumulation [69].

A most significant issue in low oxygen tolerance is the

avoidance of free-radical-mediated injury during reoxygenation. In some species, such as the crucian carp, the constitutive expression of defense molecules is not remarkable; it is increased only during hypoxia/reoxygenation [70]. Under anoxia, antioxidant enzyme activity increases, whereas most other enzyme activities, e.g., brain Se-GPX, liver catalase, and G6PDH, do not [71]. After several hours of extreme hypoxia, liver, brain, and gill superoxide dismutase (SOD) activity increases in carp, *Cyprinus carpio* [72].

Identifying the HIF- α down-stream targets and determining their functions will elucidate hypoxia adaptation mechanisms in fishes. Zhong et al. [73] have identified some hypoxia induced genes, e.g., ERO1-L, p53, CPO, HO-1, MKP2, PFK-2, cystatin B, Glut1, BTG1, TGF-beta1 and PGAM1, in crucian carp blastulae embryonic cells by suppression subtractive hybridization. Liao et al. [74] carried out transcriptome analysis for crucian carp and found hypoxia induced genes related to glycolysis/gluconeogenesis. Everett et al. [75] analyzed the effect of short-term hypoxic exposure on metabolic gene expression. Additionally, Chen et al. [76] analyzed hypoxia-induced changes in the zebrafish skeletal muscle proteome.

To date, many hypoxia-induced genes related to fish hypoxia adaptation have been identified and reported. GLUTs (Glucose transporters) are hypoxia-induced, metabolic genes that facilitate cellular glucose uptake. Under hypoxia, GLUTs are induced to enhance glucose transport, resulting in cellular energy demands. The full length cDNA of grass carp GLUT1, which is highly expressed under hypoxia, has been cloned [77]. When Atlantic cod were exposed to a hypoxic challenge, GLUT1 expression in the gill was significantly up-regulated [78]. Terova et al. [79] found GLUT2 mRNA changes in response to hypoxia in sea bass, a hypoxia-sensitive species, supporting the view that GLUT2 is involved in the hypoxia adaptation response.

Erythropoietin (Epo) is a hypoxia-induced gene related to erythropoiesis, which is a glycoprotein hormone that controls red blood cell production. The first Epo gene fish cloned was that of the pufferfish (Fugu; *Takifugu rubripes*). In Fugu, Epo is induced by hypoxia and is mainly expressed in the heart [80–82]. However, in other teleost fishes, such as rainbow trout, carp, and eel, Epo was expressed in the kidney, the kidney is the major organ for erythropoiesis in fishes [83].

VEGF is a hypoxia-induced gene related to angiogenesis, which stimulates the proliferation of blood vessels to increase oxygen supply. In the orange-spotted grouper (*Epinephelus coioides*), hypoxia can induce VEGF expression, which in turn increases oxygen delivery. Hypoxia also induces VEGF expression in salmon [84,85].

Both different cell lines and the zebrafish model have been employed to determine the function of hypoxia-induced genes. Wang et al. [86] cloned the hypoxia-induced HO-1 gene in the crucian carp. Using cell lines, they found that HO-1 plays an important role against hy-

poxia-induced cell death. Feng et al. [49] found that zebrafish p53 can protect hypoxia-induced cell death by suppressing BNIP3 expression, a hypoxia-induced cell death gene. Additionally, HIF-1 α is crucial for early zebrafish embryogenesis, particularly for neural crest chemotaxis and the central nervous system [87,88].

Compared with the mouse model, gene knock-out assays in zebrafish are difficult to perform due to lack of suitable zebrafish embryonic stem cell lines. Thus, little is known about the *in vivo* function of hypoxia-adaptation-related genes in fishes. Recently, TALEN and CRISP/Cas9 techniques have been established and rapidly developed, making knocking out gene assays in zebrafish more convenient. Therefore, information on the *in vivo* function of hypoxia-adaptation-related genes in fishes is bound to be available in the near future [89]. As a result of differing degrees of pollution in aquaculture waters, protecting the aquatic environment, intensive cultivation modes such as intensive pond aquaculture, and industrial aquaculture have become fundamental to aquaculture in China. Avoiding water pollution is the main means of increasing density and yield per unit of water body. Because of the increase in aquaculture density and yield, it is necessary to improve fish species/strains hypoxia tolerance capabilities.

To breed new fish species or strains with higher hypoxia tolerance, we must first determine the mechanisms of hypoxia tolerance and adaptation in fish. Here are some basic strategies: First, comparative omics, such as genomics, transcriptomics, and proteomics to compare different hypoxia-tolerant fish species distributed in the same waters, or closely related species distributed in different waters with varying oxygen concentration. Second, microarray, transcriptome sequencing, and proteomics to analyze the representative hypoxia-sensitive and tolerant species following hypoxia treatment. Subsequently, using bioinformatic methods, the gene regulatory network for fish hypoxia adaptation and tolerance can be constructed and the key genes identified. By taking advantage of gene function analysis approaches, the functions of key genes can be elucidated, the gene's contributions to fish hypoxia tolerance can be estimated, and the molecular markers related to hypoxia adaptation and tolerance can be screened. Based on these data, we can breed new species or strains with higher hypoxia tolerance using either transgenic techniques or classic hybridization.

- 1 Semenza GL. Hypoxia-inducible factors in physiology and medicine. *Cell*, 2012, 148: 399–408
- 2 Semenza GL. Oxygen sensing, hypoxia-inducible factors, and disease pathophysiology. *Ann Rev Pathol*, 2014, 9: 47–71
- 3 Majmundar AJ, Wong WJ, Simon MC. Hypoxia-inducible factors and the response to hypoxic stress. *Mol Cell*, 2010, 40: 294–309
- 4 Dunwoodie SL. The role of hypoxia in development of the mammalian embryo. *Dev Cell*, 2009, 17: 755–773
- 5 Schofield CJ, Ratcliffe PJ. Oxygen sensing by HIF hydroxylases. *Nat*

- Rev Mol Cell Biol, 2004, 5: 343–354
- 6 Bickler PE, Buck LT. Hypoxia tolerance in reptiles, amphibians, and fishes: life with variable oxygen availability. *Ann Rev Pathol*, 2007, 69: 145–170
 - 7 Greer SN, Metcalf JL, Wang Y, Ohh M. The updated biology of hypoxia-inducible factor. *EMBO J*, 2012, 31: 2448–2460
 - 8 Aragonés J, Fraisl P, Baes M, Carmeliet P. Oxygen sensors at the crossroad of metabolism. *Cell Metab*, 2009, 9: 11–22
 - 9 Kaelin WG. Von Hippel-Lindau disease. *Ann Rev Pathol*, 2007, 2: 145–173
 - 10 Rius J, Guma M, Schachtrup C, Akassoglou K, Zinkernagel AS, Nizet V, Johnson RS, Haddad GG, Karin M. NF-kappaB links innate immunity to the hypoxic response through transcriptional regulation of HIF-1alpha. *Nature*, 2008, 453: 807–811
 - 11 Flugel D, Gorlach A, Michiels C, Kietzmann T. Glycogen synthase kinase 3 phosphorylates hypoxia-inducible factor 1alpha and mediates its destabilization in a VHL-independent manner. *Mol Cell Biol*, 2007, 27: 3253–3265
 - 12 Ryu JH, Li SH, Park HS, Park JW, Lee B, Chun YS. Hypoxia-inducible factor alpha subunit stabilization by NEDD8 conjugation is reactive oxygen species-dependent. *J Biol Chem*, 2011, 286: 6963–6970
 - 13 Sang N, Fang J, Srinivas V, Leshchinsky I, Caro J. Carboxyl-terminal transactivation activity of hypoxia-inducible factor 1 alpha is governed by a von Hippel-Lindau protein-independent, hydroxylation-regulated association with p300/CBP. *Mol Cell Biol*, 2002, 22: 2984–2992
 - 14 Mehta R, Steinkraus K A, Sutphin GL, Ramos FJ, Shamieh LS, Huh A, Davis C, Chandler-Brown D, Kaerberlein M. Proteasomal regulation of the hypoxic response modulates aging in *C. elegans*. *Science*, 2009, 324: 1196–1198
 - 15 Chen D, Thomas EL, Kapahi P. HIF-1 modulates dietary restriction-mediated lifespan extension via IRE-1 in *Caenorhabditis elegans*. *PLoS Genet*, 2009, 5: e1000486
 - 16 Zhong L, D'Urso A, Toiber D, Sebastian C, Henry RE, Vadysirisack DD, Guimaraes A, Marinelli B, Wikstrom JD, Nir T, Clish CB, Vaitheeswaran B, Iliopoulos O, Kurland I, Dor Y, Weissleder R, Shirihai OS, Ellisen LW, Espinosa JM, Mostoslavsky R. The histone deacetylase Sirt6 regulates glucose homeostasis via Hif1alpha. *Cell*, 2010, 140: 280–293
 - 17 Lim JH, Lee YM, Chun YS, Chen J, Kim JE, Park JW. Sirtuin 1 modulates cellular responses to hypoxia by deacetylating hypoxia-inducible factor 1alpha. *Mol Cell*, 2010, 38: 864–878
 - 18 Dioum EM, Chen R, Alexander MS, Zhang Q, Hogg RT, Gerard RD, Garcia JA. Regulation of hypoxia-inducible factor 2alpha signaling by the stress-responsive deacetylase sirtuin 1. *Science*, 2009, 324: 1289–1293
 - 19 Gomes AP, Price NL, Ling AJ, Moslehi JJ, Montgomery MK, Rajman L, White JP, Teodoro JS, Wrann CD, Hubbard BP, Mercken EM, Palmeira CM, de Cabo R, Rolo AP, Turner N, Bell EL, Sinclair DA. Declining NAD(+) induces a pseudohypoxic state disrupting nuclear-mitochondrial communication during aging. *Cell*, 2013, 155: 1624–1638
 - 20 Hubbi ME, Hu H, Kshitiz Gilkes DM, Semenza GL. Sirtuin-7 inhibits the activity of hypoxia-inducible factors. *J Biol Chem*, 2013, 288: 20768–20775
 - 21 Finley LW, Carracedo A, Lee J, Souza A, Egja A, Zhang J, Teruya-Feldstein J, Moreira PI, Cardoso SM, Clish CB, Pandolfi PP, Haigis MC. SIRT3 opposes reprogramming of cancer cell metabolism through HIF1alpha destabilization. *Cancer Cell*, 2011, 19: 416–428
 - 22 Shao R, Zhang FP, Tian F, Anders Friberg P, Wang X, Sjolund H, Billig H. Increase of SUMO-1 expression in response to hypoxia: direct interaction with HIF-1alpha in adult mouse brain and heart *in vivo*. *FEBS Lett*, 2004, 569: 293–300
 - 23 Carbia-Nagashima A, Gerez J, Perez-Castro C, Paez-Pereda M, Silberstein S, Stalla GK, Holsboer F, Arzt E. RSUME, a small RWD-containing protein, enhances SUMO conjugation and stabilizes HIF-1alpha during hypoxia. *Cell*, 2007, 131: 309–323
 - 24 Cheng J, Kang X, Zhang S, Yeh ET. SUMO-specific protease 1 is essential for stabilization of HIF1alpha during hypoxia. *Cell*, 2007, 131: 584–595
 - 25 Luo W, Hu H, Chang R, Zhong J, Knabel M, O'Meally R, Cole RN, Pandey A, Semenza GL. Pyruvate kinase M2 is a PHD3-stimulated coactivator for hypoxia-inducible factor 1. *Cell*, 2011, 145: 732–744
 - 26 Montagner M, Enzo E, Forcato M, Zanconato F, Parenti A, Rampazzo E, Basso G, Leo G, Rosato A, Biciato S, Cordenonsi M, Piccolo S. SHARP1 suppresses breast cancer metastasis by promoting degradation of hypoxia-inducible factors. *Nature*, 2012, 487: 380–384
 - 27 Chen Z, Liu X, Mei Z, Wang Z, Xiao W. EAF2 suppresses hypoxia-induced factor 1alpha transcriptional activity by disrupting its interaction with coactivator CBP/p300. *Mol Cell Biol*, 2014, 34: 1085–1099
 - 28 Nakayama K, Frew IJ, Hagensen M, Skals M, Habelhah H, Bhoumik A, Kadoya T, Erdjument-Bromage H, Tempst P, Frappell PB, Bowtell DD, Ronai Z. Siah2 regulates stability of prolyl-hydroxylases, controls HIF1alpha abundance, and modulates physiological responses to hypoxia. *Cell*, 2004, 117: 941–952
 - 29 Jung CR, Hwang KS, Yoo J, Cho WK, Kim JM, Kim WH, Im DS. E2-EPF UCP targets pVHL for degradation and associates with tumor growth and metastasis. *Nat Med*, 2006, 12: 809–816
 - 30 Bernardi R, Guernah I, Jin D, Grisendi S, Alimonti A, Teruya-Feldstein J, Cordon-Cardo C, Simon MC, Rafii S, Pandolfi PP. PML inhibits HIF-1alpha translation and neoangiogenesis through repression of mTOR. *Nature*, 2006, 442: 779–785
 - 31 Huang C, Han Y, Wang Y, Sun X, Yan S, Yeh ET, Chen Y, Cang H, Li H, Shi G, Cheng J, Tang X, Yi J. SENP3 is responsible for HIF-1 transactivation under mild oxidative stress via p300 de-SUMOylation. *EMBO J*, 2009, 28: 2748–2762
 - 32 Foxler DE, Bridge KS, James V, Webb TM, Mee M, Wong SC, Feng Y, Constantin-Teodosiu D, Petrusdottir TE, Bjornsson J, Ingvarsson S, Ratcliffe PJ, Longmore GD, Sharp TV. The LIMD1 protein bridges an association between the prolyl hydroxylases and VHL to repress HIF-1 activity. *Nat Cell Biol*, 2012, 14: 201–208
 - 33 Nilsson GE, Renshaw GM. Hypoxic survival strategies in two fishes: extreme anoxia tolerance in the North European crucian carp and natural hypoxic preconditioning in a coral-reef shark. *J Exp Biol*, 2004, 207: 3131–3139
 - 34 Nilsson GE. Surviving anoxia with the brain turned on. *News Physiol Sci*, 2001, 16: 217–221
 - 35 Lutz PL, Nilsson GE. Vertebrate brains at the pilot light. *Resp Physiol Neurobiol*, 2004, 141: 285–296
 - 36 Stecyk JA, Stenslokken KO, Farrell AP, Nilsson GE. Maintained cardiac pumping in anoxic crucian carp. *Science*, 2004, 306: 77
 - 37 Shoubridge EA, Hochachka PW. Ethanol: novel end product of vertebrate anaerobic metabolism. *Science*, 1980, 209: 308–309
 - 38 Roesner A, Mitz SA, Hankeln T, Burmester T. Globins and hypoxia adaptation in the goldfish, *Carassius auratus*. *FEBS J*, 2008, 275: 3633–3643
 - 39 Rytkonen KT, Akbarzadeh A, Miandare HK, Kamei H, Duan C, Leder EH, Williams TA, Nikinmaa M. Subfunctionalization of cyprinid hypoxia-inducible factors for roles in development and oxygen sensing. *Evolution*, 2013, 67: 873–882
 - 40 Chi W, Gan X, Xiao W, Wang W, He S. Different evolutionary patterns of hypoxia-inducible factor alpha (HIF-alpha) isoforms in the basal branches of Actinopterygii and Sarcopterygii. *FEBS Open Bio*, 2013, 3: 479–483
 - 41 Nilsson GE, Dymowska A, Stecyk JA. New insights into the plasticity of gill structure. *Resp Physiol Neurobiol*, 2012, 184: 214–222
 - 42 Turko AJ, Cooper CA, Wright PA. Gill remodelling during terrestrial acclimation reduces aquatic respiratory function of the amphibious fish *Kryptolebias marmoratus*. *J Exp Biol*, 2012, 215: 3973–3980
 - 43 Dhillon RS, Yao L, Matey V, Chen BJ, Zhang AJ, Cao ZD, Fu SJ, Brauner CJ, Wang YS, Richards JG. Interspecific differences in hypoxia-induced gill remodeling in carp. *Physiol Biochem Zool*, 2013, 86: 727–739
 - 44 Tzaneva V, Vadeboncoeur C, Ting J, Perry SF. Effects of hypoxia-induced gill remodeling on the innervation and distribution of

- ionocytes in the gill of goldfish, *Carassius auratus*. J Comp Neurol, 2014, 522: 118–130
- 45 Mitrovic D, Dymowska A, Nilsson GE, Perry SF. Physiological consequences of gill remodeling in goldfish (*Carassius auratus*) during exposure to long-term hypoxia. Am J Physiol Regul Integr Comp Physiol, 2009, 297: R224–234
- 46 Zachar PC, Jonz MG. Oxygen sensitivity of gill neuroepithelial cells in the anoxia-tolerant goldfish. Adv Exp Med Biol, 2012, 758: 167–172
- 47 Cameron JS, DeWitt JP, Ngo TT, Yajnik T, Chan S, Chung E, Kang E. Cardiac K(ATP) channel alterations associated with acclimation to hypoxia in goldfish (*Carassius auratus* L.). Comp Biochem Physiol A Mol Integr Physiol, 2013, 164: 554–564
- 48 Capossela KM, Brill RW, Fabrizio MC, Bushnell PG. Metabolic and cardiorespiratory responses of summer flounder *Paralichthys dentatus* to hypoxia at two temperatures. J Fish Biol, 2012, 81: 1043–1058
- 49 Feng X, Liu X, Zhang W, Xiao W. p53 directly suppresses BNIP3 expression to protect against hypoxia-induced cell death. EMBO J, 2011, 30: 3397–3415
- 50 Soitamo AJ, Rabergh CM, Gassmann M, Sistonen L, Nikinmaa M. Characterization of a hypoxia-inducible factor (HIF-1alpha) from rainbow trout. Accumulation of protein occurs at normal venous oxygen tension. J Biol Chem, 2001, 276: 19699–19705
- 51 Rahman MS, Thomas P. Molecular cloning, characterization and expression of two hypoxia-inducible factor alpha subunits, HIF-1alpha and HIF-2alpha, in a hypoxia-tolerant marine teleost, Atlantic croaker (*Micropogonias undulatus*). Gene, 2007, 396: 273–282
- 52 Rojas DA, Perez-Munizaga DA, Centanin L, Antonelli M, Wappner P, Allende ML, Reyes AE. Cloning of HIF-1alpha and HIF-2alpha and mRNA expression pattern during development in zebrafish. Gene Exp Patterns, 2007, 7: 339–345
- 53 Shen RJ, Jiang XY, Pu JW, Zou SM. HIF-1alpha and -2alpha genes in a hypoxia-sensitive teleost species *Megalobrama amblycephala*: cDNA cloning, expression and different responses to hypoxia. Comp Biochem Physiol B Biochem Mol Biol, 2010, 157: 273–280
- 54 Cao YB, Chen XQ, Wang S, Wang YX, Du JZ. Evolution and regulation of the downstream gene of hypoxia-inducible factor-1alpha in naked carp (*Gymnocypris przewalskii*) from Lake Qinghai, China. J Mol Evol, 2008, 67: 570–580
- 55 Rytkonen KT, Vuori KA, Primmer CR, Nikinmaa M. Comparison of hypoxia-inducible factor-1 alpha in hypoxia-sensitive and hypoxia-tolerant fish species. Comp Biochem Physiol Part D Genomics Proteomics, 2007, 2: 177–186
- 56 Terova G, Rimoldi S, Cora S, Bernardini G, Gornati R, Saroglia M. Acute and chronic hypoxia affects HIF-1 alpha mRNA levels in sea bass (*Dicentrarchus labrax*). Aquaculture, 2008, 279: 150–159
- 57 Mohindra V, Tripathi RK, Singh RK, Lal KK. Molecular characterization and expression analysis of three hypoxia-inducible factor alpha subunits, HIF-1alpha, -2alpha and -3alpha in hypoxia-tolerant Indian catfish, *Clarias batrachus* [Linnaeus, 1758]. Mol Biol Rep, 2013, 40: 5805–5815
- 58 Geng X, Feng J, Liu S, Wang Y, Arias C, Liu Z. Transcriptional regulation of hypoxia inducible factors alpha (HIF-alpha) and their inhibiting factor (FIH-1) of channel catfish (*Ictalurus punctatus*) under hypoxia. Comp Biochem Physiol B Biochem Mol Biol, 2014, 169: 38–50
- 59 Law SH, Wu RS, Ng PK, Yu RM, Kong RY. Cloning and expression analysis of two distinct HIF-alpha isoforms—gcHIF-1alpha and gcHIF-4alpha—from the hypoxia-tolerant grass carp, *Ctenopharyngodon idellus*. BMC Mol Biol, 2006, 7: 15
- 60 Rimoldi S, Terova G, Ceccuzzi P, Marelli S, Antonini M, Saroglia M. HIF-1alpha mRNA levels in Eurasian perch (*Perca fluviatilis*) exposed to acute and chronic hypoxia. Mol Biol Rep, 2012, 39: 4009–4015
- 61 Kaelin WG Jr., Ratcliffe PJ. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. Mol Cell, 2008, 30: 393–402
- 62 Kaelin WG, Sawyers CL, Mihich E. Nineteenth Annual Pezcoller Symposium: hypothesis-driven clinical investigation in cancer. Cancer Res, 2007, 67: 11102–11105
- 63 Nikinmaa M, Rees BB. Oxygen-dependent gene expression in fishes. Am J Physiol Regul Integr Comp Physiol, 2005, 288: R1079–1090
- 64 van Rooijen E, Voest EE, Logister I, Korving J, Schwerte T, Schulte-Merker S, Giles RH, van Eeden FJ. Zebrafish mutants in the von Hippel-Lindau tumor suppressor display a hypoxic response and recapitulate key aspects of Chuvash polycythemia. Blood, 2009, 113: 6449–6460
- 65 Lindsley JE, Rutter J. Nutrient sensing and metabolic decisions. Comp Biochem Physiol B Biochem Mol Biol, 2004, 139: 543–559
- 66 Krumschnabel G, Biasi C, Wieser W. Action of adenosine on energetics, protein synthesis and K(+) homeostasis in teleost hepatocytes. J Exp Biol, 2000, 203: 2657–2665
- 67 Nilsson GE. The adenosine receptor blocker aminophylline increases anoxic ethanol excretion in crucian carp. Am J Physiol, 1991, 261: R1057–1060
- 68 Nilsson GE, Hylland P, Lofman CO. Anoxia and adenosine induce increased cerebral blood flow rate in crucian carp. Am J Physiol, 1994, 267: R590–595
- 69 Krumschnabel G, Schwarzbaum PJ, Lisch J, Biasi C, Wieser W. Oxygen-dependent energetics of anoxia-tolerant and anoxia-intolerant hepatocytes. J Exp Biol, 2000, 203: 951–959
- 70 Hermes-Lima M, Zenteno-Savin T. Animal response to drastic changes in oxygen availability and physiological oxidative stress. Comp Biochem Physiol Toxicol Pharmacol, 2002, 133: 537–556
- 71 Lushchak VI, Lushchak LP, Mota AA, Hermes-Lima M. Oxidative stress and antioxidant defenses in goldfish *Carassius auratus* during anoxia and reoxygenation. Am J Physiol Regul Integr Comp Physiol, 2001, 280: R100–107
- 72 Vig E, Gabrielak T, Leyko W, Nemcsok J, Matkovic B. Purification and characterization of Cu,Zn-superoxide dismutase from common carp liver. Comp Biochem Physiol B Biochem Mol Biol, 1989, 94: 395–397
- 73 Zhong XP, Wang D, Zhang YB, Gui JF. Identification and characterization of hypoxia-induced genes in *Carassius auratus* blastulae embryonic cells using suppression subtractive hybridization. Comp Biochem Physiol B Biochem Mol Biol, 2009, 152: 161–170
- 74 Liao X, Cheng L, Xu P, Lu G, Wachholtz M, Sun X, Chen S. Transcriptome analysis of crucian carp (*Carassius auratus*), an important aquaculture and hypoxia-tolerant species. PLoS One, 2013, 8: e62308
- 75 Everett MV, Antal CE, Crawford DL. The effect of short-term hypoxic exposure on metabolic gene expression. J Exp Zool A Ecol Genet Physiol, 2012, 317: 9–23
- 76 Chen K, Cole RB, Rees BB. Hypoxia-induced changes in the zebrafish (*Danio rerio*) skeletal muscle proteome. J Proteomics, 2013, 78: 477–485
- 77 Zhang Z, Wu RS, Mok HO, Wang Y, Poon WW, Cheng SH, Kong RY. Isolation, characterization and expression analysis of a hypoxia-responsive glucose transporter gene from the grass carp, *Ctenopharyngodon idellus*. Eur J Biochem/FEBS, 2003, 270: 3010–3017
- 78 Hall JR, Richards RC, MacCormack TJ, Ewart KV, Driedzic WR. Cloning of GLUT3 cDNA from Atlantic cod (*Gadus morhua*) and expression of GLUT1 and GLUT3 in response to hypoxia. Biochim Biophys Acta, 2005, 1730: 245–252
- 79 Terova G, Forchino A, Rimoldi S, Brambilla F, Antonini M, Saroglia M. Bio-Mos: an effective inducer of dicentracin gene expression in European sea bass (*Dicentrarchus labrax*). Comp Biochem Physiol B Biochem Mol Biol, 2009, 153: 372–377
- 80 Chou CF, Tohari S, Brenner S, Venkatesh B. Erythropoietin gene from a teleost fish, Fugu rubripes. Blood, 2004, 104: 1498–1503
- 81 Chu CY, Cheng CH, Chen GD, Chen YC, Hung CC, Huang KY, Huang CJ. The zebrafish erythropoietin: functional identification and biochemical characterization. FEBS Lett, 2007, 581: 4265–4271
- 82 Paffett-Lugassy N, Hsia N, Fraenkel PG, Paw B, Leshinsky I, Barut B, Bahary N, Caro J, Handin R, Zon LI. Functional conservation of erythropoietin signaling in zebrafish. Blood, 2007, 110: 2718–2726
- 83 Pierron F, Baudrimont M, Gonzalez P, Bourdineaud JP, Elie P, Mas-sabauu JC. Common pattern of gene expression in response to hypoxia or cadmium in the gills of the European glass eel (*Anguilla*

- anguilla*). *Environ Sci Technol*, 2007, 41: 3005–3011
- 84 Vuori KA, Soitamo A, Vuorinen PJ, Nikinmaa M. Baltic salmon (*Salmo salar*) yolk-sac fry mortality is associated with disturbances in the function of hypoxia-inducible transcription factor (HIF-1 α) and consecutive gene expression. *Aquat Toxicol*, 2004, 68: 301–313
- 85 Yu RM, Ng PK, Tan T, Chu DL, Wu RS, Kong RY. Enhancement of hypoxia-induced gene expression in fish liver by the aryl hydrocarbon receptor (AhR) ligand, benzo[a]pyrene (BaP). *Aquat Toxicol*, 2008, 90: 235–242
- 86 Wang D, Zhong XP, Qiao ZX, Gui JF. Inductive transcription and protective role of fish heme oxygenase-1 under hypoxic stress. *J Exp Biol*, 2008, 211: 2700–2706
- 87 Stevenson TJ, Trinh T, Kogelschatz C, Fujimoto E, Lush ME, Piotrowski T, Brimley CJ, Bonkowsky JL. Hypoxia disruption of vertebrate CNS pathfinding through ephrinB2 is rescued by magnesium. *PLoS Genet*, 2012, 8: e1002638
- 88 Barriga EH, Maxwell PH, Reyes AE, Mayor R. The hypoxia factor Hif-1 α controls neural crest chemotaxis and epithelial to mesenchymal transition. *J Cell Biol*, 2013, 201: 759–776
- 89 Zhang LL, Zhou Q. CRISPR/Cas technology: a revolutionary approach for genome engineering. *Sci China Life Sci*, 2014, 57: 639–640

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.