

Inhibitory leukocyte immunoglobulin-like receptors in cancer development

ZHANG FeiFei^{1,2}, ZHENG JunKe^{1,2*}, KANG XunLei³, DENG Mi³, LU ZhiGang³,
KIM Jaehyup³ & ZHANG ChengCheng^{3*}

¹Hongqiao International Institute of Medicine, Shanghai Tongren Hospital; Faculty of Basic Medicine, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China;

²Key Laboratory of Cell Differentiation and Apoptosis of Chinese Ministry of Education, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China;

³Departments of Physiology and Developmental Biology, University of Texas Southwestern Medical Center, Dallas 75390, USA

Received July 29, 2015; accepted August 16, 2015; published online November 13, 2015

Inhibitory leukocyte immunoglobulin-like receptors (LILRB1-5) signal through immunoreceptor tyrosine-based inhibitory motifs (ITIMs) in their intracellular domains and recruit phosphatases protein tyrosine phosphatase, non-receptor type 6 (PTPN6, SHP-1), protein tyrosine phosphatase, non-receptor type 6 (PTPN6, SHP-2), or Src homology 2 domain containing inositol phosphatase (SHIP) to negatively regulate immune cell activation. These receptors are known to play important regulatory roles in immune and neuronal functions. Recent studies demonstrated that several of these receptors are expressed by cancer cells. Importantly, they may directly regulate development, drug resistance, and relapse of cancer, and the activity of cancer stem cells. Although counterintuitive, these findings are consistent with the generally immune-suppressive and thus tumor-promoting roles of the inhibitory receptors in the immune system. This review focuses on the ligands, expression pattern, signaling, and function of LILRB family in the context of cancer development. Because inhibition of the signaling of certain LILRBs directly blocks cancer growth and stimulates immunity that may suppress tumorigenesis, but does not disturb normal development, LILRB signaling pathways may represent ideal targets for treating hematological malignancies and perhaps other tumors.

immunoreceptor tyrosine-based inhibitory motifs, immunoreceptor tyrosine-based activation motif, leukocyte immunoglobulin-like receptor subfamily B, immunoglobulin-like transcript, leukocyte immunoglobulin-like receptor, phosphatase, ITIM, ITAM, LILRB, CD85, ILT, LIR, SHP-1, SHP-2, SHIP, MHC, HLA, signal transduction, leukemia, cancer

Citation: Zhang FF, Zheng JK, Kang XL, Deng M, Lu ZG, Kim J, Zhang CC. Inhibitory leukocyte immunoglobulin-like receptors in cancer development. *Sci China Life Sci*, 2015, 58: 1216–1225, doi: 10.1007/s11427-015-4925-1

1 Introduction to ITIM-containing receptors and LILRBs

First described in 1995, the immunoreceptor tyrosine-based

inhibitory motifs (ITIMs) are conserved 6 amino acid stretches (S/I/V/LxYxxI/V/L) found in the cytoplasmic tails of certain surface transmembrane receptors [1]. Ligand binding or interplay with other signaling receptors leads to a conformational change that is accompanied by the phosphorylation of tyrosines in the ITIM by Src kinases. This leads to the recruitment of SH2 domain-containing phosphatases. With the exception of IgG Fc receptor II-B

*Corresponding author (email: zhengjunke@sjtu.edu.cn; Alec.Zhang@UTSouthwestern.edu)

(Fc γ RIIB), which is the ITIM-containing receptor that only recruits the inositol-phosphatase SHIP, these receptors all bind tyrosine phosphatases SHP-1 or SHP-2 [2–4]. An isoleucine at the first position of the ITIM (IxYxxL/V) favors binding to SHP-1, whereas a leucine in the same position (LxYxxL/V) favors SHIP binding [5]. These phosphatases are known to negatively regulate immune cell activation. For example, SHP-1 dephosphorylates the activated immunoreceptor tyrosine-based activation motif (ITAM), Src proto-oncogene non-receptor tyrosine kinase (Src), spleen tyrosine kinase (Syk), zeta-chain associated protein kinase 70 kD (ZAP70), Lck/Yes-related novel protein tyrosine kinase (Lyn), phosphatidylinositol-4-phosphate 3-kinase (PI3K), phospholipase C gamma (PLC- γ), or Vav 1 guanine nucleotide exchange factor (Vav1) [6–9]. The consequences of signaling through ITIMs are not well defined. ITIMs may inhibit signaling from immune activating receptors, cytokine receptors, and tyrosine kinase receptors. A proteome-wide analysis performed a decade ago identified 109 human ITIM-containing receptors [10].

ITIMs have antagonistic functions relative to immunoreceptor tyrosine-based activation motifs (ITAMs). ITAM is a conserved amino acid sequence of YxxL/Ix(6-8)YxxL/I in the cytoplasmic tails of membrane proteins. ITIM transmits signals from various membrane receptors including T cell receptors, B cell receptors, activating leukocyte Ig-like receptors (LILRs), certain activating natural killer (NK) cell receptors and Fc receptors, and other activating receptors [11]. Like to ITIM-containing receptors, ligand engagement at a relevant receptor results in the activation of Src kinases, leading to the phosphorylation of tyrosines within the ITAM. Subsequent recruitment and activation of Syk or ZAP-70 tyrosine kinases in myeloid cells and lymphoid cells, respectively, initiate further downstream signaling events, usually resulting in immune activation [11].

The leukocyte Ig-like receptor subfamily B (LILRB) is an important group of ITIM-containing receptors that were cloned in 1997 [12–15] (see schematic in Figure 1). These receptors, also known as cluster of differentiation 85 (CD85), Ig-like transcripts (ILTs), or leukocyte Ig-like receptors (LIRs), are type I transmembrane glycoproteins containing extracellular ligand-binding Ig-like domains and intracellular ITIMs, and are thus classified as immune inhibitory receptors. The name LILRB was officially designated in 2001 to classify a group of inhibitory receptors, whereas LILRA refer to activating receptors [16]. It is known that LILRBs, including LILRB1, LILRB2, LILRB3, LILRB4, and LILRB5, are expressed on myeloid cells and certain other hematopoietic cells and can exert immunomodulatory effects on a wide range of immune cells [17]. It is noteworthy that LILRB1-5 are primate and human specific, with paired immunoglobulin-like receptor B (PirB) [18] and leukocyte immunoglobulin-like receptor, subfamily B, member 4 (Lilrb4, gp49B1) [19] as the only two mouse orthologs. The human genes encoding these receptors are found in a gene




Figure 1 (color online) Schematic of the ligands, extracellular Ig-domains (hexagons), and intracellular ITIMs (boxes) of human LILRBs.

cluster known as the leukocyte receptor complex at chromosomal region 19q13.4 [12,20].

LILRBs are encoded by rapidly evolving genes, and it is thus difficult to study their function in animal models, so the biological function and clinical significance of these receptors have remained poorly understood. Recently, we demonstrated that LILRB2 and a mouse ortholog PirB are not only expressed by immune cells but are also expressed by hematopoietic stem cells [21]. Moreover, several LILRBs, PirB, and LAIR1 (a close related ITIM-containing receptor) [22–25] are expressed by primitive and differentiated acute leukemia cells that support leukemia development [21,26]. A number of reports also showed that LILRBs are expressed in hematopoietic and solid cancer cells and in most cases exert tumor-promoting functions [27–41]. Ligands and signaling pathways mediated through LILRBs have also been identified [21,26,38,39,41–44]. Although counterintuitive, the direct tumor-supportive roles of LILRBs are consistent with the generally immune-suppressive and thus tumor-promoting functions of the inhibitory receptors in the immune system [45]. The roles of LILRBs in immune and neuronal diseases have been recently reviewed [2,4,46,47]. Here we will focus on the ligands, expression pattern, signaling, and the immune-modulatory and direct tumor-supportive roles of LILRBs in cancer development.

2 Ligands for ITIM-containing receptors

Different types of molecules were identified as ligands for ITIM-containing receptors. These include integrated membrane proteins (e.g., major histocompatibility complex (MHC) class I for LILRB1-2 [48,49]), extracellular matrix proteins (e.g., collagens for LAIR1 [22]), and soluble proteins (e.g., antibodies for Fc γ RIIB).

Not all the ligands for LILRBs were identified. LILRB1-2 binds classical and non-classical MHC molecules [48–50]. LILRB5 was shown to bind to MHC- or human leukocyte antigen (HLA)-Class I heavy chains [44]. LILRB1 and 2 were also shown to bind to non-HLA ligands, including S100A8 and S100A9 for LILRB1 [51], and CD1d [52], several angiopoietin-like proteins (Angptls) [21,42], oligo-

meric β -amyloid [43], and myelin inhibitors reticulon 4 (RTN4, Nogo66), myelin associated glycoprotein (MAG), and oligodendrocyte myelin glycoprotein (OMgp) for LILRB2 [53]. We recently demonstrated that multimeric Angptls bind and activate LILRB2 more effectively than does HLA-G [42]. The ligands for LILRB3 and 4 have not been identified. Although integrin $\alpha v \beta 3$ was shown to bind to mouse gp49B1, a possible ortholog of human LILRB4 [54], this dimeric integrin does not bind to human LILRB4.

3 Relevance to cancer

We performed an *in silico* analysis of the relationship between gene expression and the overall survival of acute myeloid leukemia (AML) patients using the TCGA database of AML patients (<https://tcga-data.nci.nih.gov/tcga/>). The expression of two out of 58 ITIM-containing receptors is positively correlated with the overall survival of AML patients; 20 of these receptors have negative correlation between expression and patient survival [26]. Importantly, we found that silencing of certain receptors individually lead to reduced cell growth [26]. These results suggest that some ITIM-containing receptors directly support human leukemia cell growth.

The *in silico* analysis further indicated that the levels of expression of several closely related LILRB family members, LILRB 1, 2, 3, and 4, and the related LAIR1 are higher in monocytic AML (M5 subtype) cells and inversely correlate with the overall survival of AML patients [26]. LAIR1 is a type I transmembrane glycoprotein that shares the same domain organization as LILRBs, containing one extracellular Ig-like domain that binds collagens or surfactant protein D and two intracellular ITIMs that recruit SHP-1 and SHP-2 [22–25]. LAIR1 is expressed on various lineages of hematopoietic cells and hematopoietic progenitor CD34⁺ cells [55]. The silencing of expression of LILRB2, 3, 4 or LAIR1 individually in human AML cell lines inhibits cell growth *in vitro* [26]. LILRB4 and LAIR1 are more highly expressed on primary human AML cells than on normal counterparts [26,56]. Importantly, the LAIR1⁺ cells are enriched for the activity of AML stem cells (AML SCs). Inhibition of expression of LAIR1 in human AML cell lines almost completely abolished leukemia development in xenografted NOD/SCID/IL2R γ null (NSG) mice [26]. In contrast, the studies of individual knockout mouse lines of *PirB* and *gp49B1* (the only mouse orthologs) and of the *LAIR1*-knockout mouse revealed no overt defects in normal hematopoiesis [21,57,58].

With the exception of LILRB1, LILRBs are not expressed at significant levels by adaptive immune T and B cells, although several *LILR* mRNAs are detected in subset of B cells. This pattern of cell expression suggests that LILRBs may primarily be involved in the innate immune

response. Also, the expression of LILRBs in tumor cells implies possible cancer-related functions. Below we summarize the cancer related roles of individual LILRBs.

3.1 LILRB1 (CD85J, ILT2, LIR1, MIR7)

LILRB1 contains four extracellular immunoglobulin domains, a transmembrane domain, and four cytoplasmic ITIMs. Among the LILRBs, LILRB1 has the most extensive cellular distribution. It is expressed on primary NK cells, B cells, various populations of T cells, monocytes/macrophages, eosinophils and basophils, dendritic cells [4], decidual macrophages [59], and osteoclasts [17]. LILRB1, 2, 3, and 4 are expressed on progenitor mast cells but not on mature mast cells [60]. LILRB1 binds various HLA class I molecules, including HLA-A, HLA-B, HLA-C, HLA-E, and HLA-G, with $\mu\text{mol L}^{-1}$ affinities [48,50]. It also binds the cytomegalovirus UL18 protein, which is a HLA class I homolog [13]. In addition, LILRB1 was reported to bind two calcium-binding proteins S100A8 and S100A9 [51]. Dimerized HLA-G induces more efficient LILRB1 signaling than the monomeric form [61]. LILRB1-mediated signaling is thought to control inflammatory responses and cytotoxicity to focus the immune response and limit autoreactivity, and LILRB1 functions as an inhibitor of immunity of T cells, B cells, NK cells, and macrophages. HLA-G is upregulated in human breast tumors, and LILRB1-expressing immune cells infiltrate breast cancer tissues, which may contribute to tumor development [62]. Soluble anti-LILRB1 stimulates, but crosslinked antibody inhibits, T cell proliferation [63]. Tumor-cell-expressed HLA-G interacts with LILRB1 on V γ 9V δ 2 T cells to inhibit cytotoxicity of these T cells [64]. Cross-linking of LILRB1 inhibits antigen-induced B cell activation and suppresses antibody production [65]. Interaction of HLA-G with LILRB1 on NK and macrophages inhibits cytotoxicity and inflammation toward trophoblasts, circumventing undesired anti-fetus immune responses during pregnancy [66,67]. In addition, LILRB1, like LILRB3 and 4, inhibits differentiation of osteoclasts [17].

LILRB1 is also expressed on certain AML cells [26], especially in M5 monocytic AML cells, neoplastic B cells (including B-cell leukemia, B-cell lymphoma, and multiple myeloma cells [27,28]), and T cell leukemia and lymphoma cells [29]. LILRB1 expression on primary cutaneous CD8⁺ and CD56⁺ T cell lymphomas may protect these tumor cells from apoptosis [29]. LILRB1 is also expressed in human gastric cancer cells and may enhance tumor growth [30]. In contrast, soluble or nanoparticle-aggregated HLA-G binds LILRB1 on neoplastic B cells and inhibits their proliferation [27]. Antibody blocking of LILRB1 on myeloma or lymphoblastic cells in culture did not change NK cell-mediated cell lysis [31]. Therefore the roles of LILRB1 in different contexts will need further investigations.

3.2 LILRB2 (CD85D, ILT4, LIR2, MIR10)

LILRB2 contains four extracellular immunoglobulin domains, a transmembrane domain, and three cytoplasmic ITIMs. The receptor is expressed on monocytes, macrophages, dendritic cells, hematopoietic stem cells, and the basophils of some individuals but not on lymphoid cells [21,49,68]. LILRB2 was also reported to be expressed by endothelial cells [69], decidual macrophages [59], and osteoclasts [17]. Progenitor mast cells express this receptor whereas mature mast cells do not [60]. LILRB2 binds to multiple types of ligands, including HLA class I molecules [49], CD1d [52], myelin inhibitors (including Nogo66, MAG, and OMgp [53]), Angptls [21,42], and β -amyloid [43]. *In vitro* cross-linking of LILRB2 with Fc γ RI inhibits Fc receptor (FcR)-mediated signaling in monocytes [68] and serotonin release in LILRB2-transfected basophilic leukemia cells [49]. Upregulation of LILRB2 induces dendritic cell tolerance [70]. LILRB2 suppresses axonal regeneration [53] and promotes Alzheimer's disease [43]. We showed that LILRB2 is required for *ex vivo* expansion of hematopoietic stem cells (HSCs) [21]. PirB, the mouse ortholog of human LILRB2 and LILRB3, suppresses myeloid-derived suppressor cell differentiation into M1 macrophages that inhibit regulatory T cell activities and tumor development [45]. PirB expression increases upon myeloid and B cell differentiation [71].

LILRB2 is expressed on various cancerous cells. AML cells, especially monocytic AML cells express LILRB2 [26]. The mouse ortholog PirB is also expressed on MLL-AF9 AML cells, including AML SCs [21]. LILRB2 expression is induced by some chronic lymphoblastic leukemia (CLL) cells [32]. LILRB2 expression is not limited to hematologic malignancies. LILRB2 is expressed in breast cancer cell lines and in 60.7% of primary ductal and lobular breast cancer tissues but is absent from normal breast tissues [33]. The receptor is also expressed on human non-small cell lung cancer cells [34,39–41] and supports their survival and cancer development [39].

3.3 LILRB3 (CD85A, ILT5, LIR3, HL9)

LILRB3 contains four extracellular immunoglobulin domains, a transmembrane domain, and four cytoplasmic ITIMs. LILRB3 is constitutively expressed on monocytes, monocyte-derived osteoclasts, neutrophils, eosinophils, basophils, and osteoclasts [17]. LILRB3 is expressed on progenitor mast cells but not on mature mast cells [60]. Co-ligation of LILRB3 with Fc ϵ RI *in vitro* inhibits Fc receptor-mediated cell activities in human basophils [72]. LILRB3 inhibits differentiation of osteoclasts [17].

LILRB3 is expressed by certain myeloid leukemia, B lymphoid leukemia, and myeloma cells, and is co-expressed with leukemia stem cell marker CD34 or myeloma marker CD138 [35]. We showed that inhibition of LILRB3 expres-

sion in certain human leukemia cell lines inhibited growth of these cells [26]. An anti-LILRB3 antibody induces cytotoxicity of LILRB3-expressing cells via complement-dependent cytotoxicity and antibody-dependent cellular cytotoxicity [35].

3.4 LILRB4 (CD85K, ILT3, LIR5, HM18)

LILRB4 contains two extracellular immunoglobulin domains, a transmembrane domain, and three ITIMs. It is the only LILRB that contains two Ig domains. The receptor is expressed on dendritic cells, monocytes and macrophages [15,73], progenitor mast cells [60], endothelial cells [74], and osteoclasts [17]. LILRB4 is expressed on progenitor mast cells but not on mature mast cells [60]. Like other inhibitory LILRB proteins, the intracellular ITIMs of LILRB4 recruit SHP-1, which transduces a negative signal that inhibits stimulation of an immune response. LILRB4 was shown to inhibit cell activation of monocytes and macrophages [15]. Up-regulation of LILRB4 and LILRB2 expression in tolerogenic antigen presenting cells, cells that may regulate the functions of suppressor and regulatory T cells, leads to immune tolerance [70,75]. Both membrane-bound and soluble LILRB4 induce anergy of T helper cells and differentiation of CD8⁺ T suppressor cells, contributing to rejection of allogeneic tumor transplants [36,76]. LILRB4 inhibits differentiation of osteoclasts [17]. LILRB4 is one of the most polymorphic receptors with at least 15 identified single nucleotide polymorphisms (SNPs) [77].

LILRB4 has been shown to play important roles in cancer development, transplantation, and autoimmune diseases [73]. LILRB4 expression marks monocytic AML cells, and it is co-expressed with stem cell marker c-Kit or CD34 in 50% and 39% of cases, respectively [56]. We showed that inhibition of LILRB4 expression in human leukemia cell lines inhibited growth of these cells [26]. LILRB4 is also expressed on CLL B cells but not normal B cells and may have prognostic value in CLL [32]. LILRB4 is expressed in human gastric cancer cells and may enhance tumor growth by inhibiting NK cell activity [30]. Soluble LILRB4 is present in more than 40% of serum samples from patients with colorectal carcinoma, pancreatic carcinoma, and melanoma, and inhibits T-cell responses *in vitro* [36,37]. The supportive role of LILRB4 in development and/or progression of these cancers was verified by treatment of anti-LILRB4 or by depletion of LILRB4 [36].

3.5 LILRB5 (CD85C, LIR8)

LILRB5 is expressed in subpopulations of monocytes, NK cells, and mast cell granules [12,60]. It was suggested to play a role in clearance of creatine kinase via the mononuclear phagocytic system in the liver [78]. LILRB5 is unique among LILRBs in that it is the only LILRB that is not highly expressed by M5 AML cells and in that its expres-

sion does not correlate with the overall survival of AML patients based on our analysis of TCGA database of AML patients (<https://tcga-data.nci.nih.gov/tcga/>).

3.6 Mechanisms that underlie the roles of LILRB signaling in cancer cells

The signaling mediated by LILRBs starts by the activation of ITIMs via the phosphorylation of tyrosines in these motifs by Src kinases, leading to the recruitment of SH2 domain-containing phosphatases or the inositol-phosphatase SHIP [2–4]. The interaction between phospho-tyrosine in ITIMs and phosphatases are exemplified by an analysis of the mouse PirB [79,80]. The known substrates of ITIM-recruited phosphatases include ITAMs, Src, Syk, ZAP70, Lyn, PI3K, PLC- γ , and Vav1 [6–9]. However, it is likely that LILRBs have additional substrates, and the downstream signaling molecules have not been well characterized. It is also likely that ITIM receptors have diverse signaling branches instead of a linear signaling cascade like that of the janus kinase and signal transducer and activator of transcription (JAK/STAT) pathway. In addition, the LILRBs and different ligands presumably induce distinct signaling networks in a manner that is cell context dependent.

We studied the downstream signaling of PirB, LILRB2, and LAIR1 as representative ITIM-containing receptors in retrovirus transplantation acute leukemia models and in the lung cancer implantation model.

3.7 Signaling downstream of LILRB2 and PirB in cancer cells

We have studied the role of the LILRB2 mouse homolog PirB in MLL-AF9 AML cells and found that PirB is associated with SHP-1 and SHP-2. Defective PirB signaling results in decreased levels of phosphorylation of SHP-1 and SHP-2 in AML cells [21]. We also found that phosphorylation of CAMKIV can be induced by Angptl binding to LILRB2 in human cord blood CD34⁺ cells, and that p-CAMKIV levels are decreased in PirB-deficient AML cells [21]. These results suggest that LILRB2-mediated signaling is transmitted through SHP-1 and SHP-2 and is coupled with CAMKIV in AML cells.

Several papers showed that both Angptl2 and LILRB2 are highly expressed in non-small cell lung cancer (NSCLC) samples, and levels of these proteins are negatively correlated with the patient prognosis [39–41]. In one case, it was shown that ILT4 drives B7-H3 expression via PI3K/ AKT/ mTOR signalling and ILT4/B7-H3 co-expression correlates with poor prognosis in non-small cell lung cancer [41]. In another case, inhibition of LILRB2 resulted in a drastic decrease in proliferation, colony formation, and migration of NSCLC cells, and Angptl2 binding to LILRB2 supports lung cancer development via the SHP2/CaMKI/ CREB axis [39].

These studies suggest that LILRB2 signaling represents a novel target for lung cancer treatment.

3.8 Signaling downstream of LAIR1 in AML cells and B-ALL cells

LAIR1 does not affect normal hematopoiesis but is essential for leukemia development [26,38,57]. LAIR1 is a type I transmembrane glycoprotein that shares the same domain organization as LILRBs, containing one extracellular Ig-like domain that binds collagens or surfactant protein D and two intracellular ITIMs that recruit SHP-1 and SHP-2 [22–25]. LAIR1 is known to be expressed on various lineages of hematopoietic cells and hematopoietic progenitor CD34⁺ cells [55]. Previous studies demonstrated that LAIR1 engagement with antibodies induced apoptosis and prevented NF- κ B nuclear translocation of AML cell lines [81], and also blocked AKT and NF- κ B activation and thus cell division of CLL cells [82]. We showed that LAIR1 is highly expressed on human AML and B cell ALL (B-ALL) cell lines, as well as on primary AML cells especially monocytic M5 AML cells [26]. Inhibition of LAIR1 expression decreases leukemia development *in vitro* and in xenograft experiments [26,38]. In retrovirus transplantation models, including MLL-AF9 (AML) [83,84], AML1-ETO9a (AML) [85], BCR-ABL1 (B-ALL) [38], and N-Myc (B-ALL) [86], LAIR1 deficiency blocks leukemia development in primary or serial transplantation [26,38], suggesting that LAIR1 is critical for maintenance of the activity of AML stem cells [26].

Importantly, we also found that SHP-1, but not SHP-2, mediates LAIR1 signaling in AML cells and prevents exhaustion of AML SCs *in vitro* and *in vivo*. Furthermore, we demonstrated that SHP-1 is a negative signaling molecule for normal myeloid differentiation but acts as a phosphatase-independent adaptor to recruit CAMKI for activation of the downstream transcription factor CREB in AML cells. The LAIR1/SHP-1/CAMKI axis may represent a target for treating AML [26].

Müschen's group [38] demonstrated that LAIR1, as well as other ITIM-containing receptors, supports development and relapse of Philadelphia chromosome-positive (Ph⁺) B-ALL. LAIR1 acts through SHP-1 and SHIP, which dephosphorylate Syk to support this particular type of B-ALL. Hyperactive Syk tyrosine kinase activity is required and sufficient to induce death of these B-ALL cells. This suggests that the basic mechanism of negative selection of B cells with over activation is functional in transformed B-ALL cells. Therefore, activating Syk and a negative B-cell selection strategy may overcome drug resistance in Ph⁺ B-ALL [38].

Together, these results suggest that ITIM-containing receptors support the development, drug resistance, relapse, or cancer stem cell activity of different types of cancer, even

though the downstream signaling differs. When SHP-1 acts as a mediator of the signaling, both the phosphatase-dependent and -independent mechanisms can be used by the cancer cells (Figure 2).

4 Perspectives and future work

Effective elimination of all cancer cells, including cancer stem cells, remains a major challenge in oncology. Our finding that LILRBs and related receptor LAIR1 directly support the survival and self-renewal of leukemia cells opened a new research front in cancer research, diagnosis, and treatment. Below we discuss several research questions of ITIM-containing receptors that should be addressed in the near future.

4.1 Identification of ligands

Cloning of unknown ligands for LILRBs is a key effort to elucidate how these receptors work. The ligands for LILRB3 and LILRB4 have not yet been identified. In addition, based on fact that LILRB2 and LAIR1 each bind to more than one ligand, it is possible that multiple ligands exist for certain ITIM-containing receptors. For example, the binding between LILRB1-2 and MHC-I occurs with low $\mu\text{mol L}^{-1}$ affinity, suggesting the possibility of the existence of additional high-affinity ligands, co-ligands, or binding proteins. To identify LILRB ligands, various methods can be applied including expression cloning, crosslinking followed by co-immunoprecipitation and mass spectroscopy, protein array, and candidate screening. Recently developed




Figure 2 (color online) ITIM-containing receptor downstream signaling in different leukemia cells. In LILRB-expressing AML cells, the ligand binding or interplays among different types of receptors activates LILRB as reflected by the phosphorylation of the tyrosines in ITIMs. SHP-1 is recruited and acts as a phosphatase-independent scaffolding protein to form a complex with the kinase CAMKI. The activated CAMKI then induces phosphorylation of the transcription factor CREB that translocates into nucleus to regulate the self-renewal, survival, and differentiation of AML cells. By contrast, in BCR-ABL⁺ B-ALL cells, BCR-ABL induced phosphorylation of Syk is balanced by the ITIM-containing receptor mediated SHP-1 (or SHIP, not shown here) phosphatase-dependent dephosphorylation. This avoids the hyperphosphorylation of Syk, thus preventing the negative selection of over-activated malignant B cells.

cloning methods employing cell microarrays [87] and ligand-based receptor capture technologies [88] may accelerate the progress.

4.2 Cell type-dependent signaling

A key question is why some inhibitory receptors have supportive effects on cancer development. The tumor-supportive role of SHP-1 in certain acute leukemia provides an explanation. A role for SHP-1 in AML development is supported by evidence from previous studies: human LAIR1 is mainly associated with SHP-1 but not SHP-2 [89]; SHP-1 suppresses differentiation in some leukemia cells [90], concordant with the reported anti-differentiation activity of LAIR1 [91]; and SHP-1 inhibits apoptosis in freshly isolated leukemia cells [92]. While SHP-1 is capable of binding to Grb2 in a phosphatase-independent manner [93], the CAMKI recruitment of SHP-1 represents a different phosphatase-independent mechanism to sustain AML SC activity in AML. By contrast, SHP-1 utilizes a phosphatase-dependent mechanism to support Ph⁺ B-ALL [38].

Furthermore, it is known that SHP-1 is a negative regulator of growth of normal hematopoietic progenitors and that overexpression of SHP-1 inhibits growth of cancer cell lines [94–98]. It was also suggested that LAIR1 signaling negatively regulates myeloid leukemia and CLL [81,82]. Therefore, it will be interesting to determine the cell specificity for ITIM-containing receptors and SHP-1 to exert the tumor-supportive and tumor-suppressive functions.

The potentially overlapping but distinct roles of SHP-1, SHP-2, and SHIP that mediate ITIM-containing receptors' signaling in different cells are largely unknown. It is generally agreed that SHP-2 plays a positive signaling role in the hematopoietic system, whereas SHP-1 is a negative regulator of cell signaling. Therefore, the finding that SHP-1, but not SHP-2, mediates LAIR1 signaling to support AML development is surprising and deserves attention. In addition, SHP-2 is also known to have tumor-specific functions, acting as an oncogene or a tumor-suppressor in different types of cancers [99]. A similarly interesting observation is that, while SHP-1 appears to be solely responsible for LAIR1's tumor-promoting function in AML cells [26], both SHP-1 and SHIP are capable of supporting the development of Ph⁺ B-ALL [38]. Identification of the cell context for SHP-1, SHP-2, and SHIP's signaling roles and their respective substrates or interacting proteins in different cell types will be critical.

Better understanding of the likely divergent downstream signaling of LILRBs in cancer is another area that deserves further investigations. For example, SHP-1 and CAMKI may have downstream effectors other than CREB in LAIR1-expressing AML cells. The identification of the full spectrum of LILRB downstream signaling in various cells is an important task.

4.3 Roles of LILRBs and additional ITIM-containing receptors in solid cancer

The supportive role of LILRBs is unlikely to be limited to acute leukemia. While our *in silico* analyses indicate that expression of a number of ITIM inhibitory receptors negatively correlates with AML patient survival, it is possible that some of these receptors may also play positive roles in development of other types of cancer. Indeed, LILRB1, 2, and 4 are expressed in solid cancer cells such as gastric cancer, breast cancer, and lung cancer cells [30,33,34,39–41]. Because there are numerous types of ITIM-containing receptors, different receptors may have different expression patterns in different types or subtypes of cancers. It will be interesting to determine the individual and combined effects of these receptors in the same cancer cells, and the extent to which LILRB signaling in acute leukemia cells can be generalized to other immune inhibitory receptors and other types of cancer.

4.4 Therapeutic development

The identification of LILRBs and their downstream signaling as potential therapeutic targets may reshape our views regarding how cancer develops, how cancer cells differ from other cells, and how to treat this difficult disease. Our study suggests that some leukemia cells have unique signaling pathways downstream of ITIM-containing receptors. These inhibitory receptors may enable the leukemia cells to survive conventional therapies, resulting in tumor relapse.

Because inhibition of the signaling of certain LILRBs directly blocks cancer growth, and stimulates immunity that may suppress tumorigenesis but does not disturb normal development, these receptors may represent ideal targets for treating cancer.

The blockade of inhibitory receptor signaling in combination with conventional therapies may prove to be an effective strategy for elimination of leukemia cells as well as other types of cancer cells. Inhibition of factors involved in the intracellular signaling will represent a more powerful means of blocking the effects of ITIM-containing receptors. In particular, the identification of the specific inhibitors of the phosphatase-dependent and -independent SHP-1 activity for cancer treatment may have significant benefits. Therapeutic modalities for LILRBs may also include recombinant soluble extracellular domains of these receptors and blocking antibodies against the receptors' signaling. Moreover, it will be interesting to test whether chimeric antigen receptors [100] engineered to target LILRBs and other ITIM-containing receptors are effective in treating certain types of cancer especially hematopoietic malignancies.

The author(s) declare that they have no conflict of interest.

This work was supported by the National Institute of Health

(1R01CA172268), the Leukemia & Lymphoma Society (1024-14, TRP-6024-14), the Robert A. Welch Foundation (I-1834), the Cancer Prevention and Research Institute of Texas (RP140402 and DP150056), the Innovation Program of Shanghai Municipal Education Commission (13G20), the Program for Professor of Special Appointment (Eastern Scholar) at Shanghai Institutions of Higher Learning, and the National Natural Science Foundation of China (81370654, 81422001, 81471524). We regret that we have been unable to cite many relevant primary references due to space limitations.

- Daeron M, Latour S, Malbec O, Espinosa E, Pina P, Pasmans S, Fridman WH. The same tyrosine-based inhibition motif, in the intracytoplasmic domain of Fc gamma RIIB, regulates negatively BCR-, TCR-, and FcR-dependent cell activation. *Immunity*, 1995, 3: 635–646
- Takai T, Nakamura A, Endo S. Role of PIR-B in autoimmune glomerulonephritis. *J Biomed Biotechnol*, 2011, 2011: 275302
- Daeron M, Jaeger S, Du Pasquier L, Vivier E. Immunoreceptor tyrosine-based inhibition motifs: a quest in the past and future. *Immunol Rev*, 2008, 224: 11–43
- Katz HR. Inhibition of inflammatory responses by leukocyte Ig-like receptors. *Adv Immunol*, 2006, 91: 251–272
- Bruhns P, Vely F, Malbec O, Fridman WH, Vivier E, Daeron M. Molecular basis of the recruitment of the SH2 domain-containing inositol 5-phosphatases SHP1 and SHIP2 by fcgamma RIIB. *J Biol Chem*, 2000, 275: 37357–37364
- Huang ZY, Hunter S, Kim MK, Indik ZK, Schreiber AD. The effect of phosphatases SHP-1 and SHIP-1 on signaling by the ITIM- and ITAM-containing Fcgamma receptors FcgammaRIIB and FcgammaRIIA. *J Leukoc Biol*, 2003, 73: 823–829
- Binstadt BA, Brumbaugh KM, Dick CJ, Scharenberg AM, Williams BL, Colonna M, Lanier LL, Kinet JP, Abraham RT, Leibson PJ. Sequential involvement of Lck and SHP-1 with MHC-recognizing receptors on NK cells inhibits FcR-initiated tyrosine kinase activation. *Immunity*, 1996, 5: 629–638
- Daigle I, Yousefi S, Colonna M, Green DR, Simon HU. Death receptors bind SHP-1 and block cytokine-induced anti-apoptotic signaling in neutrophils. *Nat Med*, 2002, 8: 61–67
- Stebbins CC, Watzl C, Billadeau DD, Leibson PJ, Burshtyn DN, Long EO. Vav1 dephosphorylation by the tyrosine phosphatase SHP-1 as a mechanism for inhibition of cellular cytotoxicity. *Mol Cell Biol*, 2003, 23: 6291–6299
- Staub E, Rosenthal A, Hinzmann B. Systematic identification of immunoreceptor tyrosine-based inhibitory motifs in the human proteome. *Cell Signal*, 2004, 16: 435–456
- Ivashkiv LB. Cross-regulation of signaling by ITAM-associated receptors. *Nat Immunol*, 2009, 10: 340–347
- Borges L, Hsu ML, Fanger N, Kubin M, Cosman D. A family of human lymphoid and myeloid Ig-like receptors, some of which bind to MHC class I molecules. *J Immunol*, 1997, 159: 5192–5196
- Cosman D, Fanger N, Borges L, Kubin M, Chin W, Peterson L, Hsu ML. A novel immunoglobulin superfamily receptor for cellular and viral MHC class I molecules. *Immunity*, 1997, 7: 273–282
- Colonna M, Navarro F, Bellon T, Llano M, Garcia P, Samaridis J, Angman L, Cella M, Lopez-Botet M. A common inhibitory receptor for major histocompatibility complex class I molecules on human lymphoid and myelomonocytic cells. *J Exp Med*, 1997, 186: 1809–1818
- Cella M, Dohring C, Samaridis J, Dessing M, Brockhaus M, Lanzavecchia A, Colonna M. A novel inhibitory receptor (ILT3) expressed on monocytes, macrophages, and dendritic cells involved in antigen processing. *J Exp Med*, 1997, 185: 1743–1751
- Martin AM, Kulski JK, Witt C, Pontarotti P, Christiansen FT. Leukocyte Ig-like receptor complex (LRC) in mice and men. *Trends Immunol*, 2002, 23: 81–88
- Mori Y, Tsuji S, Inui M, Sakamoto Y, Endo S, Ito Y, Fujimura S, Koga T, Nakamura A, Takayanagi H, Itoi E, Takai T. Inhibitory immunoglobulin-like receptors LILRB and PIR-B negatively regulate

- osteoclast development. *J Immunol*, 2008, 181: 4742–4751
- 18 Kubagawa H, Burrows PD, Cooper MD. A novel pair of immunoglobulin-like receptors expressed by B cells and myeloid cells. *Proc Natl Acad Sci USA*, 1997, 94: 5261–5266
- 19 Katz HR, Vivier E, Castells MC, McCormick MJ, Chambers JM, Austen KF. Mouse mast cell gp49B1 contains two immunoreceptor tyrosine-based inhibition motifs and suppresses mast cell activation when coligated with the high-affinity Fc receptor for IgE. *Proc Natl Acad Sci USA*, 1996, 93: 10809–10814
- 20 Wende H, Colonna M, Ziegler A, Volz A. Organization of the leukocyte receptor cluster (LRC) on human chromosome 19q13.4. *Mamm Genome*, 1999, 10: 154–160
- 21 Zheng J, Umikawa M, Cui C, Li J, Chen X, Zhang C, Huynh H, Kang X, Silvany R, Wan X, Ye J, Cantó AP, Chen SH, Wang HY, Ward ES, Zhang CC. Inhibitory receptors bind ANGPTLs and support blood stem cells and leukaemia development. *Nature*, 2012, 485: 656–660
- 22 Olde Nordkamp MJ, van Eijk M, Urbanus RT, Bont L, Haagsman HP, Meyaard L. Leukocyte-associated Ig-like receptor-1 is a novel inhibitory receptor for surfactant protein D. *J Leukoc Biol*, 2014, 96: 105–111
- 23 Meyaard L, Adema GJ, Chang C, Woollatt E, Sutherland GR, Lanier LL, Phillips JH. LAIR-1, a novel inhibitory receptor expressed on human mononuclear leukocytes. *Immunity*, 1997, 7: 283–290
- 24 Poggi A, Pella N, Morelli L, Spada F, Revello V, Sivori S, Augugliaro R, Moretta L, Moretta A. p40, a novel surface molecule involved in the regulation of the non-major histocompatibility complex-restricted cytolytic activity in humans. *Eur J Immunol*, 1995, 25: 369–376
- 25 Lebbink RJ, de Ruiter T, Adelmeijer J, Brenkman AB, van Helvoort JM, Koch M, Farndale RW, Lisman T, Sonnenberg A, Lenting PJ, Meyaard L. Collagens are functional, high affinity ligands for the inhibitory immune receptor LAIR-1. *J Exp Med*, 2006, 203: 1419–1425
- 26 Kang X, Lu Z, Cui C, Deng M, Fan Y, Dong B, Han X, Xie F, Tyner JW, Coligan JE, Collins RH, Xiao X, You MJ, Zhang CC. The ITIM-containing receptor LAIR1 is essential for acute myeloid leukaemia development. *Nat Cell Biol*, 2015, 17: 665–677
- 27 Naji A, Menier C, Maki G, Carosella ED, Rouas-Freiss N. Neoplastic B-cell growth is impaired by HLA-G/ILT2 interaction. *Leukemia*, 2012, 26: 1889–1892
- 28 Harly C, Peyrat MA, Netzer S, Dechanet-Merville J, Bonneville M, Scotet E. Up-regulation of cytolytic functions of human Vdelta2-gamma T lymphocytes through engagement of ILT2 expressed by tumor target cells. *Blood*, 2011, 117: 2864–2873
- 29 Urosevic M, Kamarashev J, Burg G, Dummer R. Primary cutaneous CD8+ and CD56+ T-cell lymphomas express HLA-G and killer-cell inhibitory ligand, ILT2. *Blood*, 2004, 103: 1796–1798
- 30 Zhang Y, Lu N, Xue Y, Zhang M, Li Y, Si Y, Bian X, Jia Y, Wang Y. Expression of immunoglobulin-like transcript (ILT)2 and ILT3 in human gastric cancer and its clinical significance. *Mol Med Rep*, 2012, 5: 910–916
- 31 Heidenreich S, Zu Eulenburg C, Hildebrandt Y, Stubig T, Sierich H, Badbaran A, Eiermann TH, Binder TM, Kroger N. Impact of the NK cell receptor LIR-1 (ILT-2/CD85j/LILRB1) on cytotoxicity against multiple myeloma. *Clin Dev Immunol*, 2012, 2012: 652130
- 32 Colovai AI, Tsao L, Wang S, Lin H, Wang C, Seki T, Fisher JG, Menes M, Bhagat G, Alobeid B, Suciufoca N. Expression of inhibitory receptor ILT3 on neoplastic B cells is associated with lymphoid tissue involvement in chronic lymphocytic leukemia. *Cytometry B Clin Cytom*, 2007, 72: 354–362
- 33 Liu J, Wang L, Gao W, Li L, Cui X, Yang H, Lin W, Dang Q, Zhang N, Sun Y. Inhibitory receptor immunoglobulin-like transcript 4 was highly expressed in primary ductal and lobular breast cancer and significantly correlated with IL-10. *Diagn Pathol*, 2014, 9: 85
- 34 Sun Y, Liu J, Gao P, Wang Y, Liu C. Expression of Ig-like transcript 4 inhibitory receptor in human non-small cell lung cancer. *Chest*, 2008, 134: 783–788
- 35 Pfistershammer K, Lawitschka A, Klauser C, Leitner J, Weigl R, Heemskerk MH, Pickl WF, Majdic O, Böhmig GA, Fischer GF, Greinix HT, Steinberger P. Allogeneic disparities in immunoglobulin-like transcript 5 induce potent antibody responses in hematopoietic stem cell transplant recipients. *Blood*, 2009, 114: 2323–2332
- 36 Suciufoca N, Feirt N, Zhang QY, Vlad G, Liu Z, Lin H, Chang CC, Ho EK, Colovai AI, Kaufman H, D'Agati VD, Thaker HM, Remotti H, Galluzzo S, Cinti P, Rabitti C, Allendorf J, Chabot J, Caricato M, Coppola R, Berloco P, Cortesini R. Soluble Ig-like transcript 3 inhibits tumor allograft rejection in humanized SCID mice and T cell responses in cancer patients. *J Immunol*, 2007, 178: 7432–7441
- 37 Cortesini R. Pancreas cancer and the role of soluble immunoglobulin-like transcript 3 (ILT3). *JOP*, 2007, 8: 697–703
- 38 Chen Z, Shojaee S, Buchner M, Geng H, Lee JW, Klemm L, Titz B, Graeber TG, Park E, Tan YX, Satterthwaite A, Paietta E, Hunger SP, Willman CL, Melnick A, Loh ML, Jung JU, Coligan JE, Bolland S, Mak TW, Limnander A, Jumaa H, Reth M, Weiss A, Lowell CA, Müschen M. Signalling thresholds and negative B-cell selection in acute lymphoblastic leukaemia. *Nature*, 2015, 521: 357–361
- 39 Liu X, Yu X, Xie J, Zhan M, Yu Z, Xie L, Zeng H, Zhang F, Chen G, Yi X, Zheng J. ANGPTL2/LILRB2 signaling promotes the propagation of lung cancer cells. *Oncotarget*, 2014, in press
- 40 Wang L, Geng T, Guo X, Liu J, Zhang P, Yang D, Li J, Yu S, Sun Y. Co-expression of immunoglobulin-like transcript 4 and angiopoietin-like proteins in human non-small cell lung cancer. *Mol Med Rep*, 2015, 11: 2789–2796
- 41 Zhang P, Yu S, Li H, Liu C, Li J, Lin W, Gao A, Wang L, Gao W, Sun Y. ILT4 drives B7-H3 expression via PI3K/AKT/mTOR signalling and ILT4/B7-H3 co-expression correlates with poor prognosis in non-small cell lung cancer. *FEBS Lett*, 2015, 589: 2248–2256
- 42 Deng M, Lu Z, Zheng J, Wan X, Chen X, Hirayasu K, Sun H, Lam Y, Chen L, Wang Q, Song C, Huang N, Gao GF, Jiang Y, Arase H, Zhang CC. A motif in LILRB2 critical for Angptl2 binding and activation. *Blood*, 2014, 124: 924–935
- 43 Kim T, Vidal GS, Djuricic M, William CM, Birnbaum ME, Garcia KC, Hyman BT, Shatz CJ. Human LILRB2 is a beta-amyloid receptor and its murine homolog PirB regulates synaptic plasticity in an Alzheimer's model. *Science*, 2013, 341: 1399–1404
- 44 Zhang Z, Hatano H, Shaw J, Olde Nordkamp M, Jiang G, Li D, Kollnberger S. The leukocyte immunoglobulin-like receptor family member LILRB5 binds to HLA-class I heavy chains. *PLoS One*, 2015, 10: e0129063
- 45 Ma G, Pan PY, Eisenstein S, Divino CM, Lowell CA, Takai T, Chen SH. Paired immunoglobulin-like receptor-B regulates the suppressive function and fate of myeloid-derived suppressor cells. *Immunity*, 2011, 34: 385–395
- 46 Hirayasu K, Arase H. Functional and genetic diversity of leukocyte immunoglobulin-like receptor and implication for disease associations. *J Hum Genet*, 2015, doi: 10.1038/jhg.2015.64
- 47 Barrow AD, Trowsdale J. The extended human leukocyte receptor complex: diverse ways of modulating immune responses. *Immunol Rev*, 2008, 224: 98–123
- 48 Willcox BE, Thomas LM, Bjorkman PJ. Crystal structure of HLA-A2 bound to LIR-1, a host and viral major histocompatibility complex receptor. *Nat Immunol*, 2003, 4: 913–919
- 49 Colonna M, Samaridis J, Cella M, Angman L, Allen RL, O'Callaghan CA, Dunbar R, Ogg GS, Cerundolo V, Rolink A. Human myelomonocytic cells express an inhibitory receptor for classical and nonclassical MHC class I molecules. *J Immunol*, 1998, 160: 3096–3100
- 50 Shiroishi M, Tsumoto K, Amano K, Shirakihara Y, Colonna M, Braud VM, Allan DS, Makadzange A, Rowland-Jones S, Willcox B, Jones EY, van der Merwe PA, Kumagai I, Maenaka K. Human inhibitory receptors Ig-like transcript 2 (ILT2) and ILT4 compete with CD8 for MHC class I binding and bind preferentially to HLA-G. *Proc Natl Acad Sci USA*, 2003, 100: 8856–8861
- 51 Arnold V, Cummings JS, Moreno-Nieves UY, Didier C, Gilbert A, Barré-Sinoussi F, Scott-Algara D. S100A9 protein is a novel ligand

- for the CD85j receptor and its interaction is implicated in the control of HIV-1 replication by NK cells. *Retrovirology*, 2013, 10: 122
- 52 Li D, Wang L, Yu L, Freundt EC, Jin B, Screaton GR, Xu XN. Ig-like transcript 4 inhibits lipid antigen presentation through direct CD1d interaction. *J Immunol*, 2009, 182: 1033–1040
- 53 Atwal JK, Pinkston-Gosse J, Syken J, Stawicki S, Wu Y, Shatz C, Tessier-Lavigne M. PirB is a functional receptor for myelin inhibitors of axonal regeneration. *Science*, 2008, 322: 967–970
- 54 Castells MC, Klickstein LB, Hassani K, Cumplido JA, Lacouture ME, Austen KF, Katz HR. gp49B1-alpha(v)beta3 interaction inhibits antigen-induced mast cell activation. *Nat Immunol*, 2001, 2: 436–442
- 55 Meyaard L. LAIR and collagens in immune regulation. *Immunol Lett*, 2010, 128: 26–28
- 56 Dobrowolska H, Gill KZ, Serban G, Ivan E, Li Q, Qiao P, Suci-Foca N, Savage D, Alobeid B, Bhagat G, Colovai AI. Expression of immune inhibitory receptor ILT3 in acute myeloid leukemia with monocytic differentiation. *Cytometry B Clin Cytom*, 2013, 84: 21–29
- 57 Tang X, Tian L, Estes G, Choi SC, Barrow AD, Colonna M, Borrego F, Coligan JE. Leukocyte-associated Ig-like receptor-1-deficient mice have an altered immune cell phenotype. *J Immunol*, 2012, 188: 548–558
- 58 Rojo S, Stebbins CC, Peterson ME, Dombrowicz D, Wagtmann N, Long EO. Natural killer cells and mast cells from gp49B null mutant mice are functional. *Mol Cell Biol*, 2000, 20: 7178–7182
- 59 Hunt JS, Petroff MG, Morales P, Sedlmayr P, Geraghty DE, Ober C. HLA-G in reproduction: studies on the maternal-fetal interface. *Hum Immunol*, 2000, 61: 1113–1117
- 60 Tedla N, Lee CW, Borges L, Gezy CL, Arm JP. Differential expression of leukocyte immunoglobulin-like receptors on cord-blood-derived human mast cell progenitors and mature mast cells. *J Leukoc Biol*, 2008, 83: 334–343
- 61 Shiroishi M, Kuroki K, Ose T, Rasubala L, Shiratori I, Arase H, Tsumoto K, Kumagai I, Kohda D, Maenaka K. Efficient leukocyte Ig-like receptor signaling and crystal structure of disulfide-linked HLA-G dimer. *J Biol Chem*, 2006, 281: 10439–10447
- 62 Lefebvre S, Antoine M, Uzan S, McMaster M, Dausset J, Carosella ED, Paul P. Specific activation of the non-classical class I histocompatibility HLA-G antigen and expression of the ILT2 inhibitory receptor in human breast cancer. *J Pathol*, 2002, 196: 266–274
- 63 Saverino D, Merlo A, Bruno S, Pistoia V, Grossi CE, Ciccone E. Dual effect of CD85/leukocyte Ig-like receptor-1/Ig-like transcript 2 and CD152 (CTLA-4) on cytokine production by antigen-stimulated human T cells. *J Immunol*, 2002, 168: 207–215
- 64 Lesport E, Baudhuin J, Sousa S, LeMaout J, Zamborlini A, Rouas-Freiss N, Carosella ED, Favier B. Inhibition of human gamma delta [corrected] T-cell antitumoral activity through HLA-G: implications for immunotherapy of cancer. *Cell Mol Life Sci*, 2011, 68: 3385–3399
- 65 Merlo A, Tenca C, Fais F, Battini L, Ciccone E, Grossi CE, Saverino D. Inhibitory receptors CD85j, LAIR-1, and CD152 down-regulate immunoglobulin and cytokine production by human B lymphocytes. *Clin Diagn Lab Immunol*, 2005, 12: 705–712
- 66 Ponte M, Cantoni C, Biassoni R, Tradori-Cappai A, Bentivoglio G, Vitale C, Bertone S, Moretta A, Moretta L, Mingari MC. Inhibitory receptors sensing HLA-G1 molecules in pregnancy: decidua-associated natural killer cells express LIR-1 and CD94/NKG2A and acquire p49, an HLA-G1-specific receptor. *Proc Natl Acad Sci USA*, 1999, 96: 5674–5679
- 67 Petroff MG, Sedlmayr P, Azzola D, Hunt JS. Decidual macrophages are potentially susceptible to inhibition by class Ia and class Ib HLA molecules. *J Reprod Immunol*, 2002, 56: 3–17
- 68 Fanger NA, Cosman D, Peterson L, Braddy SC, Maliszewski CR, Borges L. The MHC class I binding proteins LIR-1 and LIR-2 inhibit Fc receptor-mediated signaling in monocytes. *Eur J Immunol*, 1998, 28: 3423–3434
- 69 Manavalan JS, Kim-Schulze S, Scotto L, Naiyer AJ, Vlad G, Colombo PC, Marboe C, Mancini D, Cortesini R, Suci-Foca N. Alloantigen specific CD8+CD28- FOXP3+ T suppressor cells induce ILT3+ ILT4+ tolerogenic endothelial cells, inhibiting alloreactivity. *Int Immunol*, 2004, 16: 1055–1068
- 70 Manavalan JS, Rossi PC, Vlad G, Piazza F, Yarinina A, Cortesini R, Mancini D, Suci-Foca N. High expression of ILT3 and ILT4 is a general feature of tolerogenic dendritic cells. *Transpl Immunol*, 2003, 11: 245–258
- 71 Kubagawa H, Chen CC, Ho LH, Shimada TS, Gartland L, Mashburn C, Uehara T, Ravetch JV, Cooper MD. Biochemical nature and cellular distribution of the paired immunoglobulin-like receptors, PIR-A and PIR-B. *J Exp Med*, 1999, 189: 309–318
- 72 Sloane DE, Tedla N, Awoniyi M, Macglashan DW Jr, Borges L, Austen KF, Arm JP. Leukocyte immunoglobulin-like receptors: novel innate receptors for human basophil activation and inhibition. *Blood*, 2004, 104: 2832–2839
- 73 Vlad G, Chang CC, Colovai AI, Berloco P, Cortesini R, Suci-Foca N. Immunoglobulin-like transcript 3: a crucial regulator of dendritic cell function. *Hum Immunol*, 2009, 70: 340–344
- 74 Kim-Schulze S, Seki T, Vlad G, Scotto L, Fan J, Colombo PC, Liu J, Cortesini R, Suci-Foca N. Regulation of ILT3 gene expression by processing of precursor transcripts in human endothelial cells. *Am J Transplant*, 2006, 6: 76–82
- 75 Chang CC, Ciubotariu R, Manavalan JS, Yuan J, Colovai AI, Piazza F, Lederman S, Colonna M, Cortesini R, Dalla-Favera R, Suci-Foca N. Tolerization of dendritic cells by T_S cells: the crucial role of inhibitory receptors ILT3 and ILT4. *Nat Immunol*, 2002, 3: 237–243
- 76 Kim-Schulze S, Scotto L, Vlad G, Piazza F, Lin H, Liu Z, Cortesini R, Suci-Foca N. Recombinant Ig-like transcript 3-Fc modulates T cell responses via induction of Th anergy and differentiation of CD8+ T suppressor cells. *J Immunol*, 2006, 176: 2790–2798
- 77 Chang CC, Silvia EA, Ho EK, Vlad G, Suci-Foca N, Vasilescu ER. Polymorphism and linkage disequilibrium of immunoglobulin-like transcript 3 gene. *Hum Immunol*, 2008, 69: 284–290
- 78 Dubé MP, Zetler R, Barhdadi A, Brown AM, Mongrain I, Normand V, Laplante N, Asselin G, Zada YF, Provost S, Bergeron J, Kouz S, Dufour R, Diaz A, de Denus S, Turgeon J, Rhéaume E, Phillips MS, Tardif JC. CKM and LILRB5 are associated with serum levels of creatine kinase. *Circ Cardiovasc Genet*, 2014, 7: 880–886
- 79 Blery M, Kubagawa H, Chen CC, Vely F, Cooper MD, Vivier E. The paired Ig-like receptor PIR-B is an inhibitory receptor that recruits the protein-tyrosine phosphatase SHP-1. *Proc Natl Acad Sci USA*, 1998, 95: 2446–2451
- 80 Maeda A, Kurosaki M, Ono M, Takai T, Kurosaki T. Requirement of SH2-containing protein tyrosine phosphatases SHP-1 and SHP-2 for paired immunoglobulin-like receptor B (PIR-B)-mediated inhibitory signal. *J Exp Med*, 1998, 187: 1355–1360
- 81 Poggi A, Pellegatta F, Leone BE, Moretta L, Zocchi MR. Engagement of the leukocyte-associated Ig-like receptor-1 induces programmed cell death and prevents NF-kappaB nuclear translocation in human myeloid leukemias. *Eur J Immunol*, 2000, 30: 2751–2758
- 82 Poggi A, Catellani S, Bruzzone A, Caligaris-Cappio F, Gobbi M, Zocchi MR. Lack of the leukocyte-associated Ig-like receptor-1 expression in high-risk chronic lymphocytic leukaemia results in the absence of a negative signal regulating kinase activation and cell division. *Leukemia*, 2008, 22: 980–988
- 83 Krivtsov AV, Twomey D, Feng Z, Stubbs MC, Wang Y, Faber J, Levine JE, Wang J, Hahn WC, Gilliland DG, Golub TR, Armstrong SA. Transformation from committed progenitor to leukaemia stem cell initiated by MLL-AF9. *Nature*, 2006, 442: 818–822
- 84 Somervaille TC, Cleary ML. Identification and characterization of leukemia stem cells in murine MLL-AF9 acute myeloid leukemia. *Cancer Cell*, 2006, 10: 257–268
- 85 Yan M, Kanbe E, Peterson LF, Boyapati A, Miao Y, Wang Y, Chen IM, Chen Z, Rowley JD, Willman CL, Zhang DE. A previously unidentified alternatively spliced isoform of t(8;21) transcript promotes leukemogenesis. *Nat Med*, 2006, 12: 945–949
- 86 Sugihara E, Shimizu T, Kojima K, Onishi N, Kai K, Ishizawa J,

- Nagata K, Hashimoto N, Honda H, Kanno M, Miwa M, Okada S, Andreeff M, Saya H. Ink4a and Arf are crucial factors in the determination of the cell of origin and the therapeutic sensitivity of Myc-induced mouse lymphoid tumor. *Oncogene*, 2012, 31: 2849–2861
- 87 Turner L, Lavstsen T, Berger SS, Wang CW, Petersen JE, Avril M, Brazier AJ, Freeth J, Jespersen JS, Nielsen MA, Magistrado P, Lusingu J, Smith JD, Higgins MK, Theander TG. Severe malaria is associated with parasite binding to endothelial protein C receptor. *Nature*, 2013, 498: 502–505
- 88 Frei AP, Jeon OY, Kilcher S, Moest H, Henning LM, Jost C, Plückthun A, Mercer J, Aebersold R, Carreira EM, Wollscheid B. Direct identification of ligand-receptor interactions on living cells and tissues. *Nat Biotechnol*, 2012, 30: 997–1001
- 89 Xu M, Zhao R, Zhao ZJ. Identification and characterization of leukocyte-associated Ig-like receptor-1 as a major anchor protein of tyrosine phosphatase SHP-1 in hematopoietic cells. *J Biol Chem*, 2000, 275: 17440–17446
- 90 Eklund EA, Goldenberg I, Lu Y, Andrejic J, Kakar R. SHP1 protein-tyrosine phosphatase regulates HoxA10 DNA binding and transcriptional repression activity in undifferentiated myeloid cells. *J Biol Chem*, 2002, 277: 36878–36888
- 91 Verbrugge A, de Ruiter T, Geest C, Coffey PJ, Meyskens L. Differential expression of leukocyte-associated Ig-like receptor-1 during neutrophil differentiation and activation. *J Leukoc Biol*, 2006, 79: 828–836
- 92 Tibaldi E, Brunati AM, Zonta F, Frezzato F, Gattazzo C, Zambello R, Gringeri E, Semenzato G, Pagano MA, Trentin L. Lyn-mediated SHP-1 recruitment to CD5 contributes to resistance to apoptosis of B-cell chronic lymphocytic leukemia cells. *Leukemia*, 2011, 25: 1768–1781
- 93 Minoo P, Zadeh MM, Rottapel R, Lebrun JJ, Ali S. A novel SHP-1/Grb2-dependent mechanism of negative regulation of cytokine-receptor signaling: contribution of SHP-1 C-terminal tyrosines in cytokine signaling. *Blood*, 2004, 103: 1398–1407
- 94 Tapley P, Shevde NK, Schweitzer PA, Gallina M, Christianson SW, Lin IL, Stein RB, Shultz LD, Rosen J, Lamb P. Increased G-CSF responsiveness of bone marrow cells from hematopoietic cell phosphatase deficient viable motheaten mice. *Exp Hematol*, 1997, 25: 122–131
- 95 Jiao H, Yang W, Berrada K, Tabrizi M, Shultz L, Yi T. Macrophages from motheaten and viable motheaten mutant mice show increased proliferative responses to GM-CSF: detection of potential HCP substrates in GM-CSF signal transduction. *Exp Hematol*, 1997, 25: 592–600
- 96 Wu C, Guan Q, Wang Y, Zhao ZJ, Zhou GW. SHP-1 suppresses cancer cell growth by promoting degradation of JAK kinases. *J Cell Biochem*, 2003, 90: 1026–1037
- 97 Lorenz U, Bergemann AD, Steinberg HN, Flanagan JG, Li X, Galli SJ, Neel BG. Genetic analysis reveals cell type-specific regulation of receptor tyrosine kinase c-Kit by the protein tyrosine phosphatase SHP1. *J Exp Med*, 1996, 184: 1111–1126
- 98 Paulson RF, Vesely S, Siminovitch KA, Bernstein A. Signalling by the W/Kit receptor tyrosine kinase is negatively regulated *in vivo* by the protein tyrosine phosphatase Shp1. *Nat Genet*, 1996, 13: 309–315
- 99 Bard-Chapeau EA, Li S, Ding J, Zhang SS, Zhu HH, Princen F, Fang DD, Han T, Bailly-Maitre B, Poli V, Varki NM, Wang H, Feng GS. Ptpn11/Shp2 acts as a tumor suppressor in hepatocellular carcinogenesis. *Cancer Cell*, 2011, 19: 629–639
- 100 Gill S, June CH. Going viral: chimeric antigen receptor T-cell therapy for hematological malignancies. *Immunol Rev*, 2015, 263: 68–89

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.