

Progress in the study of mercury methylation and demethylation in aquatic environments

LI YanBin & CAI Yong*

Department of Chemistry & Biochemistry, Southeast Environmental Research Center, Florida International University, Miami, Florida 33199, USA

Received March 23, 2012; accepted June 12, 2012; published online August 16, 2012

Mercury (Hg) and its compounds are a class of highly toxic and pervasive pollutants. During the biogeochemical cycling of Hg, methylmercury (MeHg), a potent neurotoxin, can be produced and subsequently bioaccumulated along the food chain in aquatic ecosystems. MeHg is among the most widespread contaminants that pose severe health risks to humans and wildlife. Methylation of inorganic mercury to MeHg and demethylation of MeHg are the two most important processes in the cycling of MeHg, determining the levels of MeHg in aquatic ecosystems. This paper reviews recent progress on the study of Hg methylation and demethylation in aquatic environments, focusing on the following three areas: (1) sites and pathways of Hg methylation and demethylation, (2) bioavailability of Hg species for methylation and demethylation, and (3) application of isotope addition techniques in quantitatively estimating the net production of MeHg.

mercury cycling, methylation, demethylation, bioavailability, methyl mercury

Citation: Li Y B, Cai Y. Progress in the study of mercury methylation and demethylation in aquatic environments. *Chin Sci Bull*, 2013, 58: 177–185, doi: 10.1007/s11434-012-5416-4

Mercury (Hg) and its compounds are a class of highly toxic and pervasive pollutants. Over the past several decades, Hg contaminating and poisoning incidents have been reported in many countries (e.g. Japan, America, Canada, Sweden, and Norway) due to the large amount of Hg emitted into the environment, in particular from anthropogenic sources [1]. It was estimated that about 630000 infants born every year in the United States had an unsafe level of mercury in their blood [2]. Hg contamination is a global problem as one of the major Hg species, elemental mercury (Hg^0) has an approximate atmospheric residence time of 6 month to 1 year [3] and therefore it can be transported to ecosystems that are far from the point sources through long range atmospheric transport.

There are five major species of Hg in the environment, including Hg^0 , divalent inorganic mercury (Hg^{2+}), monomethylmercury (MeHg), dimethylmercury (DMeHg), and monoethylmercury (EtHg). Inorganic Hg (Hg^0 and Hg^{2+}), in

particular Hg^0 , are the major chemical forms of Hg input into the environment from anthropogenic or natural sources [4,5]. During the biogeochemical cycling of Hg, organic mercury species (MeHg, DMeHg, and EtHg) can be produced. Although MeHg only accounts for a small fraction of Hg in the environment, it is the species of most concern to humans due to its high toxicity, prevalent existence, and capability of being accumulated and amplified along the food chain. The bioaccumulation of MeHg in the food chain, in particular in aquatic ecosystems, has caused the exposure of both humans and wildlife to MeHg, posing severe health risks [6] and stressing the importance of understanding the key processes controlling the levels of MeHg in aquatic environments.

In most aquatic ecosystems, *in situ* production (methylation of inorganic Hg to MeHg), rather than input from runoff water or atmospheric deposition, is the major source of MeHg. In addition to methylation of inorganic Hg, the reverse process, demethylation of MeHg, simultaneously occurs in the environment. Both processes are important for

*Corresponding author (email: cai@fiu.edu)

MeHg cycling, determining the levels of MeHg in aquatic environments. This paper reviews recent progress on the study of Hg methylation and demethylation, focusing on the following three areas: (1) sites and pathways of Hg methylation and demethylation, (2) bioavailability of Hg species for methylation and demethylation, and (3) application of isotope addition techniques in quantitatively estimating the production and degradation of MeHg in aquatic ecosystems.

1 Sites of Hg methylation and demethylation in aquatic environments

1.1 Methylation

Adequate attention was not given to the methylation of inorganic Hg in aquatic environments until researchers found that the major species of Hg present in fish was MeHg, rather than inorganic mercury or phenyl mercury, which were the species of Hg in industrial waters [7]. By adding HgCl_2 to bottom sediments and incubating for 5–10 d, Jensen and Jernelov firstly found that Hg^{2+} could be methylated to MeHg in sediments of aquatic ecosystems [7]. Since then, a large number of studies have suggested that methylation in sediment was the primary source of MeHg in most aquatic environments [8–11]. A good positive relation was often observed between MeHg concentration and *in situ* methylation potential in sediments of aquatic systems [12–16]. In addition to sediment, Hg methylation has been also observed in water (both fresh water [17–19] and marine water

[20,21]) and periphyton (floating mat or macrophyte associated) [22–29] (Figure 1) in recent years. As only a small proportion of MeHg produced in sediment can be transported to water column, methylation of Hg in compartments of water column (water and periphyton) may also contribute significantly to the MeHg pool in pelagic food webs. A recent study showed that methylation in water column can be an important source of MeHg in the Arctic, accounting for around 47% of MeHg present in Arctic waters [21]. Methylation of mercury in periphyton may facilitate the bioaccumulation of MeHg in the food chain since periphyton can be the base of food webs in aquatic ecosystems [22]. However, the relative importance of periphyton or water methylation versus sediment methylation in MeHg levels in water column has yet to be clear. This is mainly due to the lack of quantitative estimation and comparison of the net production of MeHg in water column (periphyton and/or water) and the amount of MeHg diffused from sediment to water. Such estimations are necessary for quantitative evaluation of the importance of methylation in periphyton and water to the cycling of MeHg in aquatic ecosystems.

1.2 Demethylation

MeHg demethylation is a reverse process of Hg methylation. It simultaneously occurs in the sites (sediment, periphyton, and water) that methylation takes place (Figure 1). However, methylation is supposed to overrule the demethylation process and determine the net production of MeHg in sediment




Figure 1 (Color online) Cycling of mercury in aquatic environments.

and periphyton. This opinion is supported by the phenomenon that MeHg concentrations were often positively correlated to the methylation potentials [12–16], other than the demethylation potentials in these compartments. Unlike in sediment and periphyton, demethylation in water, in particular photodemethylation was suggested to be more important than the methylation process [30–32]. The importance of demethylation in water was firstly reported in the mid-1990s by Sellers et al. [30] They found that MeHg in water could be rapidly demethylated and about 83% of MeHg flowing into the lake would be removed by photodemethylation. A number of recent studies using various means have confirmed the importance of photodemethylation in a wide range of aquatic ecosystems. Based on measuring the rates of photodemethylation in water, this process was estimated to account for about 80% and 31.4% of the MeHg mobilized annually from sediments in a lake [31] and a wetland [32], respectively. In addition, a good inverse relationship was observed between MeHg concentrations and photodemethylation potentials in water column [32]. Mass independent fractionation (MIF) of Hg isotopes in fish could also be used to quantify the loss of MeHg by photodemethylation as this process leads to significant MIF of odd-mass isotopes [33]. About 25%–68% of MeHg in lakes were estimated to be lost via photodemethylation by measuring MIF in fish [33]. In addition, a study [34] on the diurnal cycling of methylmercury in a wetland found that dissolved MeHg concentrations in water consistently decreased during daylight periods and increased during non-daylight periods. These studies on MeHg demethylation have suggested that MeHg photodemethylation in water can be a major sink of MeHg in various aquatic ecosystems.

2 Pathways of Hg methylation and demethylation in aquatic environments

2.1 Methylation

Inorganic Hg can be methylated to MeHg in water phase through biotic pathways [35–50] or abiotic pathways (photo-mediated or non photo-mediated chemical methylation) [51–66]. In natural aquatic environments, biotic process was generally suggested to be the dominant pathway of Hg methylation due to the fact that sterilization procedure could prohibit the methylation of Hg [7,67,68]. Sulfate reducing bacteria (SRB) were deemed to be responsible for the methylation of Hg^{2+} in most aquatic systems [69–73]. Methylation of Hg^{2+} by SRB in aquatic sediments was firstly confirmed by Compeau and Bartha [74]. In that study, they found that Hg^{2+} methylation in anoxic salt marsh sediment decreased more than 95% in presence of sodium molybdate, a specific inhibitor of sulfate reducers. They isolated a strain of SRB, *Desulfovibrio desulfuricans*, from the sediment and found that it could vigorously methylate Hg^{2+} . Since then, SRB has been proposed to dominate the methylation of Hg

in a variety of aquatic ecosystems based on the sodium molybdate inhibition [22,26,40,69] or good relations observed between sulfate reducing rates and Hg methylation rates [70]. Other microorganisms may also dominate the methylation of Hg, although this may only occur in limited aquatic ecosystems. For instance, iron-reducing bacteria were found to have the capability of methylating Hg^{2+} in both natural sediment [75] and pure culture [76]. A recent study [27] found that methylation rates in periphyton were totally inhibited by a methanogenesis inhibitor and highly stimulated by molybdate, indicating the involvement of methanogens in Hg methylation.

Many chemicals, e.g., amino acids [77], humic substances [53,55,58], silicones [78], and low molecular weight organic acids [66], were found to be able to methylate inorganic mercury in lab settings. Irradiance is necessary for some of these reactions [66,77]. However, importance of these reactions has yet to be confirmed in natural environments. In a recent study, methylation of Hg^{2+} by dissolved organic matter (DOM) under sunlight was proposed to occur in lake waters and cause the increase in MeHg concentration during sunlight hours [19]. However, photomethylation or other chemical pathways of Hg methylation was rarely found to contribute significantly to MeHg production in other ecosystems. A recent study [79] indicated that methylation of Hg under sunlight played a minor role in a wetland as its rate was much slower than that of MeHg photodemethylation. Most previous studies only focused on the methylation of Hg^{2+} in the environment, while there is a lack of knowledge on the methylation of Hg^0 . Our recent study implied that photomethylation of Hg^0 by CH_3I can occur in Florida Everglades water and a pond water (Yin et al., in submission). Contribution of Hg^0 methylation to MeHg pool also needs to be considered as Hg^{2+} can provide sufficient and continuous source of Hg^0 through photo-mediated reduction in natural water.

2.2 Demethylation

Similar to methylation of Hg, demethylation of MeHg can also proceed through biotic [8,80–82] or abiotic pathways (photodemethylation [30–32,34,83–91] or non photo-mediated demethylation [92,93]). Biotic process was suggested to be the dominant pathway of Hg demethylation in sediment and periphyton. Both SRB and methanogens could be the primary microorganisms for this process [27,80,81,94]. Although microbial demethylation of MeHg was also observed to occur in water column [18,20,95], photodemethylation was commonly deemed to be the most important in water column [30–32,89]. Non photo-mediated demethylation of MeHg can also occur in lab settings (e.g., by selenoamino acids [92,93]), however this process has not been confirmed in natural environments.

Since the initial demonstration of the importance of MeHg photodemethylation in aquatic ecosystems [30], a

number of studies have attempted to investigate the mechanisms underlying this process [90,91]. UV radiations (UV-A and UV-B) have been confirmed to be the primary driver of MeHg photodegradation in both a northern temperate lake and a subtropical wetland [32,89]. However, the chemical processes governing MeHg photodemethylation remain unclear. By reviewing the pathways of Hg photoreactions in the literature, five potential pathways could be responsible for MeHg photodemethylation (Figure 2). Although direct photodemethylation of MeHg species such as MeHgOH and MeHgCl (Pathway 1) is theoretically possible, recent studies showed that MeHg present in DI water cannot be demethylated under sunlight [90,91]. Free oxygen radical-induced degradation of MeHg (Pathway 2 or 3) was speculated to be the primary pathway of MeHg photodegradation in previous studies [87,88,90,91]. Both hydroxyl radical ($\cdot\text{OH}$) and singlet oxygen ($^1\text{O}_2$) have been proposed to dominate the photodemethylation of MeHg in aquatic ecosystems. By adding scavengers of singlet oxygen, superoxide anion, hydrogen peroxide and hydroxyl radical, Suda et al. [85] found that singlet oxygen could be responsible for the degradation of MeHg and EtHg in sea water. Another study suggested that $^1\text{O}_2$ induced degradation of methylmercury-DOM complexes (Pathway 2 in Figure 2) was the major pathway of MeHg photodemethylation [91]. A recent work in Arctic lakes found that MeHg photodemethylation was driven by hydroxyl radical produced from photo-Fenton reactions [90]. Some researchers proposed that Hg^0 is the product of MeHg photodemethylation [33,88], while the others reported Hg^{2+} as the main product of this reaction [87,91]. If Hg^0 is the main product in the aquatic environment, photodemethylation of MeHg could also be treated as a reduction process of Hg. Direct

transfer of electrons from photosensitized DOM to Hg within Hg-DOM complexes have been proposed to be one of the possible pathways for Hg^{2+} photoreduction [96]. Thus, direct degradation of DOM-MeHg complexes (Pathway 4) may also be a possible pathway of MeHg photodemethylation in some aquatic ecosystems. A recent study using DOM isolated from natural waters [97] found that MeHg photodemethylation rate was not significantly decreased after adding scavengers of $\cdot\text{OH}$ and $^1\text{O}_2$, implying that Pathway 4 may be the dominant pathway in this ecosystem. These results indicate that the pathways of MeHg photodemethylation may vary in different aquatic ecosystems, as evidenced by inconsistent results reported in previous studies [97]. The variation of MeHg photodemethylation pathway in different aquatic systems may be caused by their differences in chemical characteristics, e.g., DOM. A recent study showed that MIF of MeHg by photodemethylation was significantly affected by the amount of reduced organic sulfur [98], implying that concentrations and characterization of DOM may play an important role in determining the dominant pathway of MeHg photodemethylation in water.

3 Bioavailability of Hg species for Hg methylation and demethylation

Bioavailability of Hg species is among the most important factors that determine the production of MeHg in aquatic ecosystems. Previous studies showed that measured Hg methylation rates were usually positively correlated not to the total Hg^{2+} concentrations, but to the calculated concentrations of Hg^{2+} available for methylation [99–101]. It is also known that the newly deposited Hg^{2+} are more bioavailable for methylation compared to the legacy Hg^{2+} in sediment [102]. The bioavailability of Hg species in aquatic ecosystems is mainly determined by two processes: (1) distribution of Hg between solid and aqueous phase, and (2) speciation of Hg species in water phase (Figure 3).

Distribution of Hg species between solid and aqueous phase is expected to significantly affect the bioavailability of Hg species as only dissolved Hg species can transport through cell membranes and subsequently be methylated or demethylated. The importance of adsorption/desorption on Hg bioavailability has been confirmed by both laboratory and field studies. When *Pseudomonas fluorescens* was tested for its ability to methylate dissolved Hg^{2+} and Hg^{2+} absorbed on mineral colloids, the results showed that methylation rate of dissolved Hg^{2+} was much larger than that of Hg^{2+} absorbed on solids [103]. In addition, Hg methylation rates in surface sediments were observed to be inversely related to the distribution coefficients ($K(D)$) of Hg^{2+} and positively correlated to concentrations of Hg^{2+} in pore waters in some aquatic ecosystems [99,100]. The adsorption/desorption of Hg species is known to be affected by a variety of factors in aquatic environments, e.g., pH, redox potential, salinity,




Figure 2 Possible pathways of MeHg photodemethylation in aquatic environments. By reviewing the pathways of Hg photoreactions in the literature, we proposed that five potential pathways could be responsible for MeHg photodemethylation: (1) direct photodemethylation of MeHg, (2) photodemethylation of MeHg-DOM complexes by free radicals, (3) photodemethylation of MeHg by free radicals, (4) direct photodemethylation of MeHg-DOM complexes.




Figure 3 Hg²⁺ and MeHg complexes present in aquatic environments. Bold: species that have been reported to be the major species available for biotic methylation and demethylation. Italic: species that have been reported to be the major Hg species available for photodemethylation.

organic and inorganic complexing reagents, composition of solid. These factors are also expected to play an important role in controlling the bioavailability of Hg²⁺ in aquatic environments. Unlike the biotic methylation and demethylation, photodemethylation rates of MeHg were observed to be similar in filtrated and un-filtrated waters [89], implying that adsorption/desorption of MeHg in water column plays a minor role in the photodemethylation of MeHg.

Hg²⁺ and MeHg in aquatic environments are generally not free ions, but complexed to various inorganic or organic ligands, including hydroxide, chloride, sulfides, and DOM [104]. The dominant species of Hg²⁺ and MeHg in water or porewater depends upon various physical and chemical parameters, e.g., pH, Eh, sulfide, Cl⁻, and DOM [104–108]. As not all of these complexes are available for methylation and demethylation, speciation of Hg species in water and porewater is another important factor determining bioavailability of Hg species. Neutral Hg-sulfide complexes (e.g. HgS(0)_(aq) and Hg(SH)₂) have been suggested to be the major species of Hg that are available for methylation since the late 1990s [101,105,109–112]. It was proposed that these neutral Hg species could be transported to microorganism cells by passive diffusion and then methylated [109–110]. This opinion was supported by the positive relationship between Hg methylation rates and calculated concentrations of neutral Hg-sulfide complex in solution, which was observed both in laboratory pure culture of SRB [101,111] and field investigations [112,113]. Hg²⁺-thiol complexes are another Hg²⁺ species that have been proposed to be available for methylation [114,115]. In a recent study, methylation of Hg²⁺ by *Geobacter sulfurreducens* was found to be greatly enhanced in the presence of low concentrations of

cysteine. The authors proposed that the formation of Hg²⁺-cysteine complexes, which could be transported to cells by active transport [115], facilitated both the uptake and methylation of Hg²⁺ [114]. However, the importance of Hg-thiol complexes in the methylation of Hg has yet to be confirmed in natural environments. As mercury in aquatic ecosystems is expected to preferentially bind with thiol and other sulfur-containing groups in organic matter [116], it is necessary to investigate whether this process plays an important role in Hg methylation in natural environments.

Similar to the biotic methylation and demethylation of Hg, photodemethylation of MeHg can also be affected by the speciation of MeHg in water column. A previous study reported that sunlight-induced MeHg photodegradation could not occur when MeHgCl, MeHgOH or MeHg ion was the dominant species of MeHg in water, while phenyl and sulfur bonded MeHg species could be decomposed [117]. A recent study suggested that methylmercury species bound to sulphur-containing ligands such as glutathione and mercaptoacetate had a much higher demethylation rates than methylmercury-chloride complexes [91]. Despite these findings in laboratory studies, it is still unclear which species of MeHg dominate the photodemethylation process in natural waters and more attention should be paid to this area.

4 Application of isotope addition technique in the study of Hg methylation and demethylation and estimating the net production of MeHg in aquatic environments

Since the finding of the importance of methylation and demethylation in aquatic ecosystems, efforts have been made to precisely measure the Hg methylation and demethylation rates, which are important for quantitatively estimating the production of MeHg. Isotope addition technique has been applied in this field since the late 1970s [17,118] due to its high accuracy and precision, short incubation time, and ability of simultaneously determining the methylation and demethylation rates. Both radio [17,118–120] and stable isotope addition methods [15,20,32,89,102,121–128] have been applied in this field. When radio isotope addition method was adopted, a radio isotope of Hg²⁺ (e.g., ²⁰³Hg²⁺) is spiked into samples to monitor the methylation process. C-14 labeled MeHg is usually used to examine the demethylation process. Stable isotope addition methods have been widely using in studying Hg methylation and demethylation since the mid-1990s [122]. By using this technique, stable isotope labeled Hg²⁺ (e.g., ¹⁹⁹Hg²⁺) and/or MeHg (e.g., Me²⁰¹Hg) were added into samples to monitor methylation and/or demethylation process, respectively. Although both techniques promise a precise measurement of Hg methylation and demethylation rates, application of radio isotope addition techniques are limited due to its disadvantage of utilizing highly radioactive material and usually requiring

specific safety measures [129].

Net MeHg production rate is a crucial parameter for estimating the net production of MeHg in various compartments of aquatic ecosystems, which is necessary for identifying the major source and sink of MeHg in aquatic ecosystems. This parameter can be calculated by incubating the samples without the addition of any Hg species and measuring the changes in ambient MeHg. However, changes in ambient MeHg are usually difficult to be accurately measured as variation in ambient MeHg is often too small to be detected [119]. By utilizing isotope addition techniques, a method based on measuring the specific methylation and demethylation rates has been developed and used to estimate the net MeHg production rate [14,27,130]. In these studies, methylation (k_m) and demethylation (k_d) rate constants were measured and calculated (eqs. (1) and (2)) by double isotope addition methods. Then, the net production rates of MeHg were calculated by the difference of potential methylation rate ($k_m \times C_{\text{Hg}^{2+}}$ (ambient) and potential demethylation rate ($k_d \times C_{\text{MeHg}}$ (ambient)) (eq. (3)). However, such calculation does not take consideration of the differences in bioavailability of ambient and newly spiked Hg species [102]. A significant error could occur with this omission. For example, a recent study showed that the net production rate of MeHg in Everglades sediment would be overestimated by a factor of 20 without the consideration of the difference of newly spiked and ambient Hg species in methylation and demethylation efficiency [79]. Thus, it is necessary to take consideration of this difference (α and β in eq. (4)) when calculating the net production of MeHg using isotope addition techniques. In previous studies [102], α and β were obtained by measuring the methylation and demethylation rates of ambient and newly spiked Hg species, which is also based on the accurate measurement of changes in ambient MeHg concentrations. Due to the difficulty of accurate measuring the variation in ambient MeHg, it is necessary to develop a method that can accurately measure the differences of ambient and newly spiked mercury species in methylation and demethylation efficiency.

$$k_m = \frac{\Delta C_{\text{Me}^n\text{Hg}}}{C_{\text{Hg}^{2+}} \times t}, \quad (1)$$

$$k_d = \frac{\ln \frac{C_{\text{Me}^n\text{Hg}}}{C_{\text{Me}^n\text{Hg}} - \Delta C_{\text{Me}^n\text{Hg}}}}{t}, \quad (2)$$

$$\text{Net MeHg production rate (ng g}^{-1} \text{ d}^{-1}) = k_m \times C_{\text{Hg}^{2+}} \text{ (ambient)} - k_d \times C_{\text{MeHg}} \text{ (ambient)}, \quad (3)$$

$$\text{Net MeHg production rate (ng g}^{-1} \text{ d}^{-1}) = k_m \times \alpha \times C_{\text{Hg}^{2+}} \text{ (ambient)} - k_d \times \beta \times C_{\text{MeHg}} \text{ (ambient)}, \quad (4)$$

where k_m is the specific methylation rate constant of spiked $\text{Me}^n\text{Hg}^{2+}$ (d^{-1}); k_d is the specific demethylation rate constant of

spiked Me^nHg (d^{-1}); t is the incubation time (d); $C_{\text{Hg}^{2+}}$ and $C_{\text{Me}^n\text{Hg}}$ are the spiked concentrations of Hg^{2+} and Me^nHg (ng g^{-1}), respectively; α is the ratio of methylation rate constant of ambient to newly spiked Hg^{2+} ; β is the ratio of demethylation rate constant of ambient to newly spiked MeHg.

5 Concluding remarks

Methylation and demethylation of mercury are a significant part of MeHg cycling, determining the levels of MeHg in aquatic ecosystems. They can occur in various compartments of aquatic ecosystems, including sediment, periphyton and water. Biotic methylation in sediment and photodemethylation in water are suggested to be the major source and sink of MeHg in aquatic environments. Recent studies found that biotic methylation in periphyton and water could also contribute significantly to the MeHg pool, in particular for the pelagic food webs. The relative importance of methylation in periphyton and/or water versus sediment in MeHg levels in pelagic food webs has yet to be clear. This is mainly due to the lack of quantitative estimation and comparison of the net production of MeHg in water column (periphyton and/or water) and the amount of MeHg diffused from sediment to water. Such estimations are necessary for evaluating the importance of methylation in periphyton and water to the cycling of MeHg in aquatic ecosystems.

Much work is also required in the study of bioavailability of Hg species for methylation and demethylation. Distribution of Hg between solid and water and speciation of Hg species in aqueous phase are the two factors that determine the bioavailability of Hg species. Both neutral Hg^{2+} -sulfide complexes and Hg^{2+} -thiol complexes have been proposed as the species of Hg^{2+} that can be transported to bacteria cells, and subsequently being methylated. More work is required to identify which is the dominant species of Hg^{2+} that associate with the biotic methylation of Hg in natural environments. In comparison to Hg^{2+} , bioavailability of MeHg for both biotic demethylation and photodemethylation has been poorly studied and requires more attentions.

More efforts should also be made on estimating the net MeHg production rate, which is important for identifying the major source of MeHg in aquatic ecosystems. Isotope addition technique is a useful technique in this area due to its high accuracy and precision, short incubation time, and ability of simultaneously determining the methylation and demethylation rates. The difference of newly spiked and ambient mercury species in methylation and demethylation efficiency should be taken into account when estimating this rate using isotope addition techniques. There is an urgent need to develop a method that can accurately measure the differences between ambient and newly spiked mercury

species in methylation and demethylation efficiency.

This work was partially supported by the U.S. Department of Energy (DE-FG01-05EW07033) and the National Natural Science Foundation of China (21120102040).

- 1 Mason R P, Fitzgerald W F, Morel F M M. The biogeochemical cycling of elemental mercury: Anthropogenic influences. *Geochimica Et Cosmochimica Acta*, 1994, 58: 3191–3198
- 2 Renner R. Mercury woes appear to grow. *Environ Sci Technol*, 2004, 38: 144A
- 3 Fitzgerald W F. *Atmospheric and Oceanic Cycling of Mercury*. San Diego: Academic Press, 1989
- 4 Stein E D, Cohen Y, Winer A M. Environmental distribution and transformation of mercury compounds. *Crit Rev Environ Sci Technol*, 1996, 26: 1–43
- 5 Pacyna E G, Pacyna J M, Fudala J, et al. Mercury emissions to the atmosphere from anthropogenic sources in Europe in 2000 and their scenarios until 2020. *Sci Total Environ*, 2006, 370: 147–156
- 6 Mergler D, Anderson H A, Chan L H M, et al. Methylmercury Exposure and health effects in humans: A worldwide concern. *AMBIO*, 2007, 36: 3–11
- 7 Jensen S, Jernelov A. Biological methylation of mercury in aquatic organisms. *Nature*, 1969, 223: 753–754
- 8 Compeau G, Bartha R. Methylation and demethylation of mercury under controlled redox, Ph, and salinity conditions. *Appl Environ Microbiol*, 1984, 48: 1203–1207
- 9 Berman M, Bartha R. Control of the methylation process in a mercury-polluted aquatic sediment. *Environ Pollut B*, 1986, 11: 41–53
- 10 Steffan R J, Korhals E T, Winfrey M R. Effects of acidification on mercury methylation, demethylation, and volatilization in sediments from an acid-susceptible lake. *Appl Environ Microbiol*, 1988, 54: 2003–2009
- 11 Gilmour C C, Henry E A, Mitchell R. Sulfate stimulation of mercury methylation in fresh-water sediments. *Environ Sci Technol*, 1992, 26: 2281–2287
- 12 Gilmour C C, Riedel G S, Ederington M C, et al. Methylmercury concentrations and production rates across a trophic gradient in the northern Everglades. *Biogeochemistry*, 1998, 40: 327–345
- 13 Marvin-DiPasquale M C, Agee J L, Bouse R M, et al. Microbial cycling of mercury in contaminated pelagic and wetland sediments of San Pablo Bay, California. *Environ Geol (Berl)*, 2003, 43: 260–267
- 14 Gray J E, Hines M E, Higuera P L, et al. Mercury speciation and microbial transformations in mine wastes, stream sediments, and surface waters at the Almaden Mining District, Spain. *Environ Sci Technol*, 2004, 38: 4285–4292
- 15 Heyes A, Mason R P, Kim E H, et al. Mercury methylation in estuaries: Insights from using measuring rates using stable mercury isotopes. *Mar Chem*, 2006, 102: 134–147
- 16 Drott A, Lambertsson L, Bjorn E, et al. Do potential methylation rates reflect accumulated methyl mercury in contaminated sediments? *Environ Sci Technol*, 2008, 42: 153–158
- 17 Furutani A, Rudd J W M. Measurement of mercury methylation in lake water and sediment samples. *Appl Environ Microbiol*, 1980, 40: 770–776
- 18 Matilainen T, Verta M. Mercury methylation and demethylation in aerobic surface waters. *Can J Fish Aquat Sci*, 1995, 52: 1597–1608
- 19 Siciliano S D, O'Driscoll N J, Tordon R, et al. Abiotic production of methylmercury by solar radiation. *Environ Sci Technol*, 2005, 39: 1071–1077
- 20 Monperrus M, Tessier E, Amouroux D, et al. Mercury methylation, demethylation and reduction rates in coastal and marine surface waters of the Mediterranean Sea. *Mar Chem*, 2007, 107: 49–63
- 21 Lehnher I, St Louis V L, Hintelmann H, et al. Methylation of inorganic mercury in polar marine waters. *Nat Geosci*, 2011, 4: 298–302
- 22 Cleckner L B, Gilmour C C, Hurley J P, et al. Mercury methylation in periphyton of the Florida Everglades. *Limnol Oceanogr*, 1999, 44: 1815–1825
- 23 Mauro J B N, Guimaraes J R D, Hintelmann H, et al. Mercury methylation in macrophytes, periphyton, and water-comparative studies with stable and radio-mercury additions. *Anal Bioanal Chem*, 2002, 374: 983–989
- 24 Desrosiers M, Planas D, Mucci A. Mercury methylation in the epilithon of boreal shield aquatic ecosystems. *Environ Sci Technol*, 2006, 40: 1540–1546
- 25 Yu R-Q, Adatto I, Montesdeoca M R, et al. Mercury methylation in Sphagnum moss mats and its association with sulfate-reducing bacteria in an acidic Adirondack forest lake wetland. *FEMS Microbiol Ecol*, 2010, 74: 655–668
- 26 Acha D, Hintelmann H, Yee J. Importance of sulfate reducing bacteria in mercury methylation and demethylation in periphyton from Bolivian Amazon region. *Chemosphere*, 2011, 82: 911–916
- 27 Hamelin S, Amyot M, Barkay T, et al. Methanogens: Principal methylators of mercury in Lake Periphyton. *Environ Sci Technol*, 2011, 45: 7693–7700
- 28 Correia R R S, Miranda M R, Guimaraes J R D. Mercury methylation and the microbial consortium in periphyton of tropical macrophytes: Effect of different inhibitors. *Environ Res*, 2012, 112: 86–91
- 29 Guimaraes J R D, Roulet M, Lucotte M, et al. Mercury methylation along a lake-forest transect in the Tapajos river floodplain, Brazilian Amazon: Seasonal and vertical variations. *Sci Total Environ*, 2000, 261: 91–98
- 30 Sellers P, Kelly C A, Rudd J W M, et al. Photodegradation of methylmercury in lakes. *Nature*, 1996, 380: 694–697
- 31 Hammerschmidt C R, Fitzgerald W F. Photodecomposition of methylmercury in an arctic Alaskan lake. *Environ Sci Technol*, 2006, 40: 1212–1216
- 32 Li Y B, Mao Y X, Liu G L, et al. Degradation of methylmercury and its effects on mercury distribution and cycling in the Florida Everglades. *Environ Sci Technol*, 2010, 44: 6661–6666
- 33 Bergquist B A, Blum J D. Mass-dependent and -independent fractionation of Hg isotopes by photoreduction in aquatic systems. *Science*, 2007, 318: 417–420
- 34 Naftz D L, Cederberg J R, Krabbenhoft D P, et al. Diurnal trends in methylmercury concentration in a wetland adjacent to Great Salt Lake, Utah, USA. *Chem Geol*, 2011, 283: 78–86
- 35 Reisinger K, Stoeppeler M, Nurnberg H W. Biological methylation of inorganic mercury by *Saccharomyces-Cerevisiae*—A possible environmental process. *Fresenius Zeitschrift Fur Analytische Chemie*, 1983, 316: 612–615
- 36 Trevors J T. Mercury methylation by bacteria. *J Basic Microbiol*, 1986, 26: 499–504
- 37 Choi S C, Bartha R. Cobalamin-mediated mercury methylation by *Desulfovibrio-Desulfuricans* Ls. *Appl Environ Microbiol*, 1993, 59: 290–295
- 38 Choi S C, Chase T, Bartha R. Metabolic pathways leading to mercury methylation in *Desulfovibrio-Desulfuricans* Ls. *Appl Environ Microbiol*, 1994, 60: 4072–4077
- 39 Choi S C, Chase T, Bartha R. Enzymatic catalysis of mercury methylation by *Desulfovibrio-Desulfuricans* Ls. *Appl Environ Microbiol*, 1994, 60: 1342–1346
- 40 Chen Y, Bonzongo J C J, Lyons W B, et al. Inhibition of mercury methylation in anoxic freshwater sediment by group VI anions. *Environ Toxicol Chem*, 1997, 16: 1568–1574
- 41 Pak K R, Bartha R. Mercury methylation by interspecies hydrogen and acetate transfer between sulfidogens and methanogens. *Appl Environ Microbiol*, 1998, 64: 1987–1990
- 42 Pak K R, Bartha R. Mercury methylation and demethylation in anoxic lake sediments and by strictly anaerobic bacteria. *Appl Environ Microbiol*, 1998, 64: 1013–1017
- 43 King J K, Saunders F M, Lee R F, et al. Coupling mercury methylation rates to sulfate reduction rates in marine sediments. *Environ Toxicol Chem*, 1999, 18: 1362–1369
- 44 King J K, Kostka J E, Frischer M E, et al. Sulfate-reducing bacteria methylate mercury at variable rates in pure culture and in marine sediments. *Appl Environ Microbiol*, 2000, 66: 2430–2437
- 45 Macalady J L, Mack E E, Nelson D C, et al. Sediment microbial community structure and mercury methylation in mercury-polluted Clear Lake, California. *Appl Environ Microbiol*, 2000, 66: 1479–1488

- 46 Siciliano S D, Lean D R S. Methyltransferase: An enzyme assay for microbial methylmercury formation in acidic soils and sediments. *Environ Toxicol Chem*, 2002, 21: 1184–1190
- 47 Ekstrom E B, Morel F M M, Benoit J M. Mercury methylation independent of the acetyl-coenzyme a pathway in sulfate-reducing bacteria. *Appl Environ Microbiol*, 2003, 69: 5414–5422
- 48 Marvin-DiPasquale M, Agee J L. Microbial mercury cycling in sediments of the San Francisco Bay-Delta. *Estuaries*, 2003, 26: 1517–1528
- 49 Ranchou-Peyruse M, Monperrus M, Bridou R, et al. Overview of mercury methylation capacities among anaerobic bacteria including representatives of the sulfate-reducers: Implications for environmental studies. *Geomicrobiol J*, 2009, 26: 1–8
- 50 Avramescu M L, Yumvihoze E, Hintelmann H, et al. Biogeochemical factors influencing net mercury methylation in contaminated freshwater sediments from the St. Lawrence River in Cornwall, Ontario, Canada. *Sci Total Environ*, 2011, 409: 968–978
- 51 Larock R C, Hershberger S S. Mercury in organic-chemistry. 19. Rhodium promoted methylation of organomercurials. *Tetrahedron Lett*, 1981, 22: 2443–2446
- 52 Larock R C, Hershberger S S. Mercury in organic-chemistry 21. Methylation of organomercurials via organorhodium species. *J Organomet Chem*, 1982, 225: 31–41
- 53 Nagase H, Ose Y, Sato T, et al. Methylation of mercury by humic substances in an aquatic environment. *Sci Total Environ*, 1982, 25: 133–142
- 54 Nagase H, Ose Y, Sato T, et al. Production of methylmercury by abiological methylation of inorganic mercury in the environment. *Jpn J Toxicol Environ Health*, 1983, 29: 55
- 55 Nagase H, Ose Y, Sato T, et al. Mercury methylation by compounds in humic material. *Sci Total Environ*, 1984, 32: 147–156
- 56 Woggon H, Klein S, Jehle D, et al. Transformation reactions of special metals in organisms and in the environment. 2. Abiological methylation reactions of mercury, especially by methyltin compounds, and humic and fulvic-acids. *Nahrung-Food*, 1984, 28: 851–862
- 57 Lee Y H, Hultberg H, Andersson I. Catalytic effect of various metalions on the methylation of mercury in the presence of humic substances. *Water Air Soil Pollut*, 1985, 25: 391–400
- 58 Weber J H, Reisinger K, Stoeppler M. Methylation of mercury(II) by fulvic-acid. *Environ Technol Lett*, 1985, 6: 203–208
- 59 Bellama J M, Jewett K L, Nies J D. Methylation of mercury(II) species in water by organosilicon compounds. *Abstr Pap Am Chem Soc*, 1986, 192: 127–GEOC
- 60 Nagase H, Ose Y, Sato T, et al. Mercury methylation by ash from refuse incineration. *Sci Total Environ*, 1986, 53: 133–138
- 61 Watanabe N, Nagase H, Nakamura T, et al. Chemical methylation of mercury(II) salts by polydimethylsiloxanes in aqueous-solution. *Ecotoxicol Environ Saf*, 1986, 11: 174–178
- 62 Bellama J M, Jewett K L, Manders W F, et al. A comparison of the rates of methylation of mercury(II) species in aquatic media by various organotin and organo-silicon moieties. *Sci Total Environ*, 1988, 73: 39–51
- 63 Cerrati G, Bernhard M, Weber J H. Model reactions for abiotic mercury(II) methylation-kinetics of methylation of mercury(II) by mono-methyltin, di-methyltin, and tri-methyltin in seawater. *Appl Organomet Chem*, 1992, 6: 587–595
- 64 Weber J H. Review of possible paths for abiotic methylation of mercury(II) in the aquatic environment. *Chemosphere*, 1993, 26: 2063–2077
- 65 Gardfeldt K, Munthe J, Stromberg D, et al. A kinetic study on the abiotic methylation of divalent mercury in the aqueous phase. *Sci Total Environ*, 2003, 304: 127–136
- 66 Yin Y, Chen B, Mao Y, et al. Possible alkylation of inorganic Hg(II) by photochemical processes in the environment. *Chemosphere*, 2012 (in press)
- 67 Berman M, Bartha R. Levels of chemical versus biological methylation of mercury in sediments. *Bull Environ Contam Toxicol*, 1986, 36: 401–404
- 68 Mauro J B N, Guimaraes J R D, Melamed R. Mercury methylation in macrophyte roots of a tropical lake. *Water Air Soil Pollut*, 2001, 127: 271–280
- 69 Kerry A, Welbourn P M, Prucha B, et al. Mercury methylation by sulfate-reducing bacteria from sediments of an acid stressed lake. *Water Air Soil Pollut*, 1991, 56: 565–575
- 70 King J K, Kostka J E, Frischer M E, et al. A quantitative relationship that remonstrates mercury methylation rates in marine sediments are based on the community composition and activity of sulfate-reducing bacteria. *Environ Sci Technol*, 2001, 35: 2491–2496
- 71 Dias M, Salvado J C, Monperrus M, et al. Characterization of *Desulfomicrobium salsuginis* sp. nov. and *Desulfomicrobium aestuarii* sp. nov., two new sulfate-reducing bacteria isolated from the Adour estuary (French Atlantic coast) with specific mercury methylation potentials. *Syst Appl Microbiol*, 2008, 31: 30–37
- 72 Ekstrom E B, Morel F M M. Cobalt limitation of growth and mercury methylation in sulfate-reducing bacteria. *Environ Sci Technol*, 2008, 42: 93–99
- 73 Jeremiason J D, Engstrom D R, Swain E B, et al. Sulfate addition increases methylmercury production in an experimental wetland. *Environ Sci Technol*, 2006, 40: 3800–3806
- 74 Compeau G C, Bartha R. Sulfate-reducing bacteria: Principal methylators of mercury in anoxic estuarine sediment. *Appl Environ Microbiol*, 1985, 50: 498–502
- 75 Fleming E J, Mack E E, Green P G, et al. Mercury methylation from unexpected sources: Molybdate-inhibited freshwater sediments and an iron-reducing bacterium. *Appl Environ Microbiol*, 2006, 72: 457–464
- 76 Kerin E J, Gilmour C C, Roden E, et al. Mercury methylation by dissimilatory iron-reducing bacteria. *Appl Environ Microbiol*, 2006, 72: 7919–7921
- 77 Hayashi K, Kawai S, Ohno T, et al. Photomethylation of inorganic mercury by aliphatic alpha-amino acids. *J Chem Soc Chem Commun*, 1977: 158–159
- 78 Nagase H, Ose Y, Sato T. Possible methylation of inorganic mercury by silicones in the environment. *Sci Total Environ*, 1988, 73: 29–38
- 79 Li Y, Yin Y, Liu G, et al. Estimation of the major source and sink of methylmercury in the Florida Everglades. *Environ Sci Technol*, 2012, 46: 5885–5893
- 80 Oremland R S, Culbertson C W, Winfrey M R. Methylmercury decomposition in sediments and bacterial cultures: Involvement of methanogens and sulfate reducers in oxidative demethylation. *Appl Environ Microbiol*, 1991, 57: 130–137
- 81 Baldi F, Pepi M, Filippelli M. Methylmercury resistance in *Desulfovibrio desulfuricans* strains in relation to methylmercury degradation. *Appl Environ Microbiol*, 1993, 59: 2479–2485
- 82 Oremland R S, Miller L G, Dowdle P, et al. Methylmercury oxidative degradation potentials in contaminated and pristine sediments of the carson river, nevada. *Appl Environ Microbiol*, 1995, 61: 2745–2753
- 83 Inoko M. Studies on the photochemical decomposition of organomercurials-methylmercury(II) chloride. *Environ Pollut B*, 1981, 2: 3–10
- 84 Suda I, Takahashi H. Degradation of methyl and ethyl mercury into inorganic mercury by other reactive oxygen species besides hydroxyl radical. *Arch Toxicol*, 1992, 66: 34–39
- 85 Suda I, Suda M, Hirayama K. Degradation of methyl and ethyl mercury by singlet oxygen generated from sea-water exposed to sunlight or ultraviolet-light. *Arch Toxicol*, 1993, 67: 365–368
- 86 Marvin-DiPasquale M, Agee J, McGowan C, et al. Methyl-mercury degradation pathways: A comparison among three mercury-impacted ecosystems. *Environ Sci Technol*, 2000, 34: 4908–4916
- 87 Gardfeldt K, Sommar J, Stromberg D, et al. Oxidation of atomic mercury by hydroxyl radicals and photoinduced decomposition of methylmercury in the aqueous phase. *Atmos Environ*, 2001, 35: 3039–3047
- 88 Chen J, Pehkonen S O, Lin C J. Degradation of monomethylmercury chloride by hydroxyl radicals in simulated natural waters. *Water Res*, 2003, 37: 2496–2504
- 89 Lehnher I, St Louis V L. Importance of ultraviolet radiation in the photodemethylation of methylmercury in freshwater ecosystems. *Environ Sci Technol*, 2009, 43: 5692–5698
- 90 Hammerschmidt C R, Fitzgerald W F. Iron-mediated photochemical decomposition of methylmercury in an Arctic Alaskan lake. *Environ Sci Technol*, 2010, 44: 6138–6143
- 91 Zhang T, Hsu-Kim H. Photolytic degradation of methylmercury

- enhanced by binding to natural organic ligands. *Nat Geosci*, 2010, 3: 473–476
- 92 Khan M A K, Wang F Y. Chemical demethylation of methylmercury by selenoamino acids. *Chem Res Toxicol*, 2010, 23: 1202–1206
- 93 Asaduzzaman A, Schreckenbach G. Degradation mechanism of methyl mercury selenoamino acid complexes: A computational study. *Inorg Chem*, 2011, 50: 2366–2372
- 94 Marvin-DiPasquale M C, Oremland R S. Bacterial methylmercury degradation in Florida Everglades peat sediment. *Environ Sci Technol*, 1998, 32: 2556–2563
- 95 Schaefer J K, Yagi J, Reinfelder J R, et al. Role of the bacterial organomercury lyase (MerB) in controlling methylmercury accumulation in mercury-contaminated natural waters. *Environ Sci Technol*, 2004, 38: 4304–4311
- 96 Allard B, Arsenie I. Abiotic reduction of mercury by humic substances in aquatic system—an important process for the mercury cycle. *Water Air Soil Pollut*, 1991, 56: 457–464
- 97 Black F J, Poulin B A, Flegal A R. Factors controlling the abiotic photo-degradation of monomethylmercury in surface waters. *Geochimica Et Cosmochimica Acta*, 2012, 84: 492–507
- 98 Chandan P. Mercury Isotope Fractionation During Aqueous Photo-reduction of Methylmercury in Presence of Different Types and Amounts of Dissolved Organic Matter. Toronto: University of Toronto, 2011
- 99 Han S, Obratsova A, Pretto P, et al. Biogeochemical factors affecting mercury methylation in sediments of the Venice Lagoon, Italy. *Environ Toxicol Chem*, 2007, 26: 655–663
- 100 Hammerschmidt C R, Fitzgerald W F. Geochemical controls on the production and distribution of methylmercury in near-shore marine sediments. *Environ Sci Technol*, 2004, 38: 1487–1495
- 101 Benoit J M, Gilmour C C, Mason R P. The influence of sulfide on solid phase mercury bioavailability for methylation by pure cultures of *Desulfobulbus propionicus* (1pr3). *Environ Sci Technol*, 2001, 35: 127–132
- 102 Hintelmann H, Keppel-Jones K, Evans R D. Constants of mercury methylation and demethylation rates in sediments and comparison of tracer and ambient mercury availability. *Environ Toxicol Chem*, 2000, 19: 2204–2211
- 103 Farrell R E, Huang P M, Germida J J. Biomethylation of mercury(II) adsorbed on mineral colloids common in freshwater sediments. *Appl Organomet Chem*, 1998, 12: 613–620
- 104 Morel F M M, Kraepiel A M L, Amyot M. The chemical cycle and bioaccumulation of mercury. *Annu Rev Ecol Syst*, 1998, 29: 543–566
- 105 Drott A, Lambertsson L, Björn E, et al. Importance of dissolved neutral mercury sulfides for methyl mercury production in contaminated sediments. *Environ Sci Technol*, 2007, 41: 2270–2276
- 106 Wolfenden S, Charnock J M, Hilton J, et al. Sulfide species as a sink for mercury in lake sediments. *Environ Sci Technol*, 2005, 39: 6644–6648
- 107 Ullrich S M, Tanton T W, Abdrashitova S A. Mercury in the aquatic environment: A review of factors affecting methylation. *Crit Rev Environ Sci Technol*, 2001, 31: 241–293
- 108 Fitzgerald W F, Lamborg C H, Hammerschmidt C R. Marine biogeochemical cycling of mercury. *Chem Rev*, 2007, 107: 641–662
- 109 Benoit J M, Gilmour C C, Mason R P, et al. Sulfide controls on mercury speciation and bioavailability to methylating bacteria in sediment pore waters. *Environ Sci Technol*, 1999, 33: 951–957
- 110 Benoit J M, Mason R P, Gilmour C C. Estimation of mercury-sulfide speciation in sediment pore waters using octanol-water partitioning and implications for availability to methylating bacteria. *Environ Toxicol Chem*, 1999, 18: 2138–2141
- 111 Benoit J M, Gilmour C C, Mason R P. Aspects of bioavailability of mercury for methylation in pure cultures of *Desulfobulbus propionicus* (1pr3). *Appl Environ Microbiol*, 2001, 67: 51–58
- 112 Lin C C, Jay J A. Mercury methylation by planktonic and biofilm cultures of *Desulfobulbus desulfuricans*. *Environ Sci Technol*, 2007, 41: 6691–6697
- 113 Hammerschmidt C R, Fitzgerald W F, Balcom P H, et al. Organic matter and sulfide inhibit methylmercury production in sediments of New York/New Jersey Harbor. *Mar Chem*, 2008, 109: 165–182
- 114 Schaefer J K, Morel F M M. High methylation rates of mercury bound to cysteine by *Geobacter sulfurreducens*. *Nat Geosci*, 2009, 2: 123–126
- 115 Schaefer J K, Rocks S S, Zheng W, et al. Active transport, substrate specificity, and methylation of Hg(II) in anaerobic bacteria. *Proc Natl Acad Sci USA*, 2011, 108: 8714–8719
- 116 Ravichandran M. Interactions between mercury and dissolved organic matter—A review. *Chemosphere*, 2004, 55: 319–331
- 117 Baughman G L, Gordon J A, Wolfe N L, et al. Chemistry of Organomercurials in Aquatic Systems. Washington DC: US Environmental Protection Agency, 1973
- 118 Akagi H, Mortimer D C, Miller D R. Mercury methylation and partition in aquatic systems. *Bull Environ Contam Toxicol*, 1979, 23: 372–376
- 119 Gilmour C C, Riedel G S. Measurement of Hg methylation in sediments using high specific-activity Hg-203 and ambient incubation. *Water Air Soil Pollut*, 1995, 80: 747–756
- 120 Koron N, Bratkic A, Guevara S R, et al. Mercury methylation and reduction potentials in marine water: An improved methodology using (197)Hg radiotracer. *Appl Radiat Isotop*, 2012, 70: 46–50
- 121 Monperrus M, Tessier E, Point D, et al. The biogeochemistry of mercury at the sediment-water interface in the Thau Lagoon. 2. Evaluation of mercury methylation potential in both surface sediment and the water column. *Estuar Coast Shelf Sci*, 2007, 72: 485–496
- 122 Hintelmann H, Evans R D, Villeneuve J Y. Measurement of mercury methylation in sediments by using enriched stable mercury isotopes combined with methylmercury determination by gas-chromatography inductively-coupled plasma-mass spectrometry. *J Anal At Spectrom*, 1995, 10: 619–624
- 123 Hintelmann H, Harris R, Heyes A, et al. Reactivity and mobility of new and old mercury deposition in a Boreal forest ecosystem during the first year of the METAALICUS study. *Environ Sci Technol*, 2002, 36: 5034–5040
- 124 Lambertsson L, Lundberg E, Nilsson M, et al. Applications of enriched stable isotope tracers in combination with isotope dilution GC-ICP-MS to study mercury species transformation in sea sediments during *in situ* ethylation and determination. *J Anal At Spectrom*, 2001, 16: 1296–1301
- 125 Martin-Doimeadios R C, Tessier E, Amouroux D, et al. Mercury methylation/demethylation and volatilization pathways in estuarine sediment slurries using species-specific enriched stable isotopes. *Mar Chem*, 2004, 90: 107–123
- 126 Whalin L M, Mason R P. A new method for the investigation of mercury redox chemistry in natural waters utilizing deflatable Teflon (R) bags and additions of isotopically labeled mercury. *Anal Chim Acta*, 2006, 558: 211–221
- 127 Monperrus M, Gonzalez P R, Amouroux D, et al. Evaluating the potential and limitations of double-spiking species-specific isotope dilution analysis for the accurate quantification of mercury species in different environmental matrices. *Anal Bioanal Chem*, 2008, 390: 655–666
- 128 Bridou R, Monperrus M, Gonzalez P R, et al. Simultaneous determination of mercury methylation and demethylation capacities of various sulfate-reducing bacteria using species-specific isotopic tracers. *Environ Toxicol Chem*, 2011, 30: 337–344
- 129 Hintelmann H, Evans R D. Application of stable isotopes in environmental tracer studies—Measurement of monomethylmercury (CH₃Hg⁺) by isotope dilution ICP-MS and detection of species transformation. *Fresen J Anal Chem*, 1997, 358: 378–385
- 130 Gray J E, Hines M E, Biester H. Mercury methylation influenced by areas of past mercury mining in the Terlingua district, Southwest Texas, USA. *Appl Geochem*, 2006, 21: 1940–1954