

ERRATUM

Erratum to: Machine learning identification of symmetrized base states of Rydberg atoms

Daryl Ryan Chong¹, Minhyuk Kim², Jaewook Ahn², Heejeong Jeong^{1,†}

¹Department of Physics, Faculty of Science, University of Malaya, Kuala Lumpur 50603, Malaysia

²Department of Physics, KAIST, Daejeon 34141, Korea

Corresponding author. E-mail: †jhj413@gmail.com

ERRATUM TO: *Front. Phys.* 17(1), 12504 (2022),
<https://doi.org/10.1007/s11467-021-1099-0>.

at <https://doi.org/10.1007/s11467-021-1099-0> and <http://journal.hep.com.cn/fop/EN/10.1007/s11467-021-1099-0>.

The x -axis of Fig. 3(b) should be labelled “Observation duration” instead of “Total evolution time” as shown below.

The online version of the original article can be found




Fig. 3 Accuracy performance depending on (a) total evolution time, and (b) observation duration between data points for the case of identification of Na in Figs. 2(a, c, e). The accuracy values are obtained by utilizing both SVM and RFC. The default total evolution time is 1 μ s, and the default duration is 50 ns for all our ML sample data in this paper. The results show our default experimental parameters provides above 90% accuracy as in Figs. 2(a, c, e).

*Special Topic: Trapped Atoms and Ions for Quantum Science (Eds. Le Luo, Kenji Toyoda, Kihwan Kim, Jaewook Ahn & Dzmitry Matsukevich).