




# Identification and Pathogenicity of Species Isolated from Stored Potato Tubers Showing Symptoms of Dry Rot Disease

Nam Sook Kim<sup>1</sup> · Sae Jin Hong<sup>1</sup> · Heon Seop Won<sup>2</sup> · Byung Sup Kim<sup>1</sup> · Se Hwi Gwon<sup>1</sup> 



Received: 23 July 2023 / Accepted: 19 February 2024  
© The Author(s) 2024

## Abstract

Disease arises during the storage of potatoes as a result of pathogens introduced during growing and harvest of the tubers. In this study, the causative fungi of domestic potato dry rot disease were identified, and their pathogenicity was confirmed. A total of 76 species were isolated from 93 potato necropods collected from samples inoculated with dry rot disease. These 76 isolates were identified as *Fusarium boothii*, *F. circinatum*, *F. citricola*, *F. foetens*, *F. iranicum*, *F. longifundum*, *F. oxysporum*, *F. pseudoanthophilum*, *F. solani*, *Botryotinia ranunculi*, *Clonostachys rosea*, and *Humicola nigrescens*. The average size of the inoculation site was  $\geq 4.6$  mm in *F. oxysporum* and *F. solani*, which were pathogenic to dry rot in potatoes but were not pathogenic compared to other strains up to 5 weeks in a 15 to 20 °C and 99% RH environment. The pathogenicity of *F. foetens* and *F. pseudoanthophilum* was related to a strong relationship by forming a single system with *F. oxysporum*. However, except for *F. oxysporum* and *F. solani*, these strains have not yet been reported to be associated with dry rot disease. Additionally, the length of the cross-section and longitudinal section of the potato damage symptom inoculated with *C. rosea* increased the most among all strains. This suggests that *C. rosea* is the dominant species involved in domestic potato dry rot disease. By contrast, there are no reports of the involvement of *B. ranunculi* and *H. nigrescens* in dry rot disease. Therefore, these strains can be seen as parasitic using potato sclerosis as nutrients in in vivo experiments through wounds and are not directly related to dry rot disease.

**Keywords** *Clonostachys rosea* · *Fusarium* spp. · Postharvest disease · *Solanum tuberosum*

---

Communicated by Se Hwi Gwon

---

Nam Sook Kim and Sae Jin Hong contributed equally to this work.

---

Extended author information available on the last page of the article

## Introduction

Potatoes belong to the Solanaceae family, with approximately 230 species found in natural ecosystems, among which approximately 160 species form tubers (Kwon et al. 2005). Today, *Solanum tuberosum* is grown worldwide, with approximately 388 million tons of tubers produced annually across 130 countries (Kim 2020), 690,419 MT (metric ton) of which were produced in Korea in 2019 (KOSIS 2021). As the highest source of energy and of production per unit area among the world's major cultivated crops, potato production is expected to continue to increase (RDA 2020). However, potatoes are vulnerable to a variety of diseases when stored for long periods of time. For example, losses caused by dry rot disease are estimated to range from 6 to 25% worldwide, with losses of up to 60% during long-term storage periods (Desjardins 2006; Secor and Salas 2001). The estimated economic damage of crops lost in the United States, the largest potato producer, ranges from \$100 million to \$250 million per year (Slinger and Schisler 2002), and accounts for up to 88% of the total post-harvest losses of potatoes in Gansu, China (He et al. 2004). Despite a lack of accurate statistics or reports, most potatoes with pathological disorders during potato storage in Korea have been identified as having dry rot disease (data not shown).

Dry rot disease is caused by the intrusion of various *Fusarium* species through wounds on the surface of potatoes during their harvest (Nelson et al. 1981). This pathogen invades potatoes lodging in the soil and seeds and causes crop loss during storage (Tiwari et al. 2020). Starting as a small brown lesion, potato dry rot disease gradually expands, developing into a concentric dry region around dead tissue (RDA 2020). Subsequently, the wrinkled tissue changes from brown to black and can be distinguished with the naked eye. The lesion of the wrinkled tissue produces creamy white, pink, or orange spores and mycelium during long-term storage at temperatures ranging from 5 to 30 °C (Bojanowski et al. 2013; Elsherbiny et al. 2016).

Dry rot caused by *Fusarium* occurs in all potato cultivation areas, and the distribution of *Fusarium* species varies depending on the cultivation time and geographical location (Tiwari et al. 2020). More than 13 species, including *F. acuminatum*, *F. avenaceum*, *F. crookwellense*, *F. culmorum*, *F. equiseti*, *F. graminearum*, *F. oxysporum*, *F. sambucinum*, *F. scirpi*, *F. semitectum*, *F. solani*, *F. sporotrichioides*, and *F. tricinctum*, are involved in the dry rot disease of potatoes worldwide (Cullen et al. 2005). The dominant species that cause dry rot in potatoes are *F. sambucinum* and *F. graminearum* in North America, *F. sambucinum* in Europe, *F. solani* in the United Kingdom, *F. solani* and *F. oxysporum* in South Africa, and *F. sambucinum* in China (Tiwari et al. 2020). Until 2000, *F. solani* was suggested as a fungus that caused potato dry rot disease in Korea, and *F. oxysporum* was reported as a pathogen that caused withering diseases in various crops; however, this was not explained in connection with potato dry rot disease (Kim 2000; Park et al. 1988). Currently, dry rot disease pathogens in potatoes are registered as *Fusarium* spp. in the National Agricultural Products Disease and Pest Management System, and various *Fusarium* spp. have been

reported to cause the disease (NCPMS 2023). In addition, one study showed that *Clonostachys rosea*, also known as *Gliocladium roseum*, is involved in dry rot disease and simultaneously acts as an antagonist to *F. solani* and *F. oxysporum* (Theron and Holz 1991).

Research has been conducted in North America, Europe, the UK, South Africa, and China to identify the dominant species of pathogens causing potato dry rot. However, few studies have identified the causative fungi of dry rot disease during storage or reported the dominant species in Korea. Therefore, this study was conducted to identify and confirm the pathogenicity of the causative pathogens of domestic potato dry rot disease in Korea.

## Materials and Methods

### Collection of Pathogens for Dry Rot of Potatoes


Tubers were collected from potatoes stored in traditional potato warehouses located on highland areas, including Wangsan-myeon and Daegwallyeong-myeon, and located on plain area, including Gangneung Wonju University's farm and Sacheon-myeon in Gangwon Province. The potato varieties included Atlantic, Chubaek, Doobak, and Superior. The collected pathogens of potato were stored in a 20 °C incubator at Gangneung Wonju University's Plant Pathology Laboratory for 3 days to induce growth.

### Separation and Identification of Pathogens

Dry rot pathogens were isolated from the plant pathology laboratory of Gangneung Wonju University using *Solanum tuberosum* damage symptoms with signs of disease. To separate pathogens, fragments, including the surface of the potato damage symptom containing the boundary of the disease, were prepared by cutting into 1-cm diameter slices. The potato slices were sterilized in 70% ethanol for 30 s and 1% sodium hypochlorite (NaOCl) for 2 min on a clean bench, washed once with sterilized water, and dried on a sterilized filter paper for 30 min. One dried fragment from the infected damage symptom was placed in 10 WA (water agar medium) and cultured for 3 days in a 25 °C incubator (Fig. 1). Mycelia of pathogens grown in potato fragments were cultured in a 20 °C incubator for 7 days by cutting the tip of the pathogen for pure separation and putting on potato dextrose agar (PDA) medium. Each fungus was separated from the potato agar medium after confirming the shape of mycelial growth, and the separated samples were analyzed using the ITS1/ITS4 primer pair. The phylogenetic tree was constructed using the Neighbor-joining method with MEGA7.

### Pathogenicity Test of Dried Potato Rot Strain

The potatoes used in the pathogenicity test were superior varieties grown in Wangsan-myeon, Gangneung-si, and Gangwon-do. The 12 identified strains were used as inoculators. The potato samples were immersed in 1% NaOCl for 30 min, washed



**Fig. 1** Colony morphologies of the isolated fungal species on water agar medium incubated at 25 °C for 3 days from potatoes showing symptoms of dry rot disease on tubers stored in cold storages. **A** Beckjak; **B** Dano; **C** Doobak; **D** Jayoung; **E** Superior; **F** Wangsan

three times with sterile water after sterilization, and dried for 30 min. The inoculator prepared 12 identified pathogens by implanting them in PDA and culturing them in an incubator at 20 °C for 7 days. For the treatment of the inoculum, a 2-mm-deep hole was made in the base of the potato necrosis using a  $\phi$ 8mm cork borer, and the inoculum was cut and inserted with a  $\phi$ 10mm cork borer. Potatoes inoculated with the identified fungi were separated and cultured in a sealed plastic container to prevent air contamination by pathogens during culturing. The invasion and pathogenicity of pathogens through wounds were investigated by placing the fungally inoculated sclera in a humidified translucent plastic box and maintaining a relative humidity of 99% at room temperature of 15–20 °C (Table 1).

The inoculation site and pathogenicity were visually checked at intervals of 7 days after inoculation, and the length of disease spread was measured in the cross-section (Fig. 2) and longitudinal section (Fig. 3) of the potato damage symptom at 5 weeks after inoculation (Table 2). Each treatment port, including the control port, was performed five times based on 13 unusual diameters. Statistical analysis was performed using SPSS (IBM SPSS Statistics for Windows version 26.0 (IBM Corp., Armonk, NY, USA) to analyze ANOVA, and the significance of each treatment was verified at  $P < 0.05$  using Duncan's multiple range test (DMRT).


## Results and Discussion

### Collection of Pathogens for Dry Rot in Potatoes

The total number of potatoes with dry rot disease after storage was 93, including 76 in Gangneung plain, 3 in Pyeongchang, 11 in Gangneung highland. To collect individuals with dry rot disease from potatoes, potato-only storage was


**Table 1** Type and number of fungi isolated from 93 tubers showing symptoms of dry rot disease (2021–2022)

Fungal species	No. of isolates
<i>Fusarium boothii</i>	8
<i>Fusarium circinatum</i>	5
<i>Fusarium citricola</i>	14
<i>Fusarium foetens</i>	16
<i>Fusarium iranicum</i>	5
<i>Fusarium longifundum</i>	2
<i>Fusarium oxysporum</i>	1
<i>Fusarium pseudoanthophilum</i>	4
<i>Fusarium solani</i>	13
<i>Botryotinia ranunculi</i>	2
<i>Clonostachys rosea</i>	4
<i>Humicola nigrescens</i>	2
Total	76



**Fig. 2** Disease symptoms of infected areas on potato tubers 7 weeks after the artificial inoculation of 12 fungal species incubated at 15–20 °C and 99% RH. **A** Control; **B** *Fusarium boothii*; **C** *Fusarium circinatum*; **D** *Fusarium citricola*; **E** *Fusarium foetens*; **F** *Fusarium iranicum*; **G** *Fusarium longifundum*; **H** *Fusarium oxysporum*; **I** *Fusarium pseudoanthophilum*; **J** *Fusarium solani*; **K** *Botryotinia ranunculi*; **L** *Clonostachys rosea*; **M** *Humicola nigrescens*

investigated, and most potato objects with external diseases were infected with dry rot disease. There were a few cases of infection during storage, and it is believed that the spread of the disease was suppressed by screening and curing after harvest. Dry rot disease that occurred in certain varieties, such as Atlantic,



**Fig. 3** Appearance of the longitudinal sections of potato tubers 7 weeks after the artificial inoculation of 12 fungal species incubated at 15–20 °C and 99% RH. **A** Control; **B** *Fusarium boothii*; **C** *Fusarium circinatum*; **D** *Fusarium citricola*; **E** *Fusarium foetens*; **F** *Fusarium iranicum*; **G** *Fusarium longifundum*; **H** *Fusarium oxysporum*; **I** *Fusarium pseudoanthophilum*; **J** *Fusarium solani*; **K** *Botryotinia ranunculi*; **L** *Clonostachys rosea*; **M** *Humicola nigrescens*

**Table 2** Growth estimation based on the length of transverse and longitudinal sections to verify the pathogenicity of dry rot disease in potato tubers inoculated by 12 fungi collected in this study


Treatment (inoculated fungi)	Longitudinal section (mm)	Transverse section (mm)
Control	0.00 c	0.00 d <sup>z</sup>
<i>Fusarium boothii</i>	3.46 b	3.50 c
<i>Fusarium circinatum</i>	3.68 b	5.84 b
<i>Fusarium citricola</i>	4.84 b	5.44 bc
<i>Fusarium foetens</i>	3.91 b	4.56 bc
<i>Fusarium iranicum</i>	3.13 b	3.66 c
<i>Fusarium longifundum</i>	3.06 b	4.2 bc
<i>Fusarium oxysporum</i>	2.93 b	4.94 bc
<i>Fusarium pseudoanthophilum</i>	3.12 b	3.8 bc
<i>Fusarium solani</i>	3.34 b	4.62 bc
<i>Botryotinia ranunculi</i>	2.93 b	3.36 c
<i>Clonostachys rosea</i>	6.90 a	11.48 a
<i>Humicola nigrescens</i>	3.05 b	4.28 bc

<sup>z</sup>Mean separation within columns by Duncan’s multiple ranges test at  $P \leq 0.05$

Beckjak, Chubaek, Dano, Doobak, Jayoung, Superior, and Wangsan, showed typical symptoms of disease regardless of the variety (Fig. 4).

### Separation and Identification of Pathogens


Of the 93 potatoes harbouring dry rot disease, some had duplicate or separate fungal strains. As such, a total of 76 were ultimately isolated (Table 1). After analyzing the sequence of the ITS region using the ITS1/ITS4 primer pair in the 76 separate



**Fig. 4** Potato tubers with symptoms of dry rot disease under common storage conditions (2021–2022). **A** Atlantic, **B** Beckjak, **C** Chubaek, **D** Dano, **E** Doobak, **F** Jayoung, **G** Superior, and **H** Wangsan

fungal strains, a total of 12 strains were analyzed, including nine from *Fusarium*, one from *Botryotinia*, one from *Clonostachys*, and one from *Humicola*.

The nine strains of the *Fusarium* spp. were *F. boothii*, *F. circinatum*, *F. citricola*, *F. foetens*, *F. iranicum*, *F. longifundum*, *F. oxysporum*, *F. pseudoanthophilum*, and *F. solani*. Among the *Fusarium*, the most frequently isolated strains were *F. foetens* (16), *F. citricola* (14), and *F. solani*. Other *Fusarium* strains were separated in the following order: *F. boothii*, *F. circinatum*, *F. iranicum*, *F. pseudoanthophilum*, *F. longifundum*, and *F. oxysporum*. The *Fusarium* strains involved in dry rot disease, which are most commonly found in domestic potato stores, are believed to be *F. foetens*, *F. citricola*, and *F. solani*. The remaining strains were identified as *Botryotinia ranunculi*, *Clonostachys rosea*, and *Humicola nigrescens* (Fig. 5). Among these, *F. solani*, *F. oxysporum*, and *C. rosea* have been reported to cause dry rot disease (Bojanowski et al. 2013; Theron and Holz 1991). *C. rosea* is involved in dry rot disease and simultaneously acts as an antagonist of *F. solani* and *F. oxysporum* (Theron and Holz 1991), although there have been no reports of any other identified pathogens associated with dry rot disease. The following have been reported to be representative of diseases caused by each pathogen: *F. boothii* has been reported to be associated with corn ear rot, pecan malady (Gryzenhout et al. 2016) and wheat red mold disease (Wegulo et al. 2018); *F. circinatum* causes pine branch disease (Woo et al. 2011) and pine pitch canker (Maphosa et al. 2016); *F. citricola* causes tangerine branch ulcer disease (Sandoval-Denis et al. 2018); *F. foetens* is pathogenic to begonia (Schroers et al. 2004) and has been associated with the occurrence of damping-off of rooibos seedlings (Lamprecht and Tewoldemedhin 2017); *F. iranicum* has been identified in mushroom (Torbati et al. 2019) and wheat head light (Senatore et al. 2021); *F. longifundum* causes brown spots on the leaves of wild leafy vegetables (Matić et al. 2020); *F. pseudoanthophilum* is highly pathogenic in corn stalks in western Iran (Chehri et al. 2010) and is associated with *Fusarium* crown rot (Kazan and Gardiner 2018); *B. ranunculi* is a complete generation of *Botrytis* species, a plant pathogen that causes damage to economically important




**Fig. 5** Colony morphology of the twelve isolated fungal species on potato dextrose agar medium incubated at 20 °C for 7 days from potato tubers infected dry rot in this study. **A** *Fusarium boothii*; **B** *Fusarium circinatum*; **C** *Fusarium citricola*; **D** *Fusarium foetens*; **E** *Fusarium iranicum*; **F** *Fusarium longifundum*; **G** *Fusarium oxysporum*; **H** *Fusarium pseudoanthophilum*; **I** *Fusarium solani*; **J** *Botryotinia ranunculi*; **K** *Clonostachys rosea*; **L** *Humicola nigrescens*

crops worldwide and is pathogenic to Ranunculaceae, which are distributed in warm and cold regions (Staats et al. 2005); *H. nigrescens* is a soil by-product of potato packaging and is known to be pathogenic to plants (UAMH 2023).

In the phylogenetic tree, which showed a close relationship between 12 strains as a result of identification, *F. solani*, generally known as the main cause of dry rot disease, was found to be far from related among other *Fusarium* spp. (Fig. 6). *F. oxysporum* (Kwon et al. 2005) is considered another major cause of dry rot disease in Korea and exhibits a high relationship through the formation of a single system with *F. foetens*, *F. pseudoanthophilum*, and *F. circinatum*. In this study, *F. acuminatum*, *F. avenaceum*, *F. crookwellense*, *F. culmorum*, *F. equiseti*, *F. sambucinum*, *F. scirpi*, *F. semitectum*, *F. sporotrichioides*, and *F. tricinctum* were not investigated in *Fusarium* (Cullen et al. 2005). *F. sambucinum* (Tiwari et al. 2020), known as a common or dominant species in North America, Europe, and China, was not identified among the isolated strains. These results suggest that *F. sambucinum* is not common in Korea among the various types of *Fusarium* strains that do not cause dry rot disease.





**Fig. 6** Phylogenetic relationship between *Fusarium* spp. and fungi isolated from potatoes from 76 fungi of 93 tubers showing symptoms of dry rot disease based on ITS region gene sequence analysis

### Pathogenicity Test of Dried Potato Rot Strain

A pathogenic test of the 12 strains identified separately showed no signs of disease in the control potatoes, while all potatoes inoculated with the 12 strains in vivo into the sclera of the Sumi variety showed dry rot disease with a cross-sectional length of  $\geq 3.5$  mm and a longitudinal length of  $\geq 2.9$  mm. The average size of the inoculation site was  $\geq 4.6$  mm in cross-sectional length in the *F. oxysporum* and *F. solani* treatment zones, which are known to be involved in potato dried rot in Korea. When comparing the cross-sectional length of the two strains known as the main fungi of dry rot disease with 3.8–5.4 mm, the cross-sectional length of *F. citricola*, *F. foetens*, *F. longifundum*, and *F. pseudoanthophilum*, pathogenicity was found to have similar diseases with no statistically significant difference. However, *F. oxysporum* and *F. solani* had pathogenic properties of dry rot disease in potatoes, but were not as pathogenic as other strains until 5 weeks in a suitable environment (15–20 °C, 99% RH). It is possible that the *F. oxysporum* and *F. solani* strains were used after more than three series of cultures, which reduced their vitality. In addition, the optimal mycelium growth temperature of *F. oxysporum* and *F. solani* is 24–25 °C (Ezrari et al. 2021; Hibar et al. 2006; Jeon et al.

2013), and it is believed that mycelium growth was relatively inactive at 15–20 °C, which is the condition used in this experiment. Among *Fusarium*-based pathogens, *F. foetens* and *F. pseudoanthophilum* are known to exhibit high relationship through their formation of a single system with *F. oxysporum*. However, strains other than *F. oxysporum* and *F. solani* have not yet been reported to be associated with dry rot disease, and there is a need to determine whether these strains develop dry rot disease. The cross-sectional length of the potato sclera inoculated with *C. rosea* was 11.48 mm and the longitudinal length was 6.9 mm, which represents the greatest disease growth among all strains. The length of the longitudinal section of all strains used in the experiment was approximately 2.9–4.8 mm in other treatment areas except for potatoes inoculated with *C. rosea*, and there was no statistically significant difference. The results of this study are similar to those of Theron and Holz (1991), who reported that *C. rosea* is also involved in dry rot disease, and it is thought that *C. rosea* may be a dominant species involved in domestic potato dry rot disease.

In the pathogenicity test, traces of strain invasion were observed in *B. ranunculi* and *H. nigrescens*. There are no reports on the involvement of *B. ranunculi* and *H. nigrescens* in dry rot disease. In particular, *H. nigrescens* is known to be non-pathogenic due to soil by-products, and in this experiment, it appeared to have been invaded through potato sclerosis. This suggests that *B. ranunculi* and *H. nigrescens* exhibited parasitic behaviour, using potato sclera as nutrients in through wounds in vivo, and indicates that they are not directly related to the development of dry rot disease.

**Author Contribution** NS Kim: investigation and data curation. SJ Hong: supervision, validation, writing, review, and editing. HS Won: review. BS Kim: conceptualization and methodology. SH Gwon: investigation, data curation, writing, and review.

**Funding** This work was carried out with the support of “Cooperative Research Program for Agriculture Science and Technology Development (Project No. RS-2021-RD009975)” Rural Development Administration, Republic of Korea.

## Declarations

**Conflict of Interest** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References


Bojanowski A, Avis TJ, Pelletier S, Tweddell RJ (2013) Management of potato dry rot. *Postharvest Biol Technol* 84:99–109

- Chehri K, Zafari D, Nurhazrati M, Salleh B, Reddy KRN, Karami E (2010) Natural occurrence of *Fusarium* species associated with root and stalk rot of maize in Kermanshah Province, Iran. *J Biol Sci* 10:795–799
- Cullen DW, Toth IK, Pitkin Y, Boonham N, Walsh K, Barker I, Lees AK (2005) Use of quantitative molecular diagnostic assays to investigate *Fusarium* dry rot in potato stocks and soil. *Phytopathology* 95:1462–1471. <https://doi.org/10.1094/PHTO-95-1462>
- Desjardins AE (2006) *Fusarium* mycotoxins: chemistry, genetics, and biology. American Phytopathological Society, St Paul, MN
- Elsherbiny EA, Amin BH, Baka ZA (2016) Efficiency of pomegranate (*Punica granatum* L.) peels extract as a high potential natural tool towards *Fusarium* dry rot on potato tubers. *Postharvest Biol Technol* 111:256–263
- Ezrari S, Radouane N, Tahiri A, Amiri S, Lazraq A, Lahlali R (2021) Environmental effects of temperature and water potential on mycelial growth of *Neocosmospora solani* and *Fusarium* spp. causing dry root rot of citrus. *Curr Microbiol* 78:3092–3103. <https://doi.org/10.1007/s00284-021-02570-1>
- Gryzenhout M, Khooa B, Landman L (2016) First report of *Fusarium boothii* from pecan (*Carya illinoensis*) and camel thorn (*Vachellia erioloba*) trees in South Africa. *S Afr J Bot* 105:158–162. <https://doi.org/10.1016/j.sajb.2016.03.003>
- He SQ, Jin XL, Wei ZQ, Zhang TY, Du X, Luo DG (2004) Isolates and identification of pathogens causing dry rot of potato tubers in Ding Xi prefecture of Gansu province. *J Yunnan Agric Univ (Nat Sci)* 19:550–552
- Hibar K, Daami-Remadi M, Jabnoun-Khiareddine H, El Mahjoub M (2006) Temperature effect on mycelial growth and on disease incidence of *Fusarium oxysporum* f.sp. *radicis-lycopersici*. *Plant Pathol J* 5:233–238. <https://doi.org/10.3923/ppj.2006.233.238>
- Jeon CS, Kim GH, Son KI, Hur JS, Jeon KS, Yoon JH, Koh YJ (2013) Root rot of balloon flower (*Platycodon grandiflorum*) caused by *Fusarium solani* and *Fusarium oxysporum*. *Plant Pathol J* 29:440–445. <https://doi.org/10.5423/PPJ.NT.07.2013.0073>
- Kazan K, Gardiner DM (2018) *Fusarium* crown rot caused by *Fusarium pseudograminearum* in cereal crops: recent progress and future prospects. *Mole Plant Pathol* 19:1547–1562. <https://doi.org/10.1111/mpp.12639>
- Kim JW (2000) Plant pathology. Daegu University Press, Daegu
- Kim GW (2020) Evaluation of 45 potato genetic resources for application of new potato varieties with excellent processing characteristics. Thesis of Master Degree, Graduate School of Kangwon National University, Korea
- Korean Statistical Information Service (KOSIS) (2021) Agricultural production (potatoes and sweet potatoes). Korea. [https://kosis.kr/statHtml/statHtml.do?orgId=101&tblId=DT\\_2KAA406\\_OECD](https://kosis.kr/statHtml/statHtml.do?orgId=101&tblId=DT_2KAA406_OECD). Accessed 20 Mar 2022
- Kwon M, Kim SY, Kim CG, Kim JS, Kim HJ, Ryu KY, Park YE, Park, CS et al (2005) Potato book. National Institute of Highland Agriculture, Rural Development Administration (RDA), Korea
- Lamprecht SC, Tewoldemedhin YT (2017) *Fusarium* species associated with damping-off of rooibos seedlings and the potential of compost as soil amendment for disease suppression. *S Afr J Bot* 110:110–117
- Maphosa MN, Steenkamp ET, Wingfeld BD (2016) Genome-based selection and characterization of *Fusarium circinatum*-specific sequences. *G3* 6:631–639. <https://doi.org/10.1534/g3.115.025817>
- Matić S, Tabone G, Guarnaccia V, Gullino ML, Garibaldi A (2020) Emerging leafy vegetable crop diseases caused by the *Fusarium incarnatum-equiseti* species complex. *Phytopathol Medit* 59:303–317. <https://doi.org/10.14601/Phyto-10883>
- National Crop Pest Management System (NCPMS) (2023) Dry rot of potato. RURAL Development Administration, Korea. <https://ncpms.rda.go.kr/npms/ImageSearchInfoR1.np?detailKey=D00000075>. Accessed 5.1.2023
- Nelson PE, Toussoun TA, Cook RJ (1981) *Fusarium: diseases, biology and taxonomy*. The Pennsylvania State University Press, University Park, PA
- Park JS, Kwon OK, Kim KC, Kim MS, Kim MH, Kim JS, Park KH, Seo IS et al (1988) Plant pathology. Hyangmoonsa, Korea
- Rural Development Administration (RDA) (2020) Agricultural technology guide (Potatoes). Korea. <http://www.nongsaro.go.kr/portal/ps/psb/psbx/cropEbookFileViewPop.ps?indexPage=26&indexBasePage=0&cropsEbookFileNo=00001&ebookCode=20>. Accessed 20 Mar 2022
- Sandoval-Denis M, Guarnaccia V, Polizzi G, Crous PW (2018) Symptomatic citrus trees reveal a new pathogenic lineage in *Fusarium* and two new *Neocosmospora* species. *Persoonia* 40:1–25. <https://doi.org/10.3767/persoonia.2018.40.01>

- Schroers HJ, Baayen RP, Meffert JP, de Gruyter J, Hooftman M, O'Donnell K (2004) *Fusarium foetens*, a new species pathogenic to begonia elatior hybrids (*Begonia x hiemalis*) and the sister taxon of the *Fusarium oxysporum* species complex. *Mycologia* 96:393–406
- Secor GA, Salas B (2001) *Fusarium* dry rot and *Fusarium* wilt. In WR Stevenson, R Loria, GD Franc, DP Weingartner, eds, Compendium of potato diseases. Ed 2. American Phytopathological Society, St Paul, MN, USA, pp 23–25
- Senatore MT, Ward TJ, Cappelletti E, Beccari G, McCormick SP, Busman M, Laraba I, O'Donnell K, Prodi A (2021) Species diversity and mycotoxin production by members of the *Fusarium tricinctum* species complex associated with *Fusarium* head blight of wheat and barley in Italy. *Intl J Food Microbiol* 358:109298. <https://doi.org/10.1016/j.ijfoodmicro.2021.109298>
- Slininger PJ, Schisler DA (2002) Spray-on fungi stop potato rot fungus. United States Department of Agriculture (USDA). *AgResearch Magazine*, USA. <https://agresearchmag.ars.usda.gov/2002/jun/fungus>. Accessed 5.1.2023
- Staats M, van Baaren P, van Kan JAL (2005) Molecular phylogeny of the plant pathogenic genus *Botrytis* and the evolution of host specificity. *Mole Biol Evol* 22:333–346. <https://doi.org/10.1093/molbev/msi020>
- Theron DJ, Holz G (1991) Dry rot of potatoes caused by *Gliocladium roseum*. *Plant Pathol* 40:302–305
- Tiwari RK, Kumar R, Sharma S, Sagar V, Aggarwal R, Naga KC, Lal MK, Chourasia KN, Kumar D, Kumar M (2020) Potato dry rot disease: current status, pathogenomics and management. *Biotech* 10:503. <https://doi.org/10.1007/s13205-020-02496-8>
- Torbati M, Arzanlou M, Sandoval-Denis M, Crous PW (2019) Multigene phylogeny reveals new fungicolous species in the *Fusarium tricinctum* species complex and novel hosts in the genus *Fusarium* from Iran. *Mycol Prog* 18:119–133. <https://doi.org/10.1007/s11557-018-1422-5>
- University of Alberta Microfungus Herbarium (UAMH) (2023) Centre for Global Microfungal Biodiversity, Dalla Lana School of Public Health, University of Toronto, Canada. <https://www.uamh.ca/details.php?id=3078>. Accessed 5.1.2023
- Wegulo SN, Valverde-Bogantes E, Bolanos-Carriel C, Hallen-Adams H, Bianchini A, McMaster N, Schmale DG III (2018) First report of *Fusarium boothii* causing head blight of wheat in the United States. *Plant Dis* 102:2646–2646
- Woo KS, Yoon JH, Han SU, Woo SY (2011) Effects of *Fusarium circinatum* on disease development and gas exchange in the seedlings of *Pinus* spp. *Res Plant Dis* 17:177–183. <https://doi.org/10.5423/RPD.2011.17.2.177>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Authors and Affiliations

Nam Sook Kim<sup>1</sup> · Sae Jin Hong<sup>1</sup> · Heon Seop Won<sup>2</sup> · Byung Sup Kim<sup>1</sup> · Se Hwi Gwon<sup>1</sup> 

✉ Sae Jin Hong  
hongsj@gwnu.ac.kr

✉ Se Hwi Gwon  
sehwi0429@nate.com

<sup>1</sup> Department of Plant Science, Gangneung-Wonju National University, Gangneung 25457, Korea

<sup>2</sup> Potato Experiment Station, Gangwondo Agricultural Research and Extension Services, Gangneung 25437, Korea