CORRECTION

Correction to: Cryptic diversity of the genus *Beauveria* with a new species from Thailand

Artit Khonsanit¹ · Janet J. Luangsa-ard¹ · Donnaya Thanakitpipattana¹ · Wasana Noisripoom¹ · Thanyarat Chaitika¹ · Noppol Kobmoo¹

© German Mycological Society and Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: *Mycol Progress* (2020) 19, 291–315.

<https://doi.org/10.1007/s11557-020-01557-9>

Figure 6 of the original version of this article contained wrong pictures which had replaced the right pictures inadvertently during the revision of the manuscript. Please find below the correct pictures for Fig. 6. The authors apologize for such mistake.

The online version of the original article can be found at <https://doi.org/10.1007/s11557-020-01557-9>

✉ Noppol Kobmoo
noppol.kob@biotec.or.th

¹ Agriculture Plant Microbe Interaction Research Team (APMT), National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency (NSTDA), 113 Thailand Science Park, Phahonyothin Rd., Khlong Nueng, Khlong Luang, Pathum Thani 12120, Thailand

Fig. 6 *Beauveria gryllotalpidicola*. **a** Pale orange stromata on Lepidoptera larva; **b**, **d** semi-immersed perithecia occurrence on the fertile part of the stromata; **c** light pink stroma on adult beetle; **e** perithecia; **f** asci with asci caps; **g** filiform ascospores; **h** multiseptate ascospores consisting of cylindrical part-spores; **i** curved hyphae and phialides with conidia on culture media; **j** conidia; **k** colony obverse on PDA in full darkness

condition for 20 days; **l** colony obverse on PDA in white light/dark cycles for 20 days; **m** colony obverse on SDA in full darkness condition for 20 days; **n** colony obverse on SDA in white light/dark cycles for 20 days. Bars: **k**, **l**, **m**, **n** = 10mm; **a**, **c** = 5 mm; **b**, **d** = 1 mm; **e** = 100 μ m; **g** = 20 μ m; **f**, **h**, **i**, **j** = 5 μ m