



Phytophthora, *Nothophytophthora* and *Halophytophthora* diversity in rivers, streams and riparian alder ecosystems of Central Europe

Tamara Corcobado^{1,2} · Thomas L. Cech² · Andreas Daxer² · Henrieta Ďatková¹ · Josef Janoušek¹ · Sneha Patra^{1,3} · Daniella Jahn² · Christine Hüttler² · Ivan Milenković^{1,4} · Michal Tomšovský¹ · Marília Horta Jung^{1,5} · Thomas Jung^{1,5}

Received: 2 March 2023 / Revised: 11 May 2023 / Accepted: 11 May 2023 / Published online: 13 June 2023
© The Author(s) 2023

Abstract

Waterways are ideal pathways for *Phytophthora* dispersal and potential introduction to terrestrial ecosystems. While many *Phytophthora* species from phylogenetic clades 6, 9 and 10 are predominant oomycetes in watercourses due to their adaptation to a lifestyle as saprotrophs and opportunistic pathogens of riparian plants, species from clades 2, 7 and 8 are predominantly soil- or airborne using aquatic habitats as temporal niches for spreading and invading terrestrial sites along the watercourses. In contrast to forest ecosystems, knowledge of *Phytophthora* diversity in watercourses in Central Europe is limited. Between 2014 and 2019 extensive surveys of streams and rivers were undertaken across Austria, in South Moravia, Czech Republic and Žilina province, Slovakia to unveil the diversity and distribution of *Phytophthora* and related oomycetes. In addition, in Austria riparian forests of black alder (*Alnus glutinosa*) and grey alder (*A. incana*) in lowlands and in the Alps were examined. A variety of *Phytophthora* species from clades 2, 6, 7, 8, 9 and 10 were isolated, with clade 6 species showing the widest distribution and abundance. Furthermore, interspecific clade 6 hybrids and other oomycetes such as *Halophytophthora fluviatilis* and undescribed *Nothophytophthora* spp. were also obtained. In riparian alders, symptoms of *Phytophthora* infections were associated with species from the *P. ×alni* complex and *P. plurivora*. *Phytophthora plurivora* was most common in alder stands whereas *P. uniformis* was the oomycete species occurring at the highest altitude in alpine riparian areas.

Keywords Oomycetes · Hybrids · Alpine · Dieback · Water filtration · *Phytophthora plurivora*

Introduction

Many species from the oomycete genus *Phytophthora* are responsible for the destruction of natural and semi-natural forests and woodlands at a global scale (Jung et al. 2018a; Brasier et al. 2022). Until the 1990s, *Phytophthora* species were mainly reported as pathogens of orchards and horticultural crops (Kreutzer et al. 1940; Deutschmann 1954; Hildebrand 1959; Tsao 1990; Bourke 1991) and infrequently as pathogens affecting forest ecosystems (Tucker and Milbrath 1942; Podger et al. 1965). However, since the 1960s the number of important diseases of forests and other natural

ecosystems caused by invasive *Phytophthora* species has increased exponentially from six to currently 41 (Brasier et al. 2022). Since water enhances the pathogen's sporulation, spread and infection via zoospores (Hirst and Stedman 1960; McIntosh 1964; Lacey 1967; Chen et al. 2022) isolations of *Phytophthora* from irrigation water in crops and nurseries were performed for monitoring and clarifying the diversity of *Phytophthora* pathogens (Klotz et al. 1959; Thomson and Allen 1974; Oudemans 1999; Bush et al. 2003). The use of surface water for irrigation can introduce *Phytophthora* spp. to nurseries and horticultural areas and facilitate the infection of plants while recirculation systems can lead to an accumulation and concentration of *Phytophthora* inoculum enabling spread of *Phytophthora* pathogens from infested to non-infested plants, thus exacerbating the pathogen's detrimental effects. Early studies on *Phytophthora* presence in surface water of forests were performed by Kliejunas and Ko (1976) and von Broembsen (1984). These showed the occurrence of *P. cinnamomi* in run-off water and standing water (i.e.

Section Editor: Tanay Bose

This article is part of the "Topical collection—Since de Bary: Progress in *Phytophthora* research"

Extended author information available on the last page of the article

puddles) and concluded that the water would act as vehicle for pathogen spread to new locations posing a potential risk for nurseries and agricultural systems. In the early 2000s, an extensive stream monitoring program has been initiated in the USA to promote early detection of *P. ramorum*, the causal agent of the devastating sudden oak death (Sutton et al. 2009; Reeser et al. 2011) and to investigate the ecology and pathways of this pathogen (Murphy et al. 2008; Hwang et al. 2008, 2011). Subsequently, also in other continents studies of *Phytophthora* diversity in watercourses were carried out (e.g. Hüberli et al. 2013; Oh et al. 2013; Matsiakh et al. 2016). First records of *Phytophthora* pathogens causing devastating disease outbreaks and mortality along rivers in European forests was associated with alder trees (*Alnus* spp.) in the mid-1990s (Brasier et al. 1995; Gibbs et al. 1999; Hartmann 1995; Streito et al. 2002; Jung and Blaschke 2004). In Europe, the first stream surveys complemented with forest surveys were performed by Hansen and Delatour (1999) in France. Simultaneous *Phytophthora* surveys in aquatic and terrestrial ecosystems have subsequently been carried out in Sicily, Sardinia, Chile, South Africa, Taiwan and Vietnam (Nagel et al. 2013, 2015; Oh et al. 2013; Jung et al. 2017a, b, c, 2018b, 2019, 2020; Bregant et al. 2020).

Among riparian trees along riversides, only a few tree species are known to be susceptible to *Phytophthora*. In natural ecosystems, *Fraxinus* spp. was recorded to suffer from dieback and mortality associated with *Phytophthora* infections (Orlikowski et al. 2011; Akilli et al. 2013). Undoubtedly, tree species from the genus *Alnus* are recognized as the most vulnerable to *Phytophthora* attacks on riparian sites. They are affected by several species of *Phytophthora*, including *P. plurivora*, *P. gonapodyides*, *P. polonica*, *P. lacustris*, *P. syringae*, *P. pseudosyringae*, *P. cactorum*, *P. siskiyouensis*, *P. hydropathica*, *P. bilorbang* and especially the *P. ×alni* species complex, i.e. *P. uniformis*, *P. ×alni* and *P. ×multiformis*, which show host-specificity to alders (Brasier and Kirk 2001; Jung et al. 2003; Jung and Blaschke 2004; Brasier et al. 2004; Štěpánková et al. 2013; Perez-Sierra et al. 2015; Trzewik et al. 2015; Matsiakh et al. 2021). More recently, *P. acerina*, *P. pseudocryptogea*, *P. ×serendipita* and the newly described *P. alpina* were reported causing disease symptoms on *Alnus* spp. (Bregant et al. 2020; Seddaiu and Linaldeddu 2020). In Europe, alder decline has been recorded and studied mainly in black alder (*A. glutinosa*) and to a much lesser extent in grey alder (*A. incana*) (Gibbs et al. 1999; Streito et al. 2002; Jung and Blaschke 2004; Černý and Strnadová 2010; Solla et al. 2010; Trzewik et al. 2015; Seddaiu and Linaldeddu 2020). However, other alder species, such as *A. viridis* and *A. cordata* are also susceptible to *Phytophthora* (Santini et al. 2001; Jung and Blaschke 2006; Bregant et al. 2020). Grey alder commonly occurs in Northern and Central Europe and is particularly widespread in alpine areas of intermediate to high altitude (Houston Durrant et al. 2016a).

Black alder is widespread in lowland areas across Europe but can also grow at higher altitudes (Houston Durrant et al. 2016b). Since it is currently little known about the extent of *Phytophthora*-mediated decline of both alder species in alpine areas (Bregant et al. 2020) it is of interest to investigate the severity of alder decline in countries which include alpine regions with a high frequency of riparian alders.

To date, knowledge of *Phytophthora* diversity and distribution within Central European waterways is still very limited. Also, little is known about *Phytophthora* distribution in riparian alder ecosystems in Austria (Cech 2006; Balci and Cech 2004). Using various isolation methods, this study aimed to unveil the *Phytophthora* species assemblage in (i) rivers, streams and associated alder forests in Austria at different time points; and (ii) forest streams of the southern region of the Czech Republic (South Moravia) and the north-western region of Slovakia (Žilina).

Materials and methods

Sampling sites

The surveys and samplings were accomplished in watercourses and in the riparian alder forests along the rivers and streams in Austria during spring and summer 2014, summer 2015, summer and autumn 2016 and autumn 2019; and in rivers and streams of South Moravia, Czech Republic, and Žilina, Slovakia, during November 2018 (Table S1 and S2, Fig. 1). Sampling along watercourses was performed at sites with easy access to the riparian forests and the banks (Table S1). In situ stream baitings were carried out in locations with reduced visibility (e.g. under bridges or overhanging branches) and where water flow was calm to facilitate zoospore attachment to baiting leaves.

In Austria, we selected 37 rivers and streams distributed across all nine states along a wide altitude gradient from lowland to alpine areas (140.0–1408.7 m a.s.l.) (Figs. S1–S8). Four rivers and one stream are eventually draining into the North Sea (i.e. Lainsitz, Ill, Alvier, Lutz and Reißbach) while the remaining rivers and streams are draining into the Black Sea. In addition, 57 riparian forest sites with the presence of black or grey alder trees were sampled. Sampling was performed by selecting 20–100 m long sections along the riverbanks containing black or grey alder trees with typical symptoms of *Phytophthora* infections such as crown defoliation and dieback, yellowing of leaves, small leaves, excessive fructification, bleeding stem cankers and rusty or tarry spots on the outer bark (Fig. 2). In 17 of the 57 riparian sites detailed assessments were performed. A total of 824 trees were examined thoroughly and classified into non-declining (less than 15% crown transparency and absence of bleeding stem




Fig. 1 Distribution of sites (a) included in the oomycete surveys of watercourses and riparian alder forests in Austria (b), Czech Republic (c) and Slovakia (d)

cankers), declining (between 15 and 95% crown transparency and/or presence of bleeding stem cankers) and dead (more than 95% crown defoliation) trees. Other riparian tree species present at the sites included *Fraxinus excelsior*, *Populus* spp., *Salix* spp. and *Ulmus* spp., while more

distant from the sites *Carpinus betulus*, *Quercus* spp., *Acer* spp., *Tilia* spp. and *Picea abies* occurred.

In both South Moravia and the Žilina regions each ten rivers and streams were selected (Table S1 and S2). The sites in South Moravia and Žilina were covering




Fig. 2 Oomycete survey of riparian black and grey alder sites and waterways in Austria, the Czech Republic and Slovakia, with emphasis on *Phytophthora* spp.: (a) crown defoliation of *Phytophthora*-infected black alders in Austria; (b) crown dieback of *Phytophthora*-infected black alder; (c) *Phytophthora*-infected grey alders along an alpine stream showing different levels of decline; (d) water filtration in Austria; (e) baiting raft floating in an Austrian stream; (f) collection of floating naturally fallen leaves at an Austrian river; (g) bleeding stem canker on a young black alder in Austria; (h) soil baiting using leaves from several tree species as baits; (i) floating naturally fallen leaves in a stream of the Czech Republic; (j) floating naturally fallen leaves in a Slovakian stream

an altitudinal range of 242–466 m and 444–775 m a.s.l, respectively. All streams are draining into the Danube and eventually into the Black Sea (Figs. S9–S10).

Sampling methods and *Phytophthora* isolation

In Austria six different sampling methods were used to provide a detailed overview of the *Phytophthora* diversity in rivers, streams and associated alder forests, i.e. in situ baiting, water filtration and collecting naturally fallen leaves, rhizosphere soil, fine roots and bark cankers.

For in situ baiting of rivers and streams, 35 × 40 cm baiting rafts, constructed from fly mesh and styrofoam, were filled each with 15–20 young leaves of the native species *Fagus sylvatica*, *Quercus robur*, *Acer* spp., *Alnus* spp., *Prunus avium* and *Aesculus hippocastanum*, tied with a rope to a shrub or tree at the riverbank and left for 2–3 days floating on the watercourses (Reeser et al. 2011; Jung et al. 2017a, b, c, 2018a, b, 2019). This method was used at 50 sites along 28 watercourses. In addition, naturally fallen necrotic leaves were collected from the water surface at 18 sites along 16 watercourses.

In South Moravia, Czech Republic, and the Žilina region of Slovakia each 30 naturally fallen necrotic leaves were collected from the water surface of the 20 forest streams. In South Moravia the leaves belonged to *Acer* spp., *Alnus* spp., *Carpinus betulus*, *Cornus* spp., *Corylus avellana*, *Fraxinus excelsior*, *Populus* spp., *Prunus* sp., *Quercus* spp., *Robinia pseudacacia*, *Salix* spp., *Tilia* spp. and *Ulmus* spp.. In Slovakia leaves of *Acer* spp., *Alnus* spp., *Corylus avellana*, *Fagus sylvatica*, *Populus* spp., *Prunus* sp., *Salix* spp. and *Picea abies* were sampled.

Collected baiting leaves and naturally fallen leaves were transported in cool boxes to the laboratory. Then the leaves were blotted dry, cut into small pieces and plated onto selective PARPNH agar (V8 juice agar (V8A) amended with 10 µg mL⁻¹ pimaricin, 200 µg mL⁻¹ ampicillin, 10 µg mL⁻¹ rifampicin, 25 µg mL⁻¹ pentachloronitrobenzene (PCNB), 50 µg mL⁻¹ nystatin and 50 µg mL⁻¹ hymexazol) (Jung et al. 2017a, b, c). The Petri dishes were kept at 20 °C in the dark and checked for growing oomycete hyphae after 24–48 h. Oomycete hyphae were transferred from the growing margins onto V8A.

Water filtrations, according to Redondo et al. (2018ab), were performed at 29 sites along 13 rivers in Austria. At each site, 10 L of water were collected and pumped with an agricultural 5 L volume hand sprayer through a 8 µm membrane (Merck Millipore, Cork, Ireland) inserted into a polysulfone filter holder (Whatman™ Polysulfone Filter Holders, Syringe Type, Krackeler scientific, New York, USA). Several membranes were used to filter 10 L since particle accumulations obstructed the membranes. Between sampling sites, the pump was washed and disinfested by rinsing it several times with a mixture of 5% sodium hypochlorite and distilled water to avoid possible cross-contamination. The used membranes were stored in plastic bags and transported in a cool box to the laboratory where they were stored at 6 °C until processing on the following day. Membranes were blotted dry, cut into 1 × 1 cm pieces and plated onto selective PARPNH agar. The Petri dishes were kept at 20 °C in the dark and oomycete hyphae were subcultured as described before.

In Austria, rhizosphere soil sampling and oomycete isolations were performed following the methodology of Jung (2009). In the selected sections of declining riparian alder forests, symptomatic and/or asymptomatic trees were sampled by collecting at each sample tree soil monoliths from 3–4 cardinal directions in 30–50 cm distance from the stem base and at a depth of 10–30 cm avoiding the superficial organic layer, and bulking them to a composite sample of ca 1–2 L. In total 64 composite rhizosphere soil samples were transported to the lab. Well-mixed subsamples of circa 300 mL per composite soil sample were transferred to plastic containers and flooded with distilled water so that the water surface was approximately 3 cm above the soil surface. After cleaning the water surface from floating particles with kitchen paper tissue, young leaflets of the same native tree species used in the baiting rafts were placed as baits to float on the water surface. Infected necrotic leaflets were then blotted dry, cut into small fragments and plated onto selective PARPNH agar. Petri dishes were kept at 20 °C in the dark and oomycete hyphae were subcultured as described before.

Active bleeding bark lesions and, infrequently, also fine roots of declining alder trees were sampled according to Jung and Blaschke (2004) and Jung (2009). The bark samples, including the cambium, were taken from the upper section of fresh lesions using a hammer and a chisel and transported in wide-mouth bottles with distilled water to the lab. The water was exchanged 3–4 times per day over two days to remove excess polyphenols. Then small pieces of bark were cut from active parts of the lesions, blotted dry and plated onto selective PARPNH agar. Furthermore, fine roots from rhizosphere soil samples of 2–3 trees were collected at five declining alder sites. Roots were cleaned under running water, blotted on filter paper and small sections plated onto selective PARPNH agar.

Petri dishes were kept at 20 °C in the dark and oomycete hyphae were subcultured as described before.

Species identification of oomycete isolates was performed using molecular sequence analyses (see below). For isolates identified as *P. ×alni* / *P. ×multiformis*, a morphological characterisation of 3 weeks old V8A cultures at ×320 under the light microscope was performed to distinguish between *P. ×alni* and *P. ×multiformis* (Brasier et al. 2004).

Molecular identification

DNA extraction was performed using a magnetic bead-based technology. The protocol “BOMB TNA extraction from plants using TNES/GITC lysis” (high throughput protocol; Oberacker et al. 2019) was modified to extract DNA from the isolates (Jung et al. 2022). Mycelium from a colony growing on nutrient media for one up to three weeks was collected and transferred to 2 mL homogenisation tubes (116910500, MP Biomedicals; Irvine, USA) which were stored at -80 °C until further use. Homogenisation was performed at 5 800 RPM 4 × 15 s using Precellys Evolution instrument (Bertin Technologies; Montigny-le-Bretonneux, France). The samples were centrifuged briefly to collect the content at the bottom and 60 µl of TNES buffer with RNase A (T3018L; New England Biolabs; Ipswich, USA) was added. The mixture was incubated for 20 min at 56 °C. 240 µl of 1.5 × GITC lysis buffer was added and mixed thoroughly. The mixture was centrifuged for 10 min at 10 000 RPM and 180 µl of supernatant was transferred to a deep-well block (82.1970.002; Sarstedt; Nümbrecht, Germany). DNA was further mixed with 120 µl of AMPure XP beads (A63881; Beckman Coulter; Indianapolis, USA) and 240 µl of isopropanol and incubated for 10 min. All washing steps were performed according to the original protocol. DNA was eluted in 70 µl of TE buffer (12090015; Invitrogen; Carlsbad, USA) and stored at -80 °C for long-term preservation.

Initial identification was performed with ITS gene region sequence analysis using ITS1/4 (White et al. 1990) or ITS6/4 (Cooke et al. 2000) primers. The *cox1* and β -tubulin genes were sequenced using OomCoxI-Levup/OomCoxI-Levlo (Robideau et al. (2011) and TUBUF2/TUBUR1 (Kroon et al. 2004), respectively. PCR conditions and gel electrophoresis were as described by Jung et al. (2022). PCR products were purified and sequenced by Eurofins Genomics (Ebersberg, Germany) in both directions with the primers used for PCR amplification. DNA sequence data from representative isolates from this study were deposited in GenBank and their accession numbers are given in Table S3.

Results

Phytophthora diversity and distribution in rivers and streams of Austria, South Moravia (Czech Republic) and Žilina (Slovakia)


In Austria, 10 *Phytophthora* taxa and three *Phytophthora* hybrid taxa belonging to phylogenetic clades 2c, 6b, 7a and 9a, as well as *Phytophythium litorale* and a *Halophytophthora* species were obtained from 29 of 37 rivers and streams (78.4%) corresponding to 45 of 95 sites (47.4%) (Table S1; Fig. 3). In addition, several *Pythiaceae* species were isolated (e.g. *Elongisporangium undulatum*). Using baiting rafts, from 22 of the 28 rivers (78.6%) and at 32 of the 50 sites (64%) where this method was used, seven *Phytophthora* species including *P. chlamydospora*, *P. gonapodyides*, *P. hydropathica*, *P. lacustris*, *P. plurivora*, *P. riparia* and *P. uniformis* and two hybrid taxa, i.e. *P. taxon × riparia* and *P. chlamydospora × lacustris*, were isolated. From naturally fallen floating leaves four *Phytophthora* taxa, two hybrids and one *Halophytophthora* species were isolated from 14 of the 16 rivers (87.5%) and at 15 of the 18 sites (83.3%) surveyed by this method. The species assemblage included *P. bilorbang*, *P. gonapodyides*, *P. lacustris*, the new yet undescribed *P. taxon bilorbang-like 2*, *P. taxon × lacustris*, *P. taxon × riparia* and *Halophytophthora fluviatilis*. Using the water filtration method, *P. chlamydospora*, *P. gonapodyides*, *P. lacustris*, *P. plurivora*, *P. riparia* and *P. taxon × riparia* were obtained from 9 of 13 rivers (69.2%) and at 19 out of 29 sites (65.5%) where this isolation method was applied.

Phytophthora species from phylogenetic clade 6b with a primarily aquatic lifestyle showed the widest distribution in aquatic ecosystems across Austria. Using the three sampling methods they were found in 78.4% of the 37 rivers and streams. Overall, the clade 6b species *P. gonapodyides* (64.9% of all waterways) and *P. lacustris* (62.2%) were the most widespread species followed by *P. plurivora* from clade 2c (18.9%) and the clade 6b hybrid *P. taxon × riparia* (13.5%). All other *Phytophthora* species had a scattered distribution.

The Austrian watercourses with the highest numbers of *Phytophthora* taxa (Fig. 4) were the Kamp (6 taxa = 46.2% of the total number of taxa) and the Krumme Steyerling (5 taxa; 38.5%) followed equally by Traun, Enns, Zeller Ache (Ager) and Stempfelbach (4 taxa; 30.8%).

Phytophthora gonapodyides occurred between 141.1 and 1292.8 m a.s.l. and, hence, was the species with the widest altitudinal amplitude and the highest elevation, specifically in Thomatalerbach (1292.8 m), Leutascher Ache (1096.7 m) and Mur (1137.0 m). *Phytophthora lacustris* (140.0–690.8 m), *P. taxon × lacustris* (141.1–348.1 m) and *P. plurivora* (240.9–690.8 m) had a narrower altitudinal

Fig. 3 Aquatic diversity of *Phytophthora*, *Halophytophthora* and *Nothophytophthora* species in watercourses and riparian sites in Austria, Czech Republic and Slovakia; BIL = *P. bilorbang*, BIL2 = *P. taxon bilorbang-like 2*, CHL = *P. chlamydospora*, ×CHL = *P. chlamydospora* × *lacustris*, FLU = *Halophytophthora fluviatilis*; GAL = *P. gallica*; GON = *P. gonapodyides*, HYD = *P. hydropathica*, LAC = *P. lacustris*, ×LAC = *P. taxon* × *lacustris*; NOT1 = *Nothophytophthora* taxon 1; NOT2 = *Nothophytophthora* taxon 2; PLU = *P. plurivora*, RIP = *P. riparia*, ×RIP = *P. taxon* × *riparia*, SYR = *P. syringae*, UNI = *P. uniformis*




amplitude. The new undescribed *P. taxon bilorbang-like 2* was recovered at an altitude of 140.4 m.

In South Moravia, all 10 sampled streams (10 sites) were positive for *Phytophthora* using naturally fallen leaves as baits. In total six *Phytophthora* taxa from clades 6b, 8d

and 10 and one hybrid taxon from clade 6b were found (Table S1; Fig. 3). *Phytophthora gonapodyides* and *P. lacustris* were present in all watercourses followed by the clade 10 species *P. gallica* (3 streams) and *P. bilorbang* from clade 6b (2 streams) while *P. syringae* from clade 8d and the

Fig. 4 Numbers of watercourses in Austria, Czech Republic and Slovakia with different total numbers of oomycete species



clade 6b taxa *P. riparia* and *P. lacustris* × *riparia* occurred only in one stream each. In addition, a new yet undescribed *Nothophytophthora* species, informally designated as *Nothophytophthora* taxon 2, and *H. fluviatilis* were obtained from one and two streams, respectively. *Phytophythium citrinum*, *Phy. litorale* and other undescribed *Phytophythium* taxa were also present in these streams. The streams with the highest number of *Phytophthora* taxa (Fig. 4) were the Kuřimka (6 taxa = 85.7% of the total number of species) and the Punkva (5 taxa; 71.4%) followed by Časnýř and Říčka streams (3 taxa; 42.9%). The ten streams were within a narrow altitudinal range of 242–466 m.

In the Žilina region of Slovakia, three *Phytophthora* species from clade 6b were isolated from 8 of the 10 surveyed streams/sites (Table S1; Fig. 3). In one of the streams where *Phytophthora* was not found, the new yet undescribed *Nothophytophthora* taxon 1 was obtained. *Phytophthora gonapodyides* occurred in all eight *Phytophthora*-positive streams (80% of all streams) while in the western and north-western parts of Žilina *P. lacustris* and *P. bilorbang* were present in two and three streams, respectively. Other oomycete taxa included *Phy. litorale*, *Pythium kashmirensense* and other undescribed *Phytophythium* and *Pythium* spp. The highest numbers of *Phytophthora* taxa (Fig. 4) were found in the streams Črchľový potok and in Predmieranka (3 taxa = 100% of the total number of taxa). From Predmieranka stream with the highest altitude (775.0 m) of all streams sampled in Slovakia, *P. gonapodyides*, *P. lacustris* and *P. bilorbang* were isolated.

***Phytophthora* diversity and distribution in riparian alder forests in Austria and association with disease symptoms**

Alder trees showed decline symptoms in most of the selected 57 riparian alder forest sites. Across the 17 sites which were studied in detail, on average 36.0% of the 824 assessed *Alnus* trees were showing crown defoliation and/or stem cankers, 11.6% were dead and 52.4% were healthy. Overall, 9 *Phytophthora* taxa belonging to phylogenetic clades 1a, 2c, 6b, 7a, 8d and 9a were present in rhizosphere soil or bark cankers at 26 of the 57 riparian black alder and grey alder forest sites sampled in Austria (Table S1). Six *Phytophthora* species were isolated from rhizosphere soil samples of 26 of the 64 trees sampled (40.6%) including *P. cactorum*, *P. gonapodyides*, *P. lacustris*, *P. plurivora*, *P. syringae* and *P. uniformis*. Two *Phytophthora* species, i.e. *P. lacustris* and the clade 9a species *P. polonica*, were isolated from fine roots of 3 of the 25 trees sampled (12%). Three alder-specific *Phytophthora* species from clade 7a were isolated from bark lesions of 17 of the 38 trees sampled (44.7%), including *P. ×alni* (5 sites, 7 trees), *P. ×multiformis* (2 sites, 2 trees) and *P. uniformis* (5 sites, 5 trees).

The 57 riparian forest sites were associated with 30 rivers and streams. With 15 alder forest sites along 14 rivers/streams *P. plurivora* showed the widest distribution of all species (Fig. 5). *Phytophthora plurivora* was recovered from rhizosphere soil samples but not from bark cankers. With nine riparian forests associated with nine rivers/streams *P. ×alni*, *P. ×multiformis* and *P. uniformis* were also widespread. Except for *P. uniformis*, which was also obtained from a rhizosphere soil sample, they were exclusively isolated from alder bark cankers. *Phytophthora ×alni* and *P. uniformis* occurred each in five riparian forest sites along five rivers/streams, co-occurring at the river Inn (different sites) and the stream Marbach (same site). *Phytophthora ×multiformis* was found in two forest sites at two rivers. *Phytophthora gonapodyides* and *P. lacustris* were each recovered from three riparian forest sites at three rivers, co-occurring at the river Drau. While *P. gonapodyides* was only isolated from rhizosphere soil, *P. lacustris* was obtained from both fine roots and rhizosphere soil. *Phytophthora* species from phylogenetic clades 1a (*P. cactorum*), 8d (*P. syringae*) and 9a (*P. polonica*) occurred only in each one riparian forest site.

The highest numbers of *Phytophthora* species were found in the alder forest sites associated with the river Inn (4 species = 44.4% of the total number of species) followed by the rivers Enns and Salzach (3 species; 33.3%), and the Danube, Drau, Lungitzter Loben, Marbach, Reißbach and Weizbach (2 species; 22.2%) (Table S1).

The most widespread and common species, *P. plurivora*, occurred at forest sites with an altitudinal range of 144.5–793.7 m. *Phytophthora uniformis* was the species with the widest altitudinal amplitude (311.8–1266.3 m) occurring even at the highest forest sites along the streams Turrach (1266.3 m) and Gschnitzbach (1081.9 m). The other two clade 7a species, *P. ×alni* (311.8–782.4 m) and *P. ×multiformis* (149.3–504.3 m), were restricted to considerably lower altitudes. Interestingly, although *P. gonapodyides* showed the widest altitudinal range and the highest altitude in the stream surveys, in alder forest sites it was only found between 495.8 and 730.9 m a.s.l.

Generally, *Phytophthora* was isolated from declining alder trees where typical symptoms of infection were noted. The most affected states with a substantial number of observations of declining alders were Styria, Tyrol and Salzburg. Lower Austria, Upper Austria and Burgenland showed an intermediate incidence of disease while only healthy alder stands were recorded in the state of Vorarlberg. The most common and widespread *Phytophthora* species in the riparian alder forests was *P. plurivora*; it is notable that its isolation was restricted to rhizosphere soil samples whereas active bleeding cankers did not yield *P. plurivora*. Nonetheless, in most cases, an intermediate to a high incidence




Fig. 5 Diversity and frequency of *Phytophthora* species isolated from rhizosphere soil and bark cankers of alder trees in riparian forests in Austria; ×ALN=*P. ×alni*, CAC=*P. cactorum*, GON=*P. gonapody-*

ides, LAC=*P. lacustris*, ×MUL=*P. ×multiformis*, PLU=*P. plurivora*, POL=*P. polonica*, SYR=*P. syringae*, UNI=*P. uniformis*

of alder decline was associated with its presence (e.g. sites 100 and 109 with 20 and 100% declining or dead alder trees, respectively). At grey alder site 50 on the banks of the river Inn (Tyrol), where *P. plurivora* was isolated from the rhizosphere soil of a declining tree, 140 out of 200 trees were declining with 20% of them showing old inactive stem cankers with tarry spots. *Phytophthora ×multiformis* was exclusively isolated from two sites with high incidences of alder mortality (i.e. sites 4 and 128 with 18 and 52% dead alder trees, respectively), co-occurring with *P. ×alni* at one of the sites. Sites, where *P. ×alni* (e.g. sites 131 and 50) or *P. uniformis* (e.g. site 87) occurred, were usually linked with intermediate to high levels of alder decline. Co-occurrence of *P. ×alni* and *P. uniformis* caused particularly high mortality (e.g. site 129 with 58% mortality).

Discussion

This study provides for the first-time insights into the *Phytophthora* diversity of watercourses in Central Europe, namely in Austria and in each one region of Czech Republic

(South Moravia) and Slovakia (Žilina), respectively. It also shows the first extensive overview of *Phytophthora* occurrence in riparian black and grey alder forests across Austria. *Phytophthora* diversity was higher in the 95 watercourses as compared to the 57 terrestrial alder ecosystems surveyed in Austria. This is a common pattern in combined aquatic and terrestrial *Phytophthora* surveys (Oh et al. 2013; Jung et al. 2017a, 2018b, 2019, 2020; Seddaiu et al. 2020) as watercourses provide ideal niches for aquatic *Phytophthora* species and also temporal niches for various soil- and air-borne *Phytophthora* species. Aquatic *Phytophthora* species are mainly saprotrophs on floating and submerged organic debris which sporadically can establish in terrestrial ecosystems as comparatively weak opportunistic tree pathogens. Soilborne *Phytophthora* species spend most of their life cycle as root and bark pathogens of terrestrial plant species but most of them can also survive in watercourses as saprotrophs and spread downstream to other terrestrial ecosystems. Among those *Phytophthora* species found in the watercourses in Austria, Czechia and Slovakia, typical aquatic clade 6 species were most common and abundant. Aquatic clade 6 species are able to colonize green leaves

and dead or decomposing leaf litter (Brasier et al. 2003; Jung et al. 2011) and due to their high competitiveness overtake the colonization of other species (Aram and Rizzo 2018). Other aquatic species were *P. hydropathica* (clade 9) and *P. gallica* (clade 10) which occurred in Austria and South Moravia, respectively. The other *Phytophthora* species recovered from watercourses in Central Europe belong to clades 2, 7, 8 and 9 and are known to have a predominantly soilborne lifestyle. *Phytophthora plurivora* from clade 2 occurred in seven Austrian rivers/streams and has previously been recorded from rivers and streams in Chile, Ireland, Italy and the USA (Reeser et al. 2011; O' Hanlon et al. 2016; Jung et al. 2018b, 2019; Seddaiu et al. 2020; Riolo et al. 2020), demonstrating its ability to survive and spread in aquatic ecosystems. Its widespread occurrence in watercourses is concerning due to its high aggressiveness to a wide range of tree species and its involvement in oak and beech declines across Europe (Jung 2009; Jung and Burgess 2009; Jung et al. 2018a).

The stream surveys in South Moravia and in Žilina revealed the presence of several aquatic *Phytophthora* species and also one terrestrial species, *P. syringae*, in one South Moravian stream. Seasonality of these oomycetes (cf. Hwang et al. 2008, 2011) is most likely responsible for the lower *Phytophthora* diversity in these regions as compared to Austria since the surveys in Czechia and Slovakia were performed only in November while in Austria the surveys were performed in different seasons (spring, summer and autumn). The high prevalence of *P. gonapodyides* and *P. lacustris* in watercourses of Central Europe is in accordance with previous surveys performed in the Mediterranean (Jung et al. 2019; Seddaiu et al. 2020; Riolo et al. 2020) and Northern Europe (Redondo et al. 2018a). Both species are considered native to Europe and widespread on several continents (Greslebin et al. 2005; Reeser et al. 2011; Huai et al. 2013; Nagel et al. 2013; Shrestha et al. 2013; Stamler et al. 2016; Jung et al. 2018b; Xu et al. 2019). *Phytophthora lacustris* is predominantly an aquatic saprotroph on plant debris but can also act as opportunistic pathogen of riparian trees and irrigated trees and crops (Orlikowski et al. 2011; Nechwatal et al. 2013; Jung et al. 2016; Kanoun-Boulé et al. 2016). Soil infestation assays with *P. lacustris* have also shown moderate aggressiveness to several tree species (Nechwatal et al. 2013; Milenković et al. 2018). In contrast, *P. gonapodyides* has a wider range of habitats than *P. lacustris* (Jung et al. 1996; Hansen and Delatour 1999; Reeser et al. 2011; Seddaiu et al. 2020) and besides being an aquatic saprotroph has also been reported as virulent root and bark pathogen on mature forest trees and in pathogenicity tests (Jung et al. 1996, 2000; Jung 2009; Corcobado et al. 2010, 2020; Belisario et al. 2016; Cleary et al. 2016). It is notable that *P. gonapodyides* could also be isolated during the coldest months as in the present and previous studies

(Hansen and Delatour 1999; Hwang et al. 2011) demonstrating its ecological advantage to proliferate during all seasons. Moreover, *P. gonapodyides* showed the widest altitudinal amplitude (141.1–1292.8 m) and reached the highest altitudinal location surveyed in the Austrian Alps. Another clade 6 species, *P. bilorbang*, was only found in one river (2.7%) in Austria while it was present in two rivers (20%) surveyed in South Moravia and in Žilina. Also, in all three countries *P. bilorbang* was isolated only from fallen leaves floating on the waterbodies during autumn. This pathogen has been found previously in aquatic ecosystems in Italy and the USA (Reeser et al. 2011; Seddaiu et al. 2020) and is associated with declining maquis shrub species (i.e. *Juniperus phoenicea* and *Pistacia lentiscus*), riparian vegetation (i.e. *Salix* sp. and *A. glutinosa*) (Scanu et al. 2015; Riolo et al. 2020; Matsiakh et al. 2021) and with root rot in olive (*Olea europaea*) orchards (Santilli et al. 2020). *Phytophthora bilorbang* has very similar ITS sequences as *Phytophthora* taxon Oaksoil and *Phytophthora* taxon Riversoil, both informally designated in 2003 (Brasier et al. 2003). However, *P. bilorbang* has a homothallic breeding system whereas the other two taxa are sterile, considered as an adaptation to a fully aquatic saprotrophic lifestyle (Brasier et al. 2003; Jung et al. 2011). Interestingly, a previously unknown taxon closely related to *P. bilorbang*, informally designated *P. taxon bilorbang-like2*, was also isolated from floating leaves in Austria during autumn. Another common clade 6 species in the waterways of Austria and South Moravia was *P. riparia*, a widespread aquatic saprotroph in the Northern hemisphere which is often involved in hybridization processes (Brazee et al. 2016; Stamler et al. 2016; Bily et al. 2018, 2022), also demonstrated by the findings of the hybrids *P. taxon × riparia* and *P. lacustris × riparia* in Austrian and Czech streams, respectively. Inoculation tests with *P. riparia* and several crop species have demonstrated lack of pathogenicity (Stamler et al. 2016). *Phytophthora chlamydospora*, another clade 6 species, was exclusively detected in three Austrian watercourses. Considered as native in North America, *P. chlamydospora* has been found widespread in all continents except Antarctica (Greslebin et al. 2005; Jung 2009; Reeser et al. 2011; Huai et al. 2013; Oh et al. 2013; Hansen et al. 2015; Sims et al. 2015; Jung et al. 2016, 2017a, 2018b, 2019, 2020; Bily et al. 2018, 2022; Xu et al. 2019; Corcobado et al. 2020; Seddaiu et al. 2020). Although *P. chlamydospora* has a predominantly waterborne saprophytic lifestyle (Brasier et al. 2003; Hansen et al. 2015; Sims et al. 2015; Bily et al. 2018, 2022), it can act as an opportunistic tree pathogen in natural ecosystems (Bily et al. 2022) and as aggressive pathogen in orchards and nurseries (Derviş et al. 2016; Jung et al. 2016; Browne et al. 2020, 2021). In addition, a hybrid between *P. chlamydospora* and *P. lacustris*, *P. chlamydospora × lacustris*, was recovered from one Austrian river. *Phytophthora hydropathica* from a fast-growing,

high-temperature tolerant cluster of species in clade 9 was only isolated once from the Danube river in Upper Austria. It is a typical aquatic saprotroph and opportunistic pathogen of crops which is widespread in the USA (Hong et al. 2010; Shrestha et al. 2013) and was recently found in streams in Spain, Sicily and Sardinia (Pintos et al. 2016; Jung et al. 2019; Seddaiu et al. 2020). It is expected that the activity of this species in Central Europe will increase with warming climate. The only species from Clade 10 recovered in this study was *P. gallica*, occurring in 30% of the streams surveyed in South Moravia. It has been documented inhabiting waterways and riparian terrestrial ecosystems, causing occasional damage as root pathogen (Jung and Nechwatal 2008; Sims et al. 2015; Redondo et al. 2018a; Christova 2022).

Interspecific hybridization is increasingly recognised as driving evolutionary force in the genus *Phytophthora* facilitating adaptation to new or changing environmental conditions and enlarging host ranges or enabling host jumps, occasionally leading to speciation (Brasier 2000; Bertier et al. 2013; Nagel et al. 2013; Burgess 2015; Jung et al. 2017a, b, 2018b; Van Poucke et al. 2021). Aquatic environments seem to be a particularly favourable habitat for interspecific hybridizations as demonstrated by an array of aquatic hybrids within clades 6, 7 and 9 (Nagel et al. 2013; Yang et al. 2014; Burgess 2015; Brazee et al. 2016; Jung et al. 2017a, b, 2018a). Also in the present study, several clade 6 hybrids were obtained from watercourses (i.e. *P. chlamydospora* × *lacustris*, *P. taxon* × *lacustris* and *P. taxon* × *riparia*).

Nothophytophthora, a sister genus of *Phytophthora* within the *Peronosporaceae*, was recently described (Jung et al. 2017c) and members from this genus have since been reported in waterbodies and terrestrial ecosystems (Jung et al. 2018a; O'Hanlon et al. 2021; Landa et al. 2021). Also in this study, two new *Nothophytophthora* taxa, informally designated as *Nothophytophthora* taxon 1 and *Nothophytophthora* taxon 2, were obtained from each one stream in Slovakia and Czechia, respectively. Other oomycete genera from the families of the *Pythiaceae* (*Elongisporangium*, *Pythium*) and the *Peronosporaceae* (*Halophytophthora*, *Phytophythium*) were frequently present but only a few were subcultured and sequenced as they were not the goal of this study. *Phytophythium litorale* was recovered from Central European waterbodies and has a global distribution (e. g. Choudhary et al. 2016; Jung et al. 2020). It has also been identified as an aggressive pathogen of Oriental plane trees (*Platanus orientalis*) in Turkey (Derviş et al. 2020). *Elongisporangium undulatum* was isolated from a baiting raft in the Danube. The species is known as soilborne fine root pathogen from natural ecosystems worldwide (Jung et al. 1996, 2000, 2017a, 2018b) and was less often reported from waterbodies (O'Hanlon et al. 2018). The genus *Halophytophthora* is closely related to *Phytophthora* and *Nothophytophthora* and comprises mainly species from marine and

brackish-water habitats (Nigrelli and Thines 2013; Maia et al. 2022) but they can also tolerate low salinity and even inhabit freshwater ecosystems (Hüberli et al. 2013; Yang and Hong 2014). *Halophytophthora fluviatilis* was described from rivers and streams in the eastern USA (Yang and Hong 2014). It was also found widespread in rivers and streams in eastern Spain and its pathogenicity to different Fagaceae species and black alder was demonstrated (Caballol et al. 2021). In the present study, *H. fluviatilis* was recovered from naturally fallen autumn leaves in lowland rivers of Austria and Czechia, demonstrating its capacity to survive as saprotroph in freshwater environments.

The riparian alder forests surveyed in Austria showed an intermediate level of decline and low mortality rates. In Austria alder dieback associated with *Phytophthora* in Austria was first detected in 1996. Subsequently, an extensive alder survey was performed between 1996 and 1999 and the presence of the so-called 'alder-*Phytophthora*' was confirmed at a few scattered sites along the watercourses Reibach, Thaya, Kamp and Danube in Lower Austria and Upper Austria (Cech and Brandstetter 1999). A later alder survey performed in 50 sites in the state of Vienna showed a low percentage of mortality (3.74%) which was mostly associated with drought except for ten sites where collar rot cankers were detected and *Phytophthora* spp. confirmed as causal agents (Cech 2006). More extensive surveys revealed widespread occurrence of alder dieback in Austria, with higher disease incidences in Burgenland and Styria (Balci and Cech 2004). The assessments of the present study (2014–2019) demonstrated high mortality levels and chronic decline in the state of Styria and, to a lesser extent, in the states of Tyrol and Salzburg. In some parts of Styria, the decline was already extremely advanced with predominance of secondary pathogens, so that *Phytophthora* spp. could only rarely be isolated (i.e. along streams Turrach and Marbach). It was also noticed that these areas were associated with disturbances such as the construction of water channels which might have affected the water regime and vitality of alders. Since in these states several diseased alder plantings were found it is likely that *P. × alni* and other *Phytophthora* spp. were introduced with infested planting stock as demonstrated in previous studies (Jung and Blaschke 2004; Jung et al. 2016). Similar observations regarding infested nursery stock as potential pathway of *P. × alni* to natural riparian alder stands in Austria were made by Balci and Cech (2004). Other Central European countries, such as Germany, Czechia and Poland, also reported *Phytophthora*-related alder dieback during the 1990s (Hartmann 1995; Jung and Blaschke 2004) or the 2000s (Černý et al. 2003; Nagy et al. 2003; Orlikowski et al. 2003; Černý et al. 2008). In these surveys the distribution of diseased alders was mainly observed in lowland forests while alpine riparian forests were either overlooked or not

investigated in detail (Cech and Brandstetter 1999; Jung and Blaschke 2004). Later, monitoring of declining alder sites at alpine sites in Italy produced inconclusive results as symptoms but no primary agent were found (Pisetta et al. 2012). However, recently the presence of declining grey and green alders in the Italian Alps was found associated with *Phytophthora* infections (Bregant et al. 2020). This study recorded the presence of declining alders up to 1400 m altitude with *P. uniformis* being isolated from bark cankers and rhizosphere soil at altitudes of 1266 m and 1081.9 m, respectively. This is in accordance with results from the present study and with previous studies (Adams et al. 2008; Štěpánková et al. 2013; Redondo et al. 2015) which showed prevalence of *P. uniformis* at higher altitudes while *P. ×alni* was exclusively found in the lowlands. In Italy, a variety of *Phytophthora* species, including the new species *P. alpina*, were isolated from rhizosphere and stem cankers from grey and green alder at elevations between 1412 and 1803 m, but remarkably no species from the *P. ×alni* complex was found (Bregant et al. 2020).

This study revealed a diverse assemblage of nine *Phytophthora* species associated with riparian alder stands in Austria, i.e. *P. cactorum*, *P. gonapodyides*, *P. lacustris*, *P. plurivora*, *P. polonica*, *P. syringae*, *P. uniformis*, *P. ×alni* and *P. ×multiformis*, which were also recorded in previous alder surveys elsewhere (Jung and Blaschke 2004; Štěpánková et al. 2013; Jung et al. 2013, 2018a; Trzewik et al. 2015; Aday Kaya et al. 2018). In previous surveys performed in Austria, *P. ×alni*, *P. ×multiformis* and *P. plurivora* were isolated from bark cankers of alders in areas with common occurrence of alder plantations (Balci and Cech 2004; Cech 2006; Jung et al. 2013). Also in this study, the presence of *P. ×multiformis* was related to a riparian *A. incana* plantation in Vienna state. It appears that the occurrence of *P. ×multiformis* is often related to the recent introduction with infected nursery plants (Jung and Blaschke 2004; Štěpánková et al. 2013; Jung et al. 2016) which might be the reason for its infrequent occurrence in regions with rare alder planting activities (Nagy et al. 2003; Pintos et al. 2016). The most common *Phytophthora* species in Austrian alder stands in this study was *P. plurivora*. This wide-host range pathogen (previously designated as *P. citricola*; Jung & Burgess 2009) is widespread in the Northern hemisphere and one of the most important invasive pathogens in European forests. Due to its high aggressiveness *P. plurivora* is a major driver of declines and diebacks of beech and oak stands across Europe and is also responsible for the mortality of many alders and other tree species (Jung et al. 1996, 2000, 2013, 2016, 2018a, 2019; Hansen and Delatour 1999; Jung and Blaschke 2004; Jung and Burgess 2009; Mrazkova et al. 2013; Trzewik et al. 2015; Seddaiu and Linaldeddu 2020; Corcobado et al. 2020; Matsiakh et al. 2021). Its pathogenicity to alders, beech and oaks has been demonstrated in numerous trials (Jung et al.

1996; Fleischmann et al. 2002, 2004; Jung and Nechwatal 2008; Jung 2009; Zamora-Ballesteros et al. 2017; Corcobado et al. 2022). Although *P. plurivora* has been isolated previously from alder bark cankers in Austria, Bavaria and Italy (Jung and Blaschke 2004; Cech 2006; Bregant et al. 2020) it was recovered only from alder rhizosphere in the present study. The altitudinal range at which *P. plurivora* was found in this study was 144.5 to 793.7 m, confirming previous reports from beech and alder stands in Austria, Bavaria and Czech Republic (Jung and Blaschke 2004; Mrázková et al. 2013; Corcobado et al. 2020). In contrast, in the Italian Alps *P. plurivora* has been isolated from alder bark cankers and from rhizosphere at an altitude of 1412 m (Bregant et al. 2020). In one Austrian alder stand *P. plurivora* co-occurred in rhizosphere soil with *P. syringae* which has previously been found associated with alders as a weak pathogen (Trzewik et al. 2015). Isolation tests from alder fine roots yielded exclusively *P. polonica* and *P. lacustris*. *Phytophthora polonica* is considered native to Europe and has been mainly isolated from rhizosphere soil in declining riparian alder forests (Belbahri et al. 2006; Matsiakh et al. 2021), declining poplar plantations (Milenković et al. 2018) and declining *Q. robur* forests (Jankowiak et al. 2014). Soil infestation tests demonstrated its high aggressiveness to poplar roots (Milenković et al. 2018) while under-bark inoculation tests showed only mild pathogenicity to poplar and oak (Jankowiak et al. 2014; Milenković et al. 2018). In addition, the clade 6 species *P. lacustris* and *P. gonapodyides* were isolated from fine roots and rhizosphere of both declining and non-declining alders. Although being dominant species in waterways with proven pathogenicity to alders (Navarro et al. 2015; Trzewik et al. 2015) they are only rarely recorded from declining alders (Jung and Blaschke 2004; Černý et al. 2011; Sims et al. 2015; Trzewik et al. 2015; Bregant et al. 2020).

Continuous studies of *Phytophthora* diversity in the waterways of Europe are revealing almost ubiquitous infestations with an increasing number of *Phytophthora* species and hybrids and also other oomycetes such as *Nothophytophthora* and *Halophytophthora* spp.. Further studies are needed to clarify their actual distribution and their ecological roles. Due to the wide host ranges of some *Phytophthora* species, e.g. *P. cactorum* and *P. plurivora*, and the largely unknown host ranges of most *Phytophthora* species and hybrids surface water should not be used for irrigation of trees and crops. Extensive host range studies on major tree and crop species should be performed to unveil the potential threat posed by the *Phytophthora*, *Nothophytophthora* and *Halophytophthora* taxa found in Central European rivers and streams. This study also showed that *P. plurivora*, *P. uniformis*, *P. ×alni* and *P. ×multiformis* were the main causal agents in the aetiology of alder decline in Austria. Although alder cankers in alpine areas were exclusively caused by *P. uniformis*, *P. plurivora* proved to be the most widespread

oomycete in riparian alder forests. Due to its aggressiveness to alders *P. plurivora* might pose an increasing threat to alder stands in the future.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11557-023-01898-1>.

Acknowledgements We thank Katherina Schwanda (BFW, Vienna, Czech Republic) and Chloé Dulieu for much appreciated support.

Author contributions Conceptualization: T.C., T.L.C., T.J., H.D.; investigations: T.C., T.L.C., T.J., H.D., A.D., J.J., S.P.D.J., C.H., I.M., M.T., M.H.J.; writing original draft: T.C.; writing, review and editing: T.C., T.J. All authors have read and approved the final draft of the manuscript.

Funding Open access publishing supported by the National Technical Library in Prague. This research was supported by the Austrian Science Fund (FWF) for cofinancing the European BiodivERsA project RESIPATH: Responses of European Forests and Society to Invasive Pathogens (BIODIVERSA/0002/2012) and by the European Regional Development Fund, project 'Phytophthora Research Centre' Reg. No. CZ.02.1.01/0.0/0.0/15_003/0000453.

Data availability The DNA sequences generated during the current study are available in Genbank.

Declarations

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Adams GC, Catal M, Trummer L, Hansen EM, Reeser P, Worrall JJ (2008) *Phytophthora alni* subsp. *uniformis* found in Alaska beneath thinleaf alders. *Plant Health Prog* 9(1):38
- Aday Kaya AG, Lehtijärvi A, Şaşmaz Y, Nowakowska JA, Oszako T, Doğmuş Lehtijärvi HT, Woodward S (2018) *Phytophthora* species detected in the rhizosphere of *Alnus glutinosa* stands in the floodplain forests of Western Turkey. *For Pathol* 48(6):e12470
- Akilli S, Ulubaş Serçe Ç, Katırcıoğlu YZ, Maden S (2013) *Phytophthora* dieback on narrow leaved ash in the Black Sea region of Turkey. *For Pathol* 43(3):252–256
- Aram K, Rizzo DM (2018) Distinct trophic specializations affect how *Phytophthora ramorum* and clade 6 *Phytophthora* spp colonize and persist on *Umbellularia californica* leaves in streams. *Phytopathology* 108(7):858–869
- Balci Y, Cech TL (2004) Alder decline in Austria caused by a hybrid *Phytophthora*. In: XV US Department of Agriculture Interagency Research Forum on Gypsy Moth and Other Invasive Species Proceedings January 13th to 16th 2004, Annapolis, USA. USDA Forest Service: 3
- Belbahri L, Moralejo E, Calmin G, Oszako T, Garcia JA, Descals E, Lefort F (2006) *Phytophthora polonica*, a new species isolated from declining *Alnus glutinosa* stands in Poland. *FEMS Microbiol Lett* 261:165–174
- Belisario A, Luongo L, Vitale S, Galli M, Haegi A (2016) *Phytophthora gonapodyides* causes decline and death of English (Persian) walnut (*Juglans regia*) in Italy. *Plant Dis* 100(12):2537–2537
- Bertier L, Leus L, D'hondt L, De Cock AW, Höfte M (2013) Host adaptation and speciation through hybridization and polyploidy in *Phytophthora*. *PLoS One* 8(12):e85385
- Bily DS, Diehl SV, Cook M, Wallace LE, Sims LL, Watson C, Baird RE (2018) Temporal and locational variations of a *Phytophthora* spp community in an urban forested water drainage and stream-runoff system. *Southeast Nat* 17(1):176–201
- Bily DS, Nikolaeva E, Olson T, Kang S (2022) *Phytophthora* spp. associated with Appalachian oak forests and waterways in Pennsylvania, with *P. abietivora* as a pathogen of five native woody plant species. *Plant Dis* 106(4):1143–1156
- Bourke A (1991) Potato blight in Europe in 1845: the scientific controversy. In: Lucas JA, Shattock RD, Shaw DS, Cooke LR (eds) *Phytophthora*. Cambridge University Press, New York, pp 12–24
- Brasier CM (2000) Rise of the hybrid fungi. *Nature* 405:134–135
- Brasier CM, Kirk SA (2001) Comparative aggressiveness of standard and variant hybrid alder phytophthoras, *Phytophthora cambivora* and other *Phytophthora* species on bark of *Alnus*, *Quercus* and other woody hosts. *Plant Pathol* 50:218–229
- Brasier CM, Rose J, Gibbs JN (1995) An unusual *Phytophthora* associated with widespread alder mortality in Britain. *Plant Pathol* 44(6):999–1007
- Brasier CM, Cooke DE, Duncan JM, Hansen EM (2003) Multiple new phenotypic taxa from trees and riparian ecosystems in *Phytophthora gonapodyides*–*P. megasperma* ITS Clade 6, which tend to be high-temperature tolerant and either inbreeding or sterile. *Mycol Res* 107(3):277–290
- Brasier CM, Kirk SA, Delcan J, Cooke DEL, Jung T, Man In't Veld WA (2004) *Phytophthora alni* sp nov and its variants: designation of emerging heteroploid hybrid pathogens spreading on *Alnus* trees. *Mycol Res* 108(10):1172–1184
- Brasier CM, Scanu B, Cooke D, Jung T (2022) *Phytophthora*: An ancient, historic, biologically and structurally cohesive and evolutionarily successful generic concept in need of preservation. *IMA Fungus* 13(1):12
- Brazeo NJ, Wick RL, Hulvey JP (2016) *Phytophthora* species recovered from the Connecticut River Valley in Massachusetts, USA. *Mycologia* 108(1):6–19
- Bregant C, Sanna GP, Bottos A, Maddau L, Montecchio L, Linaldeddu BT (2020) Diversity and pathogenicity of *Phytophthora* species associated with declining alder trees in Italy and description of *Phytophthora alpina* sp. nov. *Forests* 11(8):848
- Browne GT, Ott NJ, Forbes H, Yagmour MA, Milliron LK (2020) First report of *Phytophthora chlamydospora* causing crown and root rot on almond in California. *Plant Dis* 104(7):2033–2033
- Browne GT, Hasey JK, Ott NJ, Forbes H, Arnold K, Milliron L (2021) Flooding by California rivers results in walnut scion infections by species of *Phytophthora*. *Plant Health Prog* 22(3):368–373
- Burgess TI (2015) Molecular characterization of natural hybrids formed between five related indigenous clade 6 *Phytophthora* species. *PLoS ONE* 10(8):e0134225
- Bush EA, Hong C, Stromberg EL (2003) Fluctuations of *Phytophthora* and *Pythium* spp in components of a recycling irrigation system. *Plant Dis* 87(12):1500–1506

- Caballol M, Štraus D, Macia H, Ramis X, Redondo MÁ, Oliva J (2021) *Halophytophthora fluvialis* pathogenicity and distribution along a Mediterranean-subalpine gradient. *J Fungi* 7(2):112
- Cech TL (2006) Monitoring of alder-*Phytophthora* in Vienna. In: IUFRO Working Party 7.03.10 Proceedings of the workshop on methodology of forest insect and disease survey in Central Europe Sept 11th to 14th 2006, Gmunden, Austria, p 136–141
- Cech TL, Brandstetter M (1999) *Phytophthora*-Erlensterben – Aktuelle Situation in Österreich. *Forstschutz Aktuell* 23(24):16–19
- Černý K, Strnadová V (2010) *Phytophthora* alder decline: disease symptoms, causal agent and its distribution in the Czech Republic. *Plant Prot Sci* 46(1):12–18
- Černý K, Gregorová B, Holub V, Strnadová V (2003) First records of “alder-*Phytophthora*” in the Czech Republic. *Czech Mycol* 55(3/4):291–296
- Černý K, Gregorova B, Strnadova V, Holub V, Tomsovsy M, Cervenka M (2008) *Phytophthora alni* causing decline of black and grey alders in the Czech Republic. *Plant Pathol* 57(2):370
- Černý K, Tomšovský M, Mrázková M, Strnadová V (2011) The present state of knowledge on *Phytophthora* spp. diversity in forest and ornamental woody plants in the Czech Republic. *NZ J for Sci* 41:75–82
- Chen Q, Bakhshi M, Balci Y, Broders KD, Cheewangkoon R, Chen SF, Fan XL, Gramaje D, Halleen F, Horta Jung M, Jiang N, Jung T, Májek T, Marincowitz S, Milenković I, Mostert L, Nakashima C, Nurul Faziha I, Pan M, Raza M, Scanu B, Spies CFJ, Suhaizan L, Suzuki H, Tian CM, Tomšovský M, Úrbez-Torres JR, Wang W, Wingfield BD, Wingfield MJ, Yang Q, Yang X, Zare R, Zhao P, Groenewald JZ, Cai L, Crous PW (2022) Genera of phytopathogenic fungi: GOPHY 4. *Stud Mycol* 101:417–564
- Choudhary CE, Burgos-Garay ML, Moorman GW, Hong C (2016) *Pythium* and *Phytophthora* species in two Pennsylvania greenhouse irrigation water tanks. *Plant Dis* 100(5):926–932
- Christova PK (2022) Detection of *Phytophthora gallica* in Bulgaria and co-existence with other *Phytophthora* species in a small river. *J Plant Dis Prot* 129(6):1377–1387
- Cleary M, Ghasemkhani M, Blomquist M, Witzell J (2016) First report of *Phytophthora gonapodyides* causing stem canker on European beech (*Fagus sylvatica*) in Southern Sweden. *Plant Dis* 100(10):2174
- Cooke DE, Drenth A, Duncan JM, Wagels G, Brasier CM (2000) A molecular phylogeny of *Phytophthora* and related oomycetes. *Fungal Genet Biol* 30:17–32
- Corcobado T, Cubera E, Pérez-Sierra A, Jung T, Solla A (2010) First report of *Phytophthora gonapodyides* involved in the decline of *Quercus ilex* in xeric conditions in Spain. *New Dis Rep* 22:33
- Corcobado T, Cech TL, Brandstetter M, Daxer A, Hüttler C, Kudláček T, Horta Jung M, Jung T (2020) Decline of European beech in Austria: Involvement of *Phytophthora* spp. and contributing biotic and abiotic factors. *Forests* 11(8):895
- Corcobado T, Milenković I, Saiz-Fernández I, Kudláček T, Plichta R, Májek T, Bačová A, Datková H, Dálya LB, Trifković M, Mureddu D, Račko V, Kardošová M, Đurković J, Rattunde R, Jung T (2022) Metabolomic and physiological changes in *Fagus sylvatica* seedlings infected with *Phytophthora plurivora* and the A1 and A2 mating types of *P. × cambivora*. *J Fungi* 8(3):298
- Derviş S, Türkölmez Ş, Çiftçi O, Ulubaş Serçe Ç (2016) First report of *Phytophthora chlamydospora* causing root rot on walnut (*Juglans regia*) trees in Turkey. *Plant Dis* 100(11):2336
- Derviş S, Türkölmez Ş, Çiftçi O, Özer G, Ulubaş Serçe Ç, Dikilitas M (2020) *Phytophthora littoralis*: A novel killer pathogen of plane (*Platanus orientalis*) causing canker stain and root and collar rot. *Plant Dis* 104(10):2642–2648
- Deutschmann VF (1954) Eine Wurzelsfäule an Erdbeeren, hervorgerufen durch *Phytophthora cactorum* (Leb et Cohn) Schröt. *Nachr Dtsch Pflanzenschutzd* 6:7–9
- Fleischmann F, Schneider D, Matyssek R, Obwald WF (2002) Investigations on Net CO₂ assimilation, transpiration and root growth of *Fagus sylvatica* infested with four different *Phytophthora* species. *Plant Biol* 4:144–152
- Fleischmann F, Göttlein A, Rodenkirchen H, Lütz C, Obwald W (2004) Biomass, nutrient and pigment content of beech (*Fagus sylvatica*) saplings infected with *Phytophthora citricola*, *P. cambivora*, *P. pseudosyringae* and *P. undulata*. *For Pathol* 34:79–92
- Gibbs JN, Lipscombe MA, Peace AJ (1999) The impact of *Phytophthora* disease on riparian populations of common alder (*Alnus glutinosa*) in southern Britain. *Eur J for Pathol* 29(1):39–50
- Greslebin AG, Hansen EM, Winton LM, Rajchenberg M (2005) *Phytophthora* species from declining *Austrocedrus chilensis* forests in Patagonia. *Argentina Mycologia* 97(1):218–228
- Hansen E, Delatour C (1999) *Phytophthora* species in oak forests of north-east France. *Ann for Sci* 56(7):539–547
- Hansen EM, Reeser P, Sutton W, Brasier CM (2015) Redesignation of *Phytophthora* taxon Pgchlamydo as *Phytophthora chlamydospora* sp. nov. *N Am Fungi* 10:1–14
- Hartmann G (1995) Wurzelsfäule der Schwarzerle (*Alnus glutinosa*) – eine bisher unbekannte Pilzkrankheit durch *Phytophthora cambivora*. *Forst Und Holz* 50:555–557
- Hildebrand AA (1959) A root and stalk rot of soybeans caused by *Phytophthora megasperma* Drechsler var. *sojae* var. nov. *Canad J Bot* 37(5):927–957
- Hirst JM, Stedman OJ (1960) The epidemiology of *Phytophthora infestans*: II The source of inoculum. *Ann Appl Biol* 48(3):489–517
- Hong CX, Gallegly ME, Richardson PA, Kong P, Moorman GW, Leacock JD, Ross DS (2010) *Phytophthora hydropathica*, a new pathogen identified from irrigation water, *Rhododendron catawbiense* and *Kalmia latifolia*. *Plant Pathol* 59(5):913–921
- Houston Durrant T, de Rigo D, Caudullo G (2016a) *Alnus incana* in Europe: Distribution, habitat, usage and threats. In: San-Miguel-Ayanz J, de Rigo D, Caudullo G, Houston Durrant T, Mauri A (Eds.) *European Atlas of Forest Tree Species*. Publications Office of the European Union, Luxembourg, e01ff87+: 66–67
- Houston Durrant T, de Rigo D, Caudullo G (2016b) *Alnus glutinosa* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayanz J, de Rigo D, Caudullo G, Houston Durrant T, Mauri A (Eds.) *European Atlas of Forest Tree Species*. Publications Office of the European Union, Luxembourg, e01f3c0+: 64–65
- Huai WX, Tian G, Hansen EM, Zhao WX, Goheen EM, Grünwald NJ, Cheng C (2013) Identification of *Phytophthora* species baited and isolated from forest soil and streams in northwestern Yunnan province. *China for Pathol* 43(2):87–103
- Hüberli D, Hardy GSJ, White D, Williams N, Burgess TI (2013) Fishing for *Phytophthora* from Western Australia’s waterways: a distribution and diversity survey. *Australas Plant Pathol* 42(3):251–260
- Hwang J, Oak SW, Jeffers SN (2008) Detecting *Phytophthora ramorum* and other species of *Phytophthora* in streams in natural ecosystems using baiting and filtration methods. In: *Proceedings of the Sudden Oak Death Third Symposium*, vol PSW-GTR-214. US Forest Service, Pacific Southwest Research Station, Albany, pp 55–58
- Hwang J, Oak SW, Jeffers SN (2011) Recovery of *Phytophthora* species from drainage points and tributaries within two forest stream networks: a preliminary report. *N Z J for Sci* 41:S83–S87
- Jankowiak R, Stepniewska H, Bilański P, Kolařík M (2014) Occurrence of *Phytophthora plurivora* and other *Phytophthora* species in oak forests of southern Poland and their association with site conditions and the health status of trees. *Folia Microbiol* 59(6):531–542
- Jung T (2009) Beech decline in Central Europe driven by the interaction between *Phytophthora* infections and climatic extremes. *For Pathol* 39:73–94







- Jung T, Blaschke M (2004) *Phytophthora* root and collar rot of alders in Bavaria: distribution, modes of spread and possible management strategies. *Plant Pathol* 53(2):197–208
- Jung T, Burgess TI (2009) Re-evaluation of *Phytophthora citricola* isolates from multiple woody hosts in Europe and North America reveals a new species. *Phytophthora Plurivora Sp Nov Persoonia* 22(1):95–110
- Jung T, Nechwatal J (2008) *Phytophthora gallica* sp. nov., a new species from rhizosphere soil of declining oak and reed stands in France and Germany. *Mycol Res* 112(10):1195–1205
- Jung T, Blaschke H, Neumann P (1996) Isolation, identification and pathogenicity of *Phytophthora* species from declining oak stands. *Eur J for Pathol* 26(5):253–272
- Jung T, Blaschke H, Osswald W (2000) Involvement of soilborne *Phytophthora* species in Central European oak decline and the effect of site factors on the disease. *Plant Pathol* 49:706–718
- Jung T, Nechwatal J, Cooke DE, Hartmann G, Blaschke M, Oßwald WF, Duncan JM, Delatour C (2003) *Phytophthora pseudosyringae* sp. nov., a new species causing root and collar rot of deciduous tree species in Europe. *Mycol Res* 107(7):772–789
- Jung T, Stukely MJC, Hardy GESTJ, White D, Paap T, Dunstan WA, Burgess TI (2011) Multiple new *Phytophthora* species from ITS Clade 6 associated with natural ecosystems in Australia: evolutionary and ecological implications. *Persoonia* 26:13–39
- Jung T, Vettraino AM, Cech TL, Vannini A (2013) The impact of invasive *Phytophthora* species on European forests. In: Lamour K (ed) *Phytophthora: A global perspective*. CABI, Wallingford, UK, pp 146–158
- Jung T, Orlikowski L, Henricot B, Abad-Campos P, Aday AG, Aguiñ Casal O, Bakonyi J, Cacciola SO, Cech T, Chavarriga D, Corcobado T, Cravador A, Decourcelle T, Denton G, Diamandis S, Dogmus-Lehtijärvi HT, Franceschini A, Ginetti B, Glavendekić M, Green S, Hantula J, Hartmann G, Herrero M, Ivic D, Horta Jung M, Lilja A, Keca N, Kramarets V, Lyubenova A, Machado H, di San M, Lio G, Mansilla Vázquez PJ, Marçais B, Matsiakh I, Milenkovic I, Moricca S, Nagy ZÁ, Nechwatal J, Olsson C, Oszako T, Pane A, Paplomatas EJ, Pintos Varela C, Prospero S, Rial Martínez C, Rigling D, Robin C, Rytönen A, Sánchez ME, Sanz Ros AV, Scanu B, Schlenzig A, Schumacher J, Slavov S, Solla A, Sousa E, Stenlid J, Talgø V, Tomić Z, Tsopelas P, Vannini A, Vettraino AM, Wenneker M, Woodward S, Pérez-Sierra A (2016) Widespread *Phytophthora* infestations in European nurseries put forest, semi-natural and horticultural ecosystems at high risk of *Phytophthora* diseases. *Forest Pathol* 46:134–163
- Jung T, Chang T-T, Bakonyi J, Seress D, Pérez-Sierra A, Yang X, Hong C, Scanu B, Fu C-H, Hsueh K-L, Maia C, Abad-Campos P, León M, Horta Jung M (2017a) Diversity of *Phytophthora* species in natural ecosystems of Taiwan and association with disease symptoms. *Plant Pathol* 66(2):194–211
- Jung T, Horta Jung M, Scanu B, Seress D, Kovács GM, Maia C, Pérez-Sierra A, Chang T-T, Chandelier A, Heungens K, Van Poucke K, Abad-Campos P, León M, Cacciola SO, Bakonyi J (2017b) Six new *Phytophthora* species from ITS clade 7a including two sexually functional heterothallic hybrid species detected in natural ecosystems in Taiwan. *Persoonia* 38:100–135
- Jung T, Scanu B, Bakonyi J, Seress D, Kovács GM, Durán A, Sanfuentes von Stowasser E, Schena L, Mosca S, Thu PQ, Nguyen CM, Fajardo S, González M, Pérez-Sierra A, Rees H, Cravador A, Maia C, Horta Jung M (2017c) *Nothophytophthora* gen nov, a new sister genus of *Phytophthora* from natural and semi-natural ecosystems. *Persoonia* 39(1):143–174
- Jung T, Durán A, Sanfuentes von Stowasser E, Schena L, Mosca S, Fajardo S, González M, Navarro Ortega AD, Bakonyi J, Seress D, Tomšovský M, Cravador A, Maia C, Horta Jung M (2018a) Diversity of *Phytophthora* species in Valdivian rainforests and association with severe dieback symptoms. *For Pathol* 48(5):e12443
- Jung T, Pérez-Sierra A, Durán A, Jung MH, Balci Y, Scanu B (2018b) Canker and decline diseases caused by soil- and air-borne *Phytophthora* species in forests and woodlands. *Persoonia* 40:182–220
- Jung T, La Spada F, Pane A, Aloï F, Evoli M, Horta Jung M, Scanu B, Faedda R, Rizza C, Puglisi I, di San Lio GM, Schena L, Cacciola SO (2019) Diversity and distribution of *Phytophthora* species in protected natural areas in Sicily. *Forests* 10(3):259
- Jung T, Scanu B, Brasier CM, Webber JF, Milenković I, Corcobado T, Tomšovský M, Pánek M, Bakonyi J, Maia C, Bačová A, Raco M, Rees H, Pérez-Sierra A, Horta Jung M (2020) A survey in natural forest ecosystems of Vietnam reveals high diversity of both new and described *Phytophthora* taxa including *P. ramorum*. *Forests* 11:93
- Jung T, Milenković I, Corcobado T, Májek T, Janoušek J, Kudláček T, Tomšovský M, Nagy Z, Durán A, Tarigan M, Sanfuentes von Stowasser E, Singh R, Ferreira M, Webber JF, Scanu B, Chi NM, Thu PQ, Junaid M, Rosmana A, Baharuddin B, Kusnianti T, Nasri N, Kageyama K, Hieno A, Masuya H, Uematsu S, Oliva J, Redondo M, Maia C, Matsiakh I, Kramarets V, O'Hanlon R, Tomić Ž, Brasier CM, Horta Jung M (2022) Extensive morphological and behavioural diversity among fourteen new and seven described species in *Phytophthora* Clade 10 and its evolutionary implications. *Persoonia* 49:1–57
- Jung T, Blaschke M (2006) Management strategies for the *Phytophthora* root and collar rot epidemic of alders in Bavaria. In: Progress in Research on *Phytophthora* Diseases of Forest Trees. (Brasier, CM, Jung, T & Oßwald, W, eds). *Proceedings of the 3rd International IUFRO Working Party 7.02.09 Meeting*, 11th–17th Sept. 2004, Freising, Germany. Forest Research, Farnham, Surrey, UK: 61–66
- Kanoun-Boulé M, Vasconcelos T, Gaspar J, Vieira S, Dias-Ferreira C, Husson C (2016) *Phytophthora* × *alni* and *Phytophthora lacustris* associated with common alder decline in Central Portugal. *Forest Pathol* 46:174–176
- Kliejunas JT, Ko WH (1976) Dispersal of *Phytophthora cinnamomi* on the island of Hawaii. *Phytopathology* 66(4):457–460
- Klotz LJ, Wong PP, DeWolfe TA (1959) Survey of irrigation water for the presence of *Phytophthora* spp. pathogenic to citrus. *Plant Dis Rep* 43:830–832
- Kreutzer WA, Bodine EW, Durrell LW (1940) Cucurbit diseases and rot of tomato fruit caused by *Phytophthora capsici*. *Phytopathology* 30:972–976
- Kroon LPNM, Bakker FT, van den Bosch GBM, Bonants PJM, Flier WG (2004) Phylogenetic analysis of *Phytophthora* species based on mitochondrial and nuclear DNA sequences. *Fungal Genet Biol* 41(8):766–782
- Lacey J (1967) The role of water in the spread of *Phytophthora infestans* in the potato crop. *Ann Appl Biol* 59(2):245–255
- Landa BB, Arias-Giraldo LF, Henricot B, Montes-Borrego M, Shuttleworth LA, Pérez-Sierra A (2021) Diversity of *Phytophthora* species detected in disturbed and undisturbed British soils using high-throughput sequencing targeting ITS rRNA and COI mtDNA regions. *Forests* 12(2):229
- Maia C, Horta Jung M, Carella G, Milenković I, Janoušek J, Tomšovský M, Mosca S, Schena L, Cravador A, Moricca S, Jung T (2022) Eight new *Halophytophthora* species from marine and brackish-water ecosystems in Portugal and an updated phylogeny for the genus. *Persoonia* 48(1):54–90
- Matsiakh I, Oszako T, Kramarets V, Nowakowska JA (2016) *Phytophthora* and *Pythium* species detected in rivers of the Polish-Ukrainian border areas. *Balt for* 22(2):230–238

- Matsiakh I, Kramarets V, Cleary M (2021) Occurrence and diversity of *Phytophthora* species in declining broadleaf forests in western Ukraine. For Pathol 51(1):e12662
- McIntosh DL (1964) *Phytophthora* spp. in soils of the Okanagan and Similkameen valleys of British Columbia. Canad J Bot 42(10):1411–1415
- Milenković I, Keča N, Karadžić D, Radulović Z, Nowakowska JA, Oszako T, Sikora K, Corcobado T, Jung T (2018) Isolation and pathogenicity of *Phytophthora* species from poplar plantations in Serbia. Forests 9(6):330
- Mrázková M, Černý K, Tomšovský M, Strnadová V, Gregorová B, Holub V, Pánek M, Havrdová L, Hejna M (2013) Occurrence of *Phytophthora multivora* and *Phytophthora plurivora* in the Czech Republic. Plant Prot Sci 49(4):155–164
- Murphy SK, Lee C, Valachovic Y, Bienapfl J, Mark W, Jirka A, Rizzo DM (2008) Monitoring *Phytophthora ramorum* distribution in streams within California watersheds. In: Proceedings of the Sudden Oak Death Third Symposium. US Forest Service, Pacific Southwest Research Station, Albany, CA, USA, Vol. PSW-GTR-214, 409–411 pp
- Nagel JH, Gryzenhout M, Slippers B, Wingfield MJ, Hardy GES, Stukely MJC, Burgess TI (2013) Characterization of *Phytophthora* hybrids from ITS clade 6 associated with riparian ecosystems in South Africa and Australia. Fungal Biol 117(5):329–347
- Nagel JH, Slippers B, Wingfield MJ, Gryzenhout M (2015) Multiple *Phytophthora* species associated with a single riparian ecosystem in South Africa. Mycologia 107(5):915–925
- Nagy ZÁ, Bakonyi J, Ersek T (2003) Standard and Swedish variant types of the hybrid alder *Phytophthora* attacking alder in Hungary. Pest Manag Sci 59(4):484–492
- Navarro S, Sims L, Hansen E (2015) Pathogenicity to alder of *Phytophthora* species from riparian ecosystems in western Oregon. For Pathol 45(5):358–366
- Nechwatal J, Bakonyi J, Cacciola SO, Cooke DEL, Jung T, Nagy ZÁ, Vannini A, Vettriano AM, Brasier CM (2013) The morphology, behaviour and molecular phylogeny of *Phytophthora* taxon Salix-soil and its redesignation as *Phytophthora lacustris* sp nov. Plant Pathol 62(2):355–369
- Nigrelli L, Thines M (2013) Tropical oomycetes in the German Bight – Climate warming or overlooked diversity? Fungal Ecol 6(2):152–160
- Oh E, Gryzenhout M, Wingfield BD, Wingfield MJ, Burgess TI (2013) Surveys of soil and water reveal a goldmine of *Phytophthora* diversity in South African natural ecosystems. IMA Fungus 4:123–131
- O’Hanlon R, Choiseul J, Corrigan M, Destefanis M (2016) Diversity and detections of *Phytophthora* species from trade and nontrade environments in Ireland. Bull OEPP 46:594–602
- O’Hanlon R, Choiseul J, Brennan JM, Grogan H (2018) Assessment of the eradication measures applied to *Phytophthora ramorum* in Irish *Larix kaempferi* forests. For Pathol 48(1):e12389
- O’Hanlon R, Destefanis M, Milenković I, Tomšovský M, Janoušek J, Bellgard SE, Weir BS, Kudláček T, Horta Jung M, Jung T (2021) Two new *Nothophytophthora* species from streams in Ireland and Northern Ireland: *Nothophytophthora irlandica* and *N. lirii* sp nov. Plos one 16(5):e0250527
- Oberacker P, Stepper P, Bond DM, Höhn S, Focken J, Meyer V, Schelle L, Sugrue VJ, Jeunen GJ, Moser T, Hore SR, von Meyenn F, Hipp K, Hore TA, Jurkowski TP (2019) Bio-On-Magnetic-Beads (BOMB): Open platform for high-throughput nucleic acid extraction and manipulation. PLoS Biol 17(1):e3000107
- Orlikowski LB, Ptaszek M, Rodziewicz A, Nechwatal J, Thinggaard K, Jung T (2011) *Phytophthora* root and collar rot of mature *Fraxinus excelsior* in forest stands in Poland and Denmark. For Pathol 41(6):510–519
- Orlikowski LB, Oszako T, Szkuta G (2003) First record of alder *Phytophthora* in Poland. J Plant Prot Res 43(1):33–39
- Oudemans PV (1999) *Phytophthora* species associated with cranberry root rot and surface irrigation water in New Jersey. Plant Dis 83(3):251–258
- Pérez-Sierra A, Kalantarzadeh M, Sancisi-Frey S, Brasier CM (2015) *Phytophthora siskiyouensis* causing stem lesions and cankers on *Alnus incana*. New Dis Rep 31:17
- Pintos C, Rial C, Aguin O, Ferreira V, Mansilla JP (2016) First report of *Phytophthora hydropathica* in river water associated with riparian alder in Spain. New Dis Rep 33:25
- Pisetta M, Montecchio L, Longa CMO, Salvadori C, Zottele F, Maresi G (2012) Green alder decline in the Italian Alps. For Ecol Manag 281:75–83
- Podger FD, Doepel RF, Zentmyer GA (1965) Association of *Phytophthora cinnamomi* with a disease of *Eucalyptus marginata* forest in Western Australia. Plant Dis Rep 49(11):943–957
- Redondo MA, Boberg J, Olsson CH, Oliva J (2015) Winter conditions correlate with *Phytophthora alni* subspecies distribution in Southern Sweden. Phytopathology 105(9):1191–1197
- Redondo MA, Boberg J, Stenlid J, Oliva J (2018a) Contrasting distribution patterns between aquatic and terrestrial *Phytophthora* species along a climatic gradient are linked to functional traits. ISME J 12:2967–2980
- Redondo MA, Boberg J, Stenlid J, Oliva J (2018b) Functional traits associated with the establishment of introduced *Phytophthora* spp in Swedish forests. J Appl Ecol 55(3):1538–1552
- Reeser PW, Sutton W, Hansen EM, Remigi P, Adams GC (2011) *Phytophthora* species in forest streams in Oregon and Alaska. Mycologia 103(1):22–35
- Riolo M, La Spada F, Aloï F, Giusso del Galdo G, Santilli E, Pane A, Cacciola SO (2020) *Phytophthora* diversity in a sentinel arbo-retum and in a nature reserve area. Biol Life Sci Forum 4(1):51
- Robideau GP, de Cock AWAM, Coffey MD, Voglmayr H, Brouwer H, Bala K, Chitty DW, Désaulniers N, Eggertson QA, Gachon CMM, Hu CH, Küpper FC, Rintoul TL, Sarhan E, Verstappen ECP, Zhang Y, Bonants PJM, Ristaino JB, Lévesque AC (2011) DNA barcoding of oomycetes with cytochrome c oxidase subunit I and internal transcribed spacer. Mol Ecol Resour 11(6):1002–1011
- Santilli E, Riolo M, La Spada F, Pane A, Cacciola SO (2020) First report of root rot caused by *Phytophthora bilorbang* on *Olea europaea* in Italy. Plants 9(7):826
- Santini A, Barzanti GP, Capretti P (2001) A new *Phytophthora* root disease of alder in Italy. Plant Dis 85(5):560–560
- Scanu B, Linaldeddu BT, Deidda A, Jung T (2015) Diversity of *Phytophthora* species from declining Mediterranean maquis vegetation, including two new species, *Phytophthora crassamura* and *P. ornamentata* sp. nov. PLoS One 10(12):e0143234
- Seddaiu S, Linaldeddu BT (2020) First Report of *Phytophthora acerina*, *P. plurivora*, and *P. pseudocryptogea* associated with declining common alder trees in Italy. Plant 104(6):1874
- Seddaiu S, Brandano A, Ruii PA, Sechi C, Scanu B (2020) An Overview of *Phytophthora* species inhabiting declining *Quercus suber* stands in Sardinia (Italy). Forests 11(9):971
- Shrestha SK, Zhou Y, Lamour K (2013) Oomycetes baited from streams in Tennessee 2010–2012. Mycologia 105:1516–1523
- Sims LL, Sutton W, Reeser P, Hansen EM (2015) The *Phytophthora* species assemblage and diversity in riparian alder ecosystems of western Oregon, USA. Mycologia 107(5):889–902
- Solla A, Pérez-Sierra A, Corcobado T, Haque MM, Diez JJ, Jung T (2010) *Phytophthora alni* on *Alnus glutinosa* reported for the first time in Spain. Plant Pathol 59(4):798–798
- Stamler RA, Sanogo S, Goldberg NP, Randall JJ (2016) *Phytophthora* species in rivers and streams of the southwestern United States. Appl Environ Microbiol 82(15):4696–4704

- Štěpánková P, Černý K, Strnadova V, Hanáček P, Tomšovský M (2013) Identification of *Phytophthora alni* subspecies in riparian stands in the Czech Republic. *Plant Prot Sci* 49:3–10
- Streito JC, Legrand PH, Tabary F, Jarnouen de Villartay GJD (2002) Phytophthora disease of alder (*Alnus glutinosa*) in France: Investigations between 1995 and 1999. *For Pathol* 32(3):179–191
- Sutton W, Hansen EM, Reeser PW, Kanaskie A (2009) Stream monitoring for detection of *Phytophthora ramorum* in Oregon tanoak forests. *Plant Dis* 93(11):1182–1186
- Thomson SV, Allen RM (1974) Occurrence of *Phytophthora* species and other potential plant pathogens in recycled irrigation water. *Plant Dis Rep* 58(10):945–949
- Trzewik A, Orlikowski LB, Oszako T, Nowakowska JA, Orlikowska T (2015) The characterization of *Phytophthora* isolates obtained from diseased *Alnus glutinosa* in Poland. *Balt for* 21(1):44–50
- Tsao PH (1990) Why many *Phytophthora* root rots and crown rots of tree and horticultural crops remain undetected 1. *Bull OEPP* 20(1):11–17
- Tucker CM, Milbrath JA (1942) Root rot of *Chamaecyparis* caused by a species of *Phytophthora*. *Mycologia* 34(1):94–103
- Van Poucke K, Haegeman A, Goedefroit T, Focquet F, Leus L, Jung MH, Nave C, Redondo MA, Husson C, Kostov K, Lyubenova A, Christova P, Chandelier A, Slavov S, De Cock A, Bonants P, Werres S, Oliva Palau J, Marçais B, Jung T, Stenlid J, Ruttink T, Heungens K (2021) Unravelling hybridization in *Phytophthora* using phylogenomics and genome size estimation. *IMA Fungus* 12(1):1–24
- Von Broembsen S (1984) Distribution of *Phytophthora cinnamomi* in rivers of the south-western Cape Province. *Phytophylactica* 16(3):227–230
- White TJ, Bruns T, Lee SJWT, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky J, White TJ (eds) *PCR Protocols: a Guide to Methods and Applications*. Academic Press, San Diego, pp 315–322
- Xu X, Huai W, Hamiti ZX, Zhao W (2019) *Phytophthora* species from Xinjiang wild apple forests in China. *Forests* 10:927
- Yang X, Hong C (2014) *Halophytophthora fluviatilis* sp. nov. from freshwater in Virginia. *FEMS Microbiol Lett* 352:230–237
- Yang X, Richardson PA, Hong C (2014) *Phytophthora* × *stagnum* nothosp nov, a new hybrid from irrigation reservoirs at ornamental plant nurseries in Virginia. *PLoS ONE* 9(7):e103450
- Zamora-Ballesteros C, Haque MMU, Diez JJ, Martín-García J (2017) Pathogenicity of *Phytophthora alni* complex and *P. plurivora* in *Alnus glutinosa* seedlings. *For Pathol* 47(2):e12299

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Tamara Corcobado^{1,2}  · Thomas L. Cech² · Andreas Daxer² · Henrieta Ďatková¹  · Josef Janoušek¹ · Sneha Patra^{1,3}  · Daniella Jahn² · Christine Hüttler² · Ivan Milenković^{1,4}  · Michal Tomšovský¹  · Marília Horta Jung^{1,5} 

✉ Thomas Jung
thomas.jung@mendelu.cz; dr.t.jung@gmail.com

Tamara Corcobado
tamara.corcobado@mendelu.cz;
tamara.corcobado@bfw.gv.at

Thomas L. Cech
thomas.cech@bfw.gv.at

Andreas Daxer
andreas.daxer@bfw.gv.at

Henrieta Ďatková
xdatkova@mendelu.cz

Josef Janoušek
janousek.jose@gmail.com

Sneha Patra
snehampatra@gmail.com

Daniella Jahn
daniella.jahn13@gmail.com

Christine Hüttler
christine.huettler@bfw.gv.at

Ivan Milenković
ivan.milenkovic@mendelu.cz

Michal Tomšovský
michal.tomsovsky@mendelu.cz

Marília Horta Jung
marilia.jung@mendelu.cz

- 1 Phytophthora Research Centre, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemědělská 3, 61300 Brno, Czech Republic
- 2 Natural Hazards and Landscape, Unit of Phytopathology, Department of Forest Protection, Federal Research and Training Centre for Forests, Seckendorff-Gudent-Weg 8, 1131 Vienna, Austria
- 3 Global Change Research Institute of the Czech Academy of Sciences, Belidla 986/4a, 603 00 Brno, Czech Republic
- 4 Faculty of Forestry, University of Belgrade, Kneza Višeslava 1, 11030 Belgrade, Serbia
- 5 Phytophthora Research and Consultancy, Am Rain 9, 83131 Nußdorf, Germany