Correction to: Toxicity and clinical outcomes of single-fraction high-dose-rate brachytherapy combined with external beam radiotherapy for high-/very high-risk prostate cancer: A dosimetric analysis of toxicity

Takayuki Sakurai¹ · Shigeyuki Takamatsu¹ · Satoshi Shibata¹ · Koji Iwata¹ · Masashi Taka¹ · Tomoyasu Kumano² · Tomoyuki Makino³ · Atsushi Mizokami³ · Toshifumi Gabata¹

Published online: 26 January 2021
© Japan Radiological Society 2021

Correction to:
Japanese Journal of Radiology (2020) 38:1197–1208
<https://doi.org/10.1007/s11604-020-01023-2>

In the original publication, Toshifumi Gabata should be listed as the final author. The revised version of author order is given in this correction.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s11604-020-01023-2>.

✉ Takayuki Sakurai
tkyk81@gmail.com

¹ Department of Radiology, Graduate School of Medicine, Kanazawa University, Kanazawa, Ishikawa 920-8641, Japan

² Department of Radiology, Graduate School of Medicine, Gifu University, Gifu 501-1194, Japan

³ Department of Integrative Cancer Therapy and Urology, Graduate School of Medicine, Kanazawa University, Kanazawa, Ishikawa 920-8641, Japan