ERRATUM

Erratum to: Provider patient-sharing networks and multiple-provider prescribing of benzodiazepines

Mei-Sing Ong, PhD^{1,2}, Karen L. Olson, PhD^{1,3}, Aurel Cami, PhD^{1,3}, Chunfu Liu, PhD⁵, Fang Tian, PhD⁵, Nandini Selvam, PhD MPH⁵, and Kenneth D. Mandl, MD, MPH^{1,3,4}

¹Children's Hospital Informatics Program, Boston Children's Hospital, Boston, MA, USA; ²Australian Institute of Health Innovation, Macquarie University, Sydney, NSW, Australia; ³Department of Pediatrics, Harvard Medical School, Boston, MA, USA; ⁴HealthCore, Inc., Alexandria, VA, USA; ⁵Center for Biomedical Informatics, Harvard Medical School, Boston, MA, USA.

J Gen Intern Med 31(5):588

DOI: 10.1007/s11606-015-3533-x

© Society of General Internal Medicine 2015

ERRATUM TO: J GEN INTERN MED

DOI: 10.1007/s11606-015-3470-8

The odds ratio for the relationship between median care density and multiple prescribing of benzodiazepine was incorrectly reported in the text as 0.76 (95% CI 0.60 – 0.95; $p=0.028$).

It should have read, “Backward stepwise logistic regression revealed that median care density was negatively associated with the risk of an overlapping benzodiazepine prescription (OR 0.76; 95% CI 0.60 – 0.95; $p=0.0178$) after adjusting for covariates (Table 3).” In the Discussion, it should have also read the following: “Consistent with these studies, in univariate analysis, we showed a significant relationship between the number of prescribers and overlapping benzodiazepine prescriptions; however, the magnitude of the association was reduced when considered alongside care density in multivariate analysis.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11606-015-3470-8>.

Published online October 30, 2015