CAPSULE COMMENTARIES

Capsule Commentary on Balaban et al., Impact of a Patient Navigator Program on Hospital-Based and Outpatient Utilization Over 180 Days in a Safety-Net Health System

Michael Hochman, MD, MPH

The Gehr Family Center for Implementation Science, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA.

J Gen Intern Med 32(9):1025

DOI: 10.1007/s11606-017-4102-2

© Society of General Internal Medicine 2017

Complex case management offers an intuitive strategy for improving care for high-risk populations, particularly during hospital-to-home care transitions.¹ Unfortunately, however, complex case management has shown only modest benefits in rigorous evaluations.¹

Nevertheless, research has begun to shed light on which approaches to complex case management offer the most promise. Specifically, the most effective programs seem to involve close, interpersonal interaction among care coordinators, patients, and primary care clinicians; a focus on care transitions; a component of medication management; and the targeting of patients amenable to intervention, such as the frail elderly.²

The study by Balaban et al.³ in this week's *JGIM* adds important insights to this burgeoning literature.

In their study, Balaban et al. report 180-day outcomes of a randomized evaluation of a care transition intervention provided by unlicensed community health workers (referred to as "patient navigators"). These navigators identified hospitalized patients at risk for readmission, visited them prior to discharge, provided post-discharge phone calls, and offered support with medications, appointments, transportation, communication, and self-care. Among patients 60 years of age and older, there was a modest but significant reduction of 0.21 hospital-based encounters per patient, which was accompanied by an increase in ambulatory utilization in the first 30 days following discharge (presumably ambulatory visits substituted for avoidable re-hospitalizations). Paradoxically, however, among patients younger than 60, there was an increase of 0.79 hospital-based encounters per patient. The authors acknowledge that the explanation for this finding among patients <60—which was also observed in a previously published short-term analysis⁴—is not clear.

What can we learn from these results? First, this study supports the notion that unlicensed staff can provide meaningful support for high-risk patients, at least in certain settings. Other studies have also demonstrated this finding.⁵

Second, the contradictory results among patients ≥ 60 vs. those <60 underscore the differential impact of complex case management in different populations. Increasingly it seems that these programs are most effective among frail, older adults.²

Finally, it is clear that we are only beginning to understand the nuances of complex case management. This is a ripe area for future research!

Corresponding Author: Michael Hochman, MD, MPH; The Gehr Family Center for Implementation Science, Keck School of Medicine University of Southern California, Los Angeles, CA, USA (e-mail: mhochman@med.usc.edu).

Compliance with Ethical Standards:

Conflict of Interest: The author has no conflicts of interest with this article.

REFERENCES

1. Hochman M, Asch SM. Disruptive models in primary care: caring for high-needs, high-cost populations. *J Gen Intern Med*. 2017;32(4):392–397.
2. Brown RS, Peikes D, Peterson G, Schore J, Razafindrakoto CM. Six features of medicare coordinated care demonstration programs that cut hospital admissions of high-risk patients. *Health Aff (Millwood)* 2012; 31(6):1156–66.
3. Balaban RB, Zhang F, Vialle-Valentin CE, Galbraith AA, Burns ME, Larochelle MR, Ross-Degnan D. Impact of a patient navigator program on hospital-based and outpatient utilization over 180 days in a safety-net health system. *J Gen Intern Med*. doi:10.1007/s11606-017-4074-2
4. Balaban RB, Galbraith AA, Burns ME, Vialle-Valentin CE, Larochelle MR, Ross-Degnan D. A patient navigator intervention to reduce hospital readmissions among high-risk safety-net patients: a randomized controlled trial. *J Gen Intern Med*. 2015;30(7):907–15.
5. Kangovi S, Mitra N, Grande D, et al. Patient-centered community health worker intervention to improve posthospital outcomes: a randomized clinical trial. *JAMA Intern Med*. 2014;174(4):535–43.