

## Comparison of enzymically glucuronidated flavonoids with flavonoid aglycones in an in vitro cellular model of oxidative stress protection

David E. Stevenson · Janine M. Cooney ·  
Dwayne J. Jensen · Reginald Wibisono · Aselle Adaim ·  
Margot A. Skinner · Jingli Zhang

Published online: 2 July 2008  
© The Society for In Vitro Biology 2008

**Erratum to: In Vitro Cell.Dev.Biol.—Animal**  
**DOI 10.1007/s11626-007-9072-y**

In the last line of the caption for Figure 3, the last word “determinations)” was inadvertently omitted.

The last line should read: H<sub>2</sub>O<sub>2</sub> under the same conditions was 52.2±5.4 (mean of ten separate determinations).

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11626-007-9072-y>

---

D. E. Stevenson (✉) · J. M. Cooney · D. J. Jensen  
The Horticulture and Food Research Institute of New Zealand,  
Private Bag 3123, Waikato Mail Centre,  
Hamilton 3240, New Zealand  
e-mail: [dstevenson@hortresearch.co.nz](mailto:dstevenson@hortresearch.co.nz)

R. Wibisono · A. Adaim · M. A. Skinner · J. Zhang  
The Horticulture and Food Research Institute of New Zealand,  
Private Bag 92–169, Mt Albert,  
Auckland, New Zealand