

Primary culture of insect midgut cells

Raziel S. Hakim · Silvia Caccia · Marcia Loeb ·
Guy Smaghe

Published online: 23 May 2009
© The Society for In Vitro Biology 2009

Erratum to: In Vitro Cell.Dev.Biol.—Animal
DOI 10.1007/s11626-009-9176-7

On page 2, right-hand column, insert the following paragraph before the heading **Procedure**.

Culture medium.

200 ml Grace's medium, 14 ml FBS, 0.2 g Yeastolate, 0.1 to 0.5% gentamicin (Sigma), 1.2 ml vitamin mixture, 0.02% antibiotic-antimycotic solution 1X, 1 ng / μ l 20-OH-E, and 10 μ l/ml FBX or alternatively 100 ng/ml arylphorin.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11626-009-9176-7>.

R. S. Hakim (✉)
Department of Anatomy, College of Medicine,
Howard University,
520 W. St NW,
Washington, DC 20059, USA
e-mail: rhakim@mac.com

S. Caccia · G. Smaghe
Laboratory of Agrozoology, Department of Crop Protection,
Faculty of Bioscience Engineering, Ghent University,
9000 Ghent, Belgium

M. Loeb
Insect Biocontrol Laboratory, USDA-ARS,
Beltsville, MD 20705–2350, USA

Present Address:
S. Caccia
Departament de Genètica, Facultat de Ciències Biològiques,
Campus de Burjassot, Universitat de València,
46100 Valencia, Spain