



Effects of different healing agents on acclimatization success of *in vitro* rooted Garnem (*Prunus dulcis* × *Prunus persica*) rootstock

Heydem Ekinci¹ · Necla Saskin² · Bekir Erol Ak¹ · Birgul Dikmetas Dogan²

Received: 23 October 2023 / Accepted: 8 February 2024 / Editor: Marco Buenrostro-Nava
© The Author(s) 2024

Abstract

Continuing productivity of the acclimatization stage of plantlets means successful micropropagation. Due to the free water and high humidity in the culture container environment, poorly developed cuticle results in rapid water loss and drying of plantlets with watery stems and leaves, incomplete stomata, and large intercellular space. When plantlets are exposed to an environmental condition different from the culture medium, problems, such as rapid water loss and drying, may be encountered, and the survival rate of plantlets *in vitro* may be reduced. The aim of this study was to reduce the stress factors that occur during the acclimatization phase. For this reason, various healing agents have been used. Some of these compounds are ascorbic acid (AsA), salicylic acid (SA), and sodium nitroprusside (SNP). In the study, the response of AsA (100.0 and 200.0 mg L⁻¹), SA (100.0 and 200.0 mg L⁻¹), and SNP (100.0 and 200.0 μM) applications on growth parameters (survival rate (%), node count (pieces per plant), count of leaves (pieces per plant), shoot length (mm), and shoot diameter (mm)) and physiological variable (Soil Plant Analysis Development (SPAD)) were examined. The AsA100 (100.0 mg L⁻¹ ascorbic acid), AsA200 (200.0 mg L⁻¹ ascorbic acid), and SNP100 (100.0 μM sodium nitroprusside) applications resulted in an improvement in node count, leaf count per plant, shoot length, and shoot diameter parameters. The highest shoot length (60.50 ± 7.81 mm) and node count (16.83 ± 1.15 pieces per plantlet) were achieved with the AsA100 application. The maximum stem diameter (2.31 ± 0.37 mm) was determined with the SNP100 application. There were no statistically significant differences found in the survival rate, leaf count, and SPAD parameter. The current study determined that AsA, SA, and SNP applications were effective in regulating several growth parameters in Garnem plantlets and in reducing acclimation stress thereby facilitating adaptation to external conditions.

Keywords Acclimatization · *In vitro* · Healing agent · Stress

Introduction

The use of rootstocks holds great importance in fruit cultivation. The use of rootstocks is imperative to counteract adverse soil conditions. Rootstocks have various effects on the variety. Utilizing rootstocks cannot only enhance fruit yield and quality but also promote the growth and development of the variety.

Rootstocks are divided into two categories based on their method of acquisition: clonal and seedling rootstocks. Rootstocks obtained from seeds demonstrate a heterozygous structure. Hence, clonal rootstocks are preferred based on this reasoning. Clonal rootstocks can be propagated through different methods, such as cutting and biotechnology. Clonal rootstocks carry the characteristics of the parent plant. Therefore, micropropagation is preferred in cultivating to obtain a uniform plant structure

✉ Heydem Ekinci
heydemekinci@harran.edu.tr

Necla Saskin
neclasaskinn@gmail.com

Bekir Erol Ak
beak@harran.edu.tr

Birgul Dikmetas Dogan
dikmetasbirgul@gmail.com

¹ Faculty of Agriculture, Department of Horticulture, Harran University, 63050 Sanliurfa, Turkey

² Graduate School of Natural and Applied Sciences, Department of Horticulture, Harran University, 63050 Sanliurfa, Turkey

(Nimbolkar *et al.* 2016; Sharma *et al.* 2020; Ak *et al.* 2021). Garnem rootstock is a widely used clonal rootstock for the *Prunus* species. This rootstock was obtained through the crossbreeding of the Spanish almond variety, Garfi, as the female parent, and the North American peach, Nemared, as the pollen donor. Garnem rootstock demonstrates good compatibility in grafting with plum, peach, almond, and nectarine. It is tolerant to drought, and weak, calcareous, and heavy soils. This rootstock can be propagated through both cutting and tissue culture methods (Felipe 2009; Erbas 2011; İpek 2015; Kankaya *et al.* 2021).

Micropropagation is a technique developed for rapid multiplication and holds significant commercial potential in obtaining disease-free plants of high-quality standards (George and Debergh 2007; Chandra *et al.* 2010; Saskin *et al.* 2022). However, one of the disadvantages of micropropagation is its costliness, the requirement for intensive labor, and the low survival rate upon transfer of the *in vitro* propagated plant to external conditions. The most important factor limiting the success of the micropropagation method is the acclimatization process of the obtained plants. During the acclimatization phase, high losses occur due to different factors, such as light intensity, temperature, and water stress (Kumar and Rao 2012). The primary reasons for this are water loss after the transfer of plants, planting shock, various pathogenic attacks, weak photosynthesis, and similar factors occurring during the post-transfer period (Krishna *et al.* 2005; Kara *et al.* 2022). The plant is adversely affected by such stressful situations.

Various healing agents have been used against stress factors in the acclimatization phase. One of these healing agents, sodium nitroprusside, which is a nitric oxide donor, is widely used in various plant tissue culture methods because it positively affects plant growth. SNP also has positive effects against abiotic and biotic stresses. When used *in vitro*, it increases shoot and root formation. Additionally, it yields positive results in callus formation, organogenesis, transformation, and embryogenesis, and it also helps against browning (Sundararajan *et al.* 2022).

Salicylic acid is found in all plant groups. It belongs to the group of plant growth regulators. It reverses the effects of abscisic acid, which is effective in leaf shedding. The main roles of salicylic acid include influencing plant growth, ion uptake, flowering, important enzyme activities, photosynthesis rate, and thermogenesis (Raskin *et al.* 1990; Popova *et al.* 1997; Yusuf *et al.* 2013; Rady *et al.* 2015).

Salicylic acid promotes seed germination and root development. Salicylic acid is directly effective against the negativities caused by various abiotic stresses (drought, salinity, and heavy metal). Roots can change the root structure by reacting to these stress factors in various ways. With the application of SA, the effects that occur according to the

degree of stress are largely eliminated (Bagautdinova *et al.* 2022).

Ascorbic acid is an organic acid with antioxidant properties. It is one of the most important antioxidant molecules. It directly affects plant growth and development positively. It provides cell division and expansion (Zhang *et al.* 2016). AsA plays an important role in plant physiological processes because it is a cofactor for various enzymes. Ascorbic acid (AsA) provides protection to the plant against photosynthesis and environmental stresses (Shah *et al.* 2019; Smirnoff and Wheeler 2000; Çoban and Aras 2022).

Research has also reported that plant losses occur during the acclimatization process in plants propagated *in vitro* (Pospóšilová *et al.* 1999; Kara *et al.* 2022). The high water content of plantlets propagated *in vitro* and the fact that stomata are not fully developed make the adaptation process difficult. The same situation occurs in Garnem plantlets. While some studies focused on the regeneration of Garnem rootstock, the resulting plantlets were transferred to external conditions. However, the acclimatization of Garnem rootstock has not been specifically examined, and growth-supporting exogenous applications have not been tried (Erfani *et al.* 2017; Ak *et al.* 2021). In the present study, different healing agents and concentrations were tried to improve the process. With the use of these healing agents, the aim is to prevent the losses that can occur during the transfer of plantlets to the external environment, to increase the success of survival in the acclimatization stage, and to obtain healthier plants.

The principle of success in plant tissue culture is to enhance the survival rate of *in vitro* propagated plantlets. Based on this notion, this study investigated the effects of foliar spray application of 100.0 mg L⁻¹ and 200.0 mg L⁻¹ doses of SA, 100.0 mg L⁻¹ and 200.0 mg L⁻¹ AsA, and 100.0 and 200.0 µM SNP on the acclimatization stage of *in vitro* propagated Garnem plantlets.

Material–methods

Plant material and treatments In the research, shoots obtained from the Garnem rootstock collection plot belonging to the Department of Horticulture, Faculty of Agriculture, Harran University, Sanliurfa, Türkiye, were utilized. Shoots taken during the active growth phase of the Garnem rootstock were propagated using the node culture method in the Plant Tissue Culture Laboratory of the Department of Horticulture, Faculty of Agriculture, Harran University. Garnem rootstock used for the *Prunus* species was one of the clonal rootstocks commonly used for plum, peach, and almond species. This rootstock was strong and tolerant to drought and calcareous heavy-textured soils (İpek 2015; Kankaya *et al.* 2021). Three replications were constructed

with five plants in each replication, based on random plots. Shoots were prepared as single nodular cuttings. For surface sterilization, they were kept in a sterile cabinet (vertical air flow) (Nuve, Ankara, Türkiye) for 1 min in 70% ethyl alcohol and then in 12 min 10.0% sodium hypochlorite solution (Domestos, Unilever, Türkiye, and 5.0% active ingredient); and at the end of these processes, the cuttings were rinsed three times with sterile distilled water (Yalçın-Mendi *et al.* 2003; İpek 2015). After surface sterilization, the cuttings were treated with 3.0% sucrose (Merck, Darmstadt, Germany), 0.4 mg L⁻¹ GA₃ (Merck, Hohenbrunn, Germany), and 2.0 mg L⁻¹ BAP (Acros Organics, Morris Plains, New Jersey), and 1.0 mg L⁻¹ Plant Preservative Mixture (PPM) (Plant Cell Technology, Washington, DC) to prevent bacterial contamination of explants, and they were transferred to jars containing Murashige and Skoog (MS) (Sigma-Aldrich®, St. Louis, MO) (Murashige and Skoog 1962) medium solidified with 0.6% agar (Duchefa, 2003 RV Haarlem, Netherlands). After emergence of the nodes, shoots were transferred to MS rooting medium containing 1.5 mg L⁻¹ IBA (Merck) (Kara and Yazar 2020; Kara *et al.* 2022). The explants were kept in the climatic chamber with a temperature of 18 to 28 °C and white fluorescent lamps (4000 lx) (Koninklijke Philips Electronics N.V., Netherlands) as the light source with 16 h of light and 8 h of darkness.

Healthy shoots (approximately 4 to 5 cm in length) with more than three nodes from the multiplied shoot clumps were used for root regeneration. Garnem plantlets, after completing root formation in the rooting medium (1.5 mg L⁻¹ IBA), were removed from the nutrient medium, and their roots were cleaned. The plantlets were transferred to cups (produced by a local manufacturer) with a volume of

450 mL and a diameter of 95 mm, containing aquarium sand (produced by a local manufacturer). These transferred plantlets were placed in a climatic chamber with conditions set at 65% relative humidity and a temperature of 24 ± 1 °C (white fluorescent tubes (produced by a local manufacturer), 100 μmol m⁻² s⁻¹ irradiance) after the transfer. In the study, AsA (100.0 mg L⁻¹ and 200.0 mg L⁻¹), SA (100.0 mg L⁻¹ and 200.0 mg L⁻¹), and SNP (100.0 μM and 200.0 μM) chemicals and concentrations were used (Zangani *et al.* 2023). AsA (Sigma-Aldrich®), SA (Merck), and SNP (Merck) chemicals were dissolved in distilled water until they reached a homogeneous structure. Subsequently, they were transferred into spray bottles (Fig. 1). The prepared solutions were applied as a spray to the plantlets undergoing acclimatization at 5-d intervals for a total period of 50 d. The control group of plants received a spray of distilled water.

Survival rate, node count, shoot length, count of leaves per plant, shoot diameter, and SPAD (Soil Plant Analysis Development) measurements The survival rate (%), node count (node per plant), count of leaves per plant (leaves per plant), shoot length (mm), stem diameter (mm), and SPAD parameters were examined. The survival rate was determined as the percentage ratio of the number of live plants to the total number of plantlets taken for acclimatization. Node count and leaf count were determined by counting the nodes and leaves present on the plantlets. Shoot diameter and shoot length were measured in millimeters using calipers (Kara *et al.* 2022). Chlorophyll content of the 3rd and 4th leaves of all shoots was measured by a chlorophyllmeter (SPAD-502, Minolta, Tokyo, Japan) (Bayraktar *et al.* 2016; Fang *et al.* 2018).



Figure 1. The appearance of Garnem (*Prunus dulcis* × *Prunus persica*) plantlets propagated under *in vitro* conditions during the acclimatization process. (A) General view of plantlets obtained from

Garnem (*Prunus dulcis* × *Prunus persica*) rootstock in *in vitro* conditions. (B) Keeping the plantlets transferred to aquarium sand under controlled conditions. (C) Applications are made by spraying.

Data Analysis Data were subjected to statistical analysis using the analysis variance (ANOVA) technique (Gomez and Gomez 1984). To determine significant differences among the treatments, Tukey multiple comparison test (with a significance level of 0.05) was used. JMP Pro 13 software was used for Tukey multiple comparison test. Hierarchical clustering analysis (HCA) and principal component analysis (PCA) were conducted *via* the Software R (Version 4.1.1, R Foundation for Statistical Computing, Vienna, Austria).

Results and discussion

Garnem rootstock plantlets some growth parameters and physiological variable The growth parameters (survival rate (%), node count (nodes per plant), leaf count (leaves per plant), shoot length (mm), shoot diameter (mm)) and the physiological variable SPAD were significantly affected by the applications (Table 1). Plantlets subjected to AsA100 application showed the highest node count and shoot length parameters. On the contrary, plantlets subjected to individual control and SNP200 applications exhibited the lowest levels in growth parameters (survival rate, node count, leaf count, and shoot diameter). Generally, healing agent applications to plantlets exposed to external stress in the acclimatization stage improved the levels of growth parameters, and the AsA100, AsA200, and SNP 100 healing agent individual treatments showed the best performance compared to SNP200 and SA200 applications.

Survival rate (%) The micropropagation method provided clonally efficient and fast production. One of the factors affecting the success of this method was the survival rate in the acclimatization stage.

In plants obtained under *in vitro* conditions, the leaf water content is high due to elevated humidity in the container. In addition, the cuticle layer and stoma pores are not fully

developed. Therefore, in the process of transferring to external conditions, plantlets lose water through transpiration and the survival rate of plantlets decreases (Shiwani *et al.* 2022). Based on this idea, beneficial applications aimed at improving plant quality during the acclimatization phase may contribute to the enhancement of the process and shorten the adaptation period. One of the critical processes during acclimatization was the survival rate. There was no statistically significant difference in the survival rates of plants in this study. The highest survival rate was seen in the AsA200 application with 53.33%, and the lowest survival rate was observed in the AsA 100 (40.00%) and SNP200 application with 40.00% (Table 1). In the study conducted by Vasar (2001), the effect of ascorbic acid and citric acid on the acclimatization of *Prunus avium* L. plantlets was examined, and the highest survival rate was recorded in the plantlets sprayed with ascorbic acid. Similar studies also observed positive effects of various beneficial agents applied during the acclimatization phase on the survival rates of plantlets obtained under *in vitro* conditions (Johkan *et al.* 2008; Kara *et al.* 2022).

Node count (nodes per plant) There was a statistically significant difference in the number of nodes in the plants. While the highest number of 16.83 nodes per plant was obtained from the AsA100 application, the lowest number, 8.83 nodes per plant, was determined in the AsA200 treatment (Table 1). AsA played an important role in plant stress tolerance. Many studies revealed that ascorbic acid played an important role in improving plant tolerance to abiotic stress (Shalata and Neumann 2001; Toksoy and Dogru 2021). The acclimatization phase of plantlets grown under *in vitro* conditions is a very important step in which plantlets are exposed to a high mortality rate. Ascorbic acid is an antioxidant that can inhibit the oxidation process that occurs in the plant under adverse conditions. In addition to its antioxidant role, ascorbic acid also plays a role in cell

Table 1. Effect of healing agents applied to Garnem (*Prunus dulcis* × *Prunus persica*) plantlets during the acclimatization stage on vegetative properties

| Treatments | Survival rate (%) | Nodes per plant | Leaves per plant | Shoot length (mm) | Stem diameter (mm) | SPAD* |
|------------|-------------------|-----------------|------------------|-------------------|--------------------|---------------|
| Control | 40.00 ± 0.00 | 9.33 ± 0.29b | 8.33 ± 1.26 | 39.72 ± 6.62abc | 1.68 ± 0.10ab | 35.90 ± 5.88 |
| SNP100 | 46.67 ± 11.55 | 12.83 ± 2.36ab | 11.67 ± 2.52 | 20.33 ± 6.77c | 2.31 ± 0.37a | 28.42 ± 5.56 |
| SNP200 | 40.00 ± 0.00 | 10.00 ± 2.29 b | 6.00 ± 0.50 | 32.37 ± 2.41bc | 1.34 ± 0.20b | 31.75 ± 2.93 |
| AsA100 | 40.00 ± 0.00 | 16.83 ± 1.15a | 10.17 ± 2.08 | 60.50 ± 7.81a | 1.80 ± 0.21ab | 21.78 ± 1.35 |
| AsA200 | 53.33 ± 11.55 | 8.83 ± 2.08b | 8.50 ± 4.77 | 37.44 ± 6.79abc | 1.92 ± 0.19ab | 18.47 ± 6.25 |
| SA100 | 46.67 ± 11.55 | 11.50 ± 1.00b | 12.17 ± 1.61 | 52.22 ± 14.07ab | 1.98 ± 0.21a | 23.40 ± 10.04 |
| SA200 | 46.67 ± 11.55 | 11.83 ± 1.89ab | 11.00 ± 2.29 | 42.08 ± 12.12abc | 1.86 ± 0.11ab | 32.29 ± 10.78 |

SNP100 sodium nitroprusside 100.0 µM, SNP200 sodium nitroprusside 200.0 µM, AsA100 ascorbic acid 100.0 mg L⁻¹, AsA200 ascorbic acid 200.0 mg L⁻¹, SA100 salicylic acid 100.0 mg L⁻¹, SA200 salicylic acid 200.0 mg L⁻¹

*SPAD, Soil Plant Analysis Development

division, elongation, and shoot regeneration (Babbar *et al.* 2010; Das and Srivastav 2015). In the present study, it was determined that ascorbic acid played an important role in increasing the number of nodes. In the studies carried out on vegetable species, it was determined that the seedling growth, the number of branches, and the number of nodes were positively affected (Sajid and Aftab 2009; Gaafar *et al.* 2020; Çoban and Aras 2022).

Number of leaves per plant (leaves per plant) Plants treated with SA100 showed higher leaf count (12.17 leaves per plant) than control plants (Table 1). These results supported the findings of Rady *et al.* (2015) who reported an increase in the number of leaves of *Phaseolus vulgaris* L. plants treated with SA. There was no statistically substantial difference between the applications in terms of the number of leaves. Salicylic acid affects many biochemical and physiological responses in plants, acts as an important signaling molecule, and has various effects on tolerance to environmental stresses (Nazar *et al.* 2011; Aras 2018). When the effects of SA were examined, positive improvements were observed in different studies. In a study conducted by Yenilmez (2016), there was a significant difference in the number of leaves of SA applied at different doses to some vine rootstocks in saline conditions, and improvements were noted.

Shoot length (mm) In *in vitro* conditions, humidity in culture vessels was high, and lighting was low compared to the conventional situation. The decrease in plant growth rate under stress during the adaptation process of these plantlets to external conditions was probably due to the high water and moisture content in leaves and other tissues. For this reason, when transferred from *in vitro* conditions to *ex vitro* conditions, the plant cannot immediately adapt to external conditions and losses increase (Munns 2002; Ejaz *et al.* 2012). As demonstrated by the growth parameters examined in this study, exogenous application of AsA significantly increased the growth of Garnem plants under stress conditions that occurred during adaptation. Plants treated with AsA100 showed higher shoot length (60.50 mm) than control plants. A statistically significant difference was found between the treatments in terms of shoot lengths (Table 1). It is known that AsA plays a direct role in shoot development of woody plants (Bilska *et al.* 2019). In Hassan *et al.* (2021), it was reported that AsA that was applied exogenously from the leaves in barley plants grown in saline conditions increased plant growth. The results supported this study (Gaafar *et al.* 2020).

Shoot diameter (mm) The stem diameter was found to be statistically significant among the treatments. The shoot diameter was highest in the SNP100 treatment with 2.31 mm

compared to the control (Table 1). Exogenous SNP application in plants is one of the strategies to improve plant photosynthetic performance and stress tolerance. It can be concluded that the increased endogenous nitric oxide with the application of SNP alleviates the negative effects on growth and photosynthesis under stress conditions (Li *et al.* 2022). Studies on appropriate nitric oxide concentration have also reported that it contributes to DNA concentration. Therefore, nitric oxide promotes cell division and growth (Bai *et al.* 2012; Novikova *et al.* 2017). It is thought that increasing stem diameter is directly proportional to the role of nitric oxide in cell division and growth (Tan *et al.* 2013).

SPAD As a result of the applications, there was no statistical difference between the SPAD values. The highest value was obtained from the control (35.90) application. However, among the conducted applications, it was determined that the SA200 (32.29) application yielded results closest to the control. This result can be explained by the salicylic acid application increasing the chlorophyll content in the plant (Gautam and Singh 2009).

Hierarchical clustering analysis (HCA) HCA is a clustering method that determines order in groups of samples and between groups taking into account the order of importance. The result of the HCA is usually presented in a dendrogram, which is a graph showing the arrangement and relationships of the samples in tree form. HCA was performed using six different parameters in line with the applications made in the study (Fig. 2). In the heatmap resulting from the analysis, applications were divided into groups as I, II, III, IV, and V. The applications were situated as follows: control and SNP200 in group I, AsA100 in group II, SA100 and SA200 in group III, SNP100 in group IV, and AsA200 in group V (Fig. 2). HCA of the six physiological parameters considered as a result of the applications was divided into four groups: A, B, C, and D. The grouping included the following: in cluster B, shoot length and node count; in cluster D, leaf count and shoot diameter; survival rate in cluster C; and the physiological variable SPAD in cluster A. During the adaptation of the plantlets obtained under *in vitro* conditions to the external environment, parameters, such as node count, shoot length, survival rate, count of leaves, and shoot diameter, were adversely affected by the hardening stress. An improvement in these values was observed with the effect of AsA100, SA100, AsA200, and SNP100. While node count and shoot length parameters were in the B cluster, as a result of the AsA100 application, it had the highest value. Finally, the survival rate parameter was placed in cluster C, and the highest value was detected in the AsA200. It has been reported by studies that exogenous ascorbic acid applications have a positive effect on cell growth and division, differentiation, and metabolism in plants as an antioxidant




Figure 2. Heat map for evaluated morphological traits under control and treatment conditions in the Garnem (*Prunus dulcis* × *Prunus persica*) rootstock. I, II, III, IV, and V represent the groups of the treatments studied. A, B, C, and D represent the groups of param-

eters studied. SNP100 (sodium nitroprusside 100.0 μM), SNP200 (sodium nitroprusside 200.0 μM), AsA100 (ascorbic acid 100.0 mg L^{-1}), AsA200 (ascorbic acid 200.0 mg L^{-1}), SA100 (salicylic acid 100.0 mg L^{-1}), and SA200 (salicylic acid 200.0 mg L^{-1}).

(Athar *et al.* 2009; Xu *et al.* 2015; Hafez and Gharib 2016). In the present study, the positive increase in node count and shoot length parameters in the AsA100 application and the survival rate parameters in the AsA200 application were parallel to similar studies (Sajid and Aftab 2009; Xu *et al.* 2015; Seleem *et al.* 2021). SNP induces carbohydrate metabolism in plants through consumption of soluble sugars by increasing photosynthesis and carbohydrate anabolism (Ghadakchiasl *et al.* 2017; Chavoushi *et al.* 2019). This increased metabolic activity may be a possible reason for the observed improvement in shoot diameter from growth parameters (Fig. 2).

Principal component analysis (PCA) PCA was used to represent the current variation of the dataset obtained from the applications made in the study (Fig. 3). Variables observed in the iteratively calculated PCA were largely explained as PC1 (71.7%) and PC2 (27.3). When the analysis map was examined, survival rate, shoot diameter, and parameters for SNP100 and AsA200 applications were in the same group, and their values that were greater than 1 made the applications meaningful. It has been observed that the parameters shoot length, node count, and count of leaves were in the same direction as the SA100 application. However, it was determined that the SPAD parameter, unlike the survival rate, number of nodes, number of leaves, shoot length, and

shoot diameter, was in the opposite group with the control and SNP200 applications. When looking at the PCA map of the applications, SNP100 and AsA100 applications had a significant effect on the parameters; SNP200 applications, which were closer to the control, were negatively affected (Fig. 3). It was determined that AsA100, SNP100, and AsA200 applications had a positive effect on the growth parameters measured in Garnem plantlets (Fig. 3). Using molecules that have been shown to have protective functions against stresses may help reduce the negative effects of plants adapted to external environments under *in vitro* conditions. Such healing substances are powerful antioxidants that protect cells from damage caused by abiotic stresses. These antioxidants (AsA, SA, and SNP) play a key role in plant growth, development, and defense responses (Smirnoff 1996; Hemavathi Upadhyaya *et al.* 2010; Kumar *et al.* 2011).

Conclusions

The micropropagation method plays an important role in obtaining a large number of plants as well as clonal propagation. The tissue culture method plays a role in the development of new varieties and the preservation of plant gen resources. Plant losses during acclimatization of valuable plants obtained by this method adversely affect the sustainability of




Figure 3. A loading plot of all the detected variables included in PCA (principal component analysis) for the physiological parameters in Garnem (*Prunus dulcis* × *Prunus persica*) rootstock. SNP100 (sodium nitroprusside 100.0 μM), SNP200 (sodium nitroprusside

200.0 μM), AsA100 (ascorbic acid 100.0 mg L^{-1}), AsA200 (ascorbic acid 200.0 mg L^{-1}), SA100 (salicylic acid 100.0 mg L^{-1}), and SA200 (salicylic acid 200.0 mg L^{-1}).

micropropagation. Therefore, minimizing plant loss during acclimatization is the most important step in making plant tissue culture sustainable. Due to low ambient humidity and high lighting, plants cannot adapt to external conditions right away when transferred from *in vitro* to *ex vitro*. Research supports the positive effects of healing agents on stress reduction. In the present study, it was determined as a result of observations that exogenous applications improved the vegetative development of the plant (shoot length, number of leaves, and stem diameter) positively and accelerated the adaptation process to external conditions. The count of nodes (16.83 pieces per plant) and shoot length (60.50 mm) increased positively with AsA (100 mg L^{-1}) while the stem diameter (2.31 mm) showed a positive increase with SNP (100 μM) applications in vegetative development. For subsequent acclimatization studies, AsA100 and SNP100 applications are recommended. Improving the acclimatization process of the Garnem rootstock is crucial, as it serves as a significant source of plant material for almond cultivation, which is widely used in the provinces of Sanliurfa and Adiyaman. Additionally, the Garnem rootstock is used in research conducted at numerous universities throughout Türkiye.

Author contribution Investigation and conceptualization, B.E.A., N.S., H.E. (Heydem Ekinci); methodology, N.S., H.E. (Heydem Ekinci); software and formal analysis, N.S., H.E. (Heydem Ekinci); writing-original,

writing-review and editing, and draft preparation, H.E., B.E.A., B.D.D and N.S.; visualization H.E. (Heydem Ekinci). All authors have read and agreed to the published version of the manuscript.

Funding Open access funding provided by the Scientific and Technological Research Council of Türkiye (TÜBİTAK).

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethical approval This original article does not include any studies with human participants or animals by any of the authors.

Consent to participate Informed consent was obtained from all individual participants included in the study.

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will

need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ak BE, Kıyar PK, Hatipoglu IH, Dikmetas B (2021) Effects of different BA and IBA concentrations on proliferation and rooting of ‘GARNEM’ rootstock *in vitro* propagation. *Intl J Agr Environ Food Sci* 5:470–476
- Aras S (2018) Plant growth regulator use in temperate zone fruit trees against abiotic stresses. *2nd Int Symp Innov Approaches Sci Stud* 3:1305–1307 (Samsun, Turkey)
- Athar H-u-R, Khan A, Ashraf M (2009) Inducing salt tolerance in wheat by exogenously applied ascorbic acid through different modes. *J Plant Nut* 32:1799–1817
- Babbar N, Oberoi HS, Uppal DS, Patil RT (2010) Total phenolic content and antioxidant capacity of extracts obtained from six important fruit residues. *Food Res Intl* 44:391–396
- Bagautdinova ZZ, Omelyanchuk N, Tyapkin AV, Kovrizhnykh VV, Lavrekha VV, Zemlyanskaya EV (2022) Salicylic acid in root growth and development. *Intl J Mol Sci* 23:2228
- Bai S, Li M, Yao T, Wang H, Zhang Y, Xiao L, He Y (2012) Nitric oxide restrain root growth by DNA damage induced cell cycle arrest in *Arabidopsis thaliana*. *Nitric Oxide* 26:54–60
- Bayraktar M, Naziri E, Akgun IH, Karabey F, Ilhan E, Akyol B, Gurel A (2016) Elicitor induced stevioside production, *in vitro* shoot growth and biomass accumulation in micropropagated *Stevia Rebaudiana*. *Plant Cell Tiss Org Cult* 127:289–300
- Bilska K, Wojciechowska N, Alipour S, Kalemba EM (2019) Ascorbic acid—the little-known antioxidant in woody plants. *Antioxidants* 8:645
- Chandra S, Bandopadhyay R, Kumar V, Chandra R (2010) Acclimatization of tissue cultured plantlets: from laboratory to land. *Biotechnol Lett* 32:1199–1205
- Chavoushi M, Najafi F, Salimi A, Angaji S (2019) Improvement in drought stress tolerance of safflower during vegetative growth by exogenous application of salicylic acid and sodium nitroprusside. *Indust Crop Prod* 134:168–176
- Çoban GA, Aras S (2022) Effects of ascorbic and oxalic acids on cucumber seedling growth and quality under mildly limey soil conditions. *Gesunde Pflanzen* 75:1925–1932
- Das P, Srivastav AK (2015) To study the effect of activated charcoal, ascorbic acid and light duration on *in-vitro* micropropagation of *Aloe vera* L. *Intl J Innovat Res Sci Engineer Technol* 4:3131–3138
- Ejaz B, Sajid ZA, Aftab F (2012) Effect of exogenous application of ascorbic acid on antioxidant enzyme activities, proline contents, and growth parameters of *Saccharum* spp. Hybrid cv. HSF-240 under Salt Stress. *Turkish J Biol* 36:630–640
- Erbas B (2011) Myrobalan ve Garnem anaçlarının *in vitro* doku kültürü teknikleri ile çoğaltımı üzerinde araştırmalar. Yüksek Lisans, Gaziantep Üniversitesi Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, p 67
- Erfani M, Miri SM, Imani A (2017) *In vitro* shoot proliferation and rooting of Garnem rootstock as influenced by basal media, plant growth regulators and carbon sources. *Plant Cell Biotechnol Mol Biol* 18:101–109
- Fang X, Li Y, Nie J, Wang C, Huang K, Zhang Y, Yi Z (2018) Effects of nitrogen fertilizer and planting density on the leaf photosynthetic characteristics, agronomic traits and grain yield in Common Buckwheat (*Fagopyrum esculentum* M.). *Field Crops Res* 219:160–168
- Felipe AJ (2009) ‘Felinem’, ‘Garnem’, and ‘Monegro’ almond×peach hybrid rootstocks. *Hort Sci* 44:196–197
- Gaafar AA, Ali SI, El-Shawadfy MA, Salama ZA, Şekara A, Ulrichs C, Abdelhamid MT (2020) Ascorbic acid induces the increase of secondary metabolites, antioxidant activity, growth and productivity of the common bean under water stress conditions. *Plants* 9:627
- Gautam S, Singh PK (2009) Salicylic acid-induced salinity tolerance in corn grown under NaCl stress. *Acta Physiol Plant* 31:1185–1190
- George EF, Debergh PC (2007) Plant propagation by tissue culture, 3rd edn. In: George EF, Hall MA, De Klerk GJ (eds) *Micropropagation uses and methods*, vol 1. Springer Science & Business Media, Dordrecht, pp 29–35
- Ghadakchiasl A, Mozafari A-a, Ghaderi N (2017) Mitigation by sodium nitroprusside of the effects of salinity on the morpho-physiological and biochemical characteristics of *Rubus idaeus* under *in vitro* conditions. *Physiol Mol Biol Plants* 23:73–83
- Gomez KA, Gomez AA (1984) *Statistical procedures for agricultural research*, 2nd edn. John Wiley & Sons, New York, 680 pp
- Hafez E, Gharib H (2016) Effect of exogenous application of ascorbic acid on physiological and biochemical characteristics of wheat under water stress. *Intl J Plant Prod* 10:579–596
- Hassan A, Amjad SF, Saleem MH, Yasmin H, Imran M, Riaz M, Ali Q, Joyia FA, Ahmed S, Ali S (2021) Foliar application of ascorbic acid enhances salinity stress tolerance in barley (*Hordeum vulgare* L.) through modulation of morpho-physio-biochemical attributes, ions uptake, osmo-protectants and stress response genes expression. *Saudi J Biol Sci* 28:4276–4290
- Hemavathi Upadhyaya CP, Akula N, Young KE, Chun SC, Kim DH, Park SW (2010) Enhanced ascorbic acid accumulation in transgenic potato confers tolerance to various abiotic stresses. *Biotechnol Lett* 32:321–330
- İpek M (2015) *In vitro* şartlarda Garnem ve Myrobalan 29C anaçlarının kurak stresine karşı tepkilerinin belirlenmesi. Doktora Tezi, Selçuk Üniversitesi Fen Bilimleri Enstitüsü, Bahçe Bitkileri Anabilim Dalı, Konya, p 142
- Johkan M, Oda M, Mori G (2008) Ascorbic acid promotes graft-take in sweet pepper plants (*Capsicum annum* L.). *Sci Hort* 116:343–347
- Kankaya A, Polat M, Eskimez İ, Mertoğlu K (2021) Şeftali fidan üretiminde aşı başarısı bakımından anaç çapı ve kalem dinlenmesinin etkileri: ‘Artemis–Garnem’ örneği. *ISUBÜ J Agric Fac* 16:150–153
- Kara Z, Yazar K (2020) Bazı üzüm çeşitlerinde *in vitro* poliploidi uyarımı. *Anadolu J Agr Sci* 35:410–418
- Kara Z, Yazar K, Ekinci H, Dogan O, Ozer A (2022) The effects of ortho silicone applications on the acclimatization process of grapevine rootstocks. *Selcuk J Agric Food Sci* 36:233–237
- Krishna H, Singh S, Sharma R, Khawale R, Grover M, Patel V (2005) Biochemical changes in micropropagated grape (*Vitis vinifera* L.) plantlets due to arbuscular-mycorrhizal fungi (AMF) inoculation during *ex vitro* acclimatization. *Sci Hort* 106:554–567
- Kumar K, Rao I (2012) Morphophysiological problems in acclimatization of micropropagated plants *in-ex vitro* conditions—a reviews. *J Ornament Hort Plants* 2:271–283
- Kumar S, Kaur R, Kaur N, Bhandhari K, Kaushal N, Gupta K, Bains T, Nayyar H (2011) Heat-stress induced inhibition in growth and chlorosis in mungbean (*Phaseolus aureus* Roxb.) is partly mitigated by ascorbic acid application and is related to reduction in oxidative stress. *Acta Physiol Plant* 33:2091–2101
- Li X, Wang S, Chen X, Cong Y, Cui J, Shi Q, Liu H, Diao M (2022) The positive effects of exogenous sodium nitroprusside on the plant growth, photosystem II efficiency and Calvin cycle of tomato seedlings under salt stress. *Sci Hort* 299:111016
- Munns R (2002) Comparative physiology of salt and water stress. *Plant Cell Environ* 25:239–250

- Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol Plant* 15:473–497
- Nazar R, Iqbal N, Syyed S, Khan NA (2011) Salicylic acid alleviates decreases in photosynthesis under salt stress by enhancing nitrogen and sulfur assimilation and antioxidant metabolism differentially in two mungbean cultivars. *J Plant Physiol* 168:807–815
- Nimbolkar P, Awachare C, Reddy Y, Chander S, Hussain F (2016) Role of rootstocks in fruit production—a review. *J Agric Eng Food Technol* 3:183–188
- Novikova GV, Mur LA, Nosov AV, Fomenkov AA, Mironov KS, Mamaeva AS, Hall MA (2017) Nitric oxide has a concentration-dependent effect on the cell cycle acting via EIN2 in *Arabidopsis thaliana* cultured cells. *Front Physiol* 8:142
- Popova L, Pancheva T, Uzunova A (1997) Salicylic acid: properties, biosynthesis and physiological role. *Bulg J Plant Physiol* 23(1–2):85–93
- Pospóšilová J, Tichá I, Kadleček P, Haisel D, Plzánková Š (1999) Acclimatization of micropropagated plants to *ex vitro* conditions. *Biol Plant* 42:481–497
- Rady MM, Mohamed GF, Abdalla AM, Ahmed YH (2015) Integrated application of salicylic acid and *Moringa oleifera* leaf extract alleviates the salt-induced adverse effects in common bean plants. *J Agric Technol* 11:1595–1614
- Raskin I (1992) Role of salicylic acid in plants. *Annu Rev Plant Biol* 43:439–463
- Raskin I, Skubatz H, Tang W, Meeuse BJ (1990) Salicylic acid levels in thermogenic and non-thermogenic plants. *Ann Bot* 66:369–373
- Sajid ZA, Aftab F (2009) Amelioration of salinity tolerance in *Solanum tuberosum* L. by exogenous application of ascorbic acid. *In Vitro Cell Dev Biol - Plant* 45:540–549
- Saskin N, Ak BE, Ekinçi H (2022) The usage of node culture *in vitro* conditions. In: Kirca L, Bak T, Guler E, Dogru-Cokran B, Kilic D (eds) *Proceeding Book. 5th Intl Agric Cong, Denizli, Turkey, 5–6 Dec 2022*, online, pp 90–99
- Seleem EA, Ibrahim HMS, Taha ZK (2021) Exogenous application of ascorbic acid and putrescine: a natural eco-friendly potential for alleviating NaCl stress in barley (*Hordeum vulgare*). *Emir J Food Agric* 33:657–670
- Shah T, Latif S, Khan H, Munsif F, Nie L (2019) Ascorbic acid priming enhances seed germination and seedling growth of winter wheat under low temperature due to late sowing in Pakistan. *Agron* 9:757
- Shalata A, Neumann PM (2001) Exogenous ascorbic acid (vitamin C) increases resistance to salt stress and reduces lipid peroxidation. *J Exp Bot* 52:2207–2211
- Sharma JB, Chauhan N, Rana K, Bakshi M (2020) Evaluation of rootstocks for temperate fruit crops—a review. *Int J Curr Microbiol Appl Sci* 9:3533–3539
- Shiwani K, Sharma D, Kumar A (2022) Improvement of plant survival and expediting acclimatization process. In: Gupta S, Chaturvedi P (eds) *Commer scale tissue cult hortic plant crops*. Springer, Singapore, pp 277–291
- Smirnoff N (1996) Botanical briefing: the function and metabolism of ascorbic acid in plants. *Ann Bot* 78:661–669
- Smirnoff N, Wheeler GL (2000) Ascorbic acid in plants: biosynthesis and function. *Crit Rev Plant Sci* 19:267–290
- Sundararajan S, Rajendran V, Sivakumar HP, Kumariah M, Ramalingam S (2022) Growth modulation by nitric oxide donor sodium nitroprusside in *in vitro* plant tissue cultures—a review. *Biol* 77:1699–1711
- Tan BC, Chin CF, Alderson P (2013) Effects of sodium nitroprusside on shoot multiplication and regeneration of *Vanilla planifolia* Andrews. *In Vitro Cell Dev Biol - Plant* 49:626–630
- Toksoy S, Dogru A (2021) Tuz stresi altundaki hiyar bitkilerinde ekzojen askorbik asit uygulamalarının fotosistem II aktivitesi üzerindeki etkileri. *KSU J Agric Nat* 24:757–765
- Vasar V (2001) Effect of ascorbic acid and citric acid on *ex vitro* rooting and acclimatization of *Prunus avium* L. microshoots. In: *Int Symp Acclimatization Establishment Micropropagated Plants, Sani-Halkidiki, Macedonia, Greece*, pp 251–254
- Xu Y, Xu Q, Huang B (2015) Ascorbic acid mitigation of water stress-inhibition of root growth in association with oxidative defense in tall fescue (*Festuca arundinacea* Schreb.). *Front Plant Sci* 6:807
- Yalçın-Mendi N, Ipek M, Kacan H, Curuk S, Sari N, Çetiner S, Gaba VA (2003) histological analysis of regeneration in watermelon. *J Plant Biochem Biotechnol* 12:147–150
- Yenilmez N (2016) Farklı salisilik asit dozlarının bazı Amerikan asma anaçlarının tuzluluğa olan dayanımı üzerine etkileri. *Ordu Üniversitesi Fen Bilimleri Enstitüsü, Bahçe Bitkileri Anabilim Dalı, Ordu*, p 83
- Yusuf M, Hayat S, Alyemeni MN, Fariduddin Q, Ahmad A (2013) Salicylic acid: physiological roles in plants, 2nd edn. In: Hayat S, Ahmad A, Alyemeni M (eds) *Salicylic acid: plant, growth and development*. Springer, Dordrecht, pp 15–30
- Zangani E, Angourani HR, Andalibi B, Rad SV, Mastinu A (2023) Sodium nitroprusside improves the growth and behavior of the stomata of *Silybum marianum* L. subjected to different degrees of drought. *Life* 13:875
- Zhang X, Yu HJ, Zhang XM, Yang XY, Zhao WC, Li Q, Jiang WJ (2016) Effect of nitrogen deficiency on ascorbic acid biosynthesis and recycling pathway in cucumber seedlings. *Plant Physiol Biochem* 108:222–230