Correction to: Osteoporosis medication use among Australian women over two decades

Amanda L Stuart¹ • Julie A Pasco^{1,2,3} • Mohammadreza Mohebbi⁴ • Mark A Kotowicz^{1,2,3} • Kara L Holloway-Kew¹ • Sarah M Hosking¹ • Lana J Williams¹

Published online: 1 July 2020

© International Osteoporosis Foundation and National Osteoporosis Foundation 2020

Correction to: Archives of Osteoporosis.

<https://doi.org/10.1007/s11657-019-0661-7>

The original version of this article, published 06 May 2020, unfortunately contained a mistake. The presentation of Figs. 2 (a, b and c) were incorrect. The corrected figures are given below.

The online version of the original article can be found at <https://doi.org/10.1007/s11657-019-0661-7>

✉ Amanda L Stuart
amandh@barwonhealth.org.au

¹ IMPACT, the Institute for Mental and Physical Health and Clinical Translation, School of Medicine, Deakin University, Geelong, Australia

² Melbourne Medical School-Western Campus, The University of, Melbourne, St Albans, Australia

³ University Hospital Geelong, Geelong, Australia

⁴ Biostatistics Unit, Faculty of Health, Deakin University, Geelong, Australia

a) Whole group

b) Women with osteoporosis by BMD criteria

c) Women at high risk of fracture according to the FRAX calculator

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.