Correction to: A 3i hip fracture liaison service with nurse and physician co-management is cost-effective when implemented as a standard clinical program

L. A. Beaupre¹ · D. Lier² · C. Smith³ · L. Evens³ · H. M. Hanson^{4,5} · A. G. Juby⁶ · P. Kivi⁷ · S. R. Majumdar² ·
On behalf of the STOP-Fracture Team

Published online: 27 August 2020

© International Osteoporosis Foundation and National Osteoporosis Foundation 2020

Correction to: Archives of Osteoporosis (2020) 15:113
<https://doi.org/10.1007/s11657-020-00781-w>

The original version of this article, published on 22 July 2020, unfortunately contained a mistake.

Throughout the article, the authors state “anti-absorptive” medication when it should be instead “anti-resorptive” medication.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s11657-020-00781-w>

✉ L. A. Beaupre
lauren.beaupre@ualberta.ca

¹ University of Alberta (Physical Therapy), 2-50 Corbett Hall, Edmonton, AB T6G2G4, Canada

² University of Alberta (Medicine), Edmonton, AB, Canada

³ Alberta Bone and Joint Health Institute, Calgary, AB, Canada

⁴ Alberta Health Services, Calgary, AB, Canada

⁵ University of Calgary, Calgary, AB, Canada

⁶ University of Alberta (Geriatric Medicine), Edmonton, AB, Canada

⁷ University of Alberta (Family Medicine), Edmonton, AB, Canada