Joint position statement on management of patient with osteoporosis during COVID-19 contingency from the AMMOM, CONAMEGER, FELAEN, FEMECOG, FEMECOT, and ICAAFYD

Francisco Torres-Naranjo^{1,2,3} · Pilar De la Peña-Rodríguez^{2,4} · Roberto Enrique López-Cervantes^{2,5,6} · Jorge Morales-Torres^{2,7} · Jorge Morales-Vargas^{2,7} · Hugo Gutiérrez-Hermosillo^{2,8,9} · Alan Christopher Guzmán-Rico¹ · Roberto Gabriel González-Mendoza¹⁰ · Pedro Nel Rueda Plata^{11,12} · Miguel Flores Castro^{9,13} · Cuauhtémoc Celis Gonzalez³ · Rolando Espinosa Morales^{2,14} · Sergio Quintero Hernández^{6,15} · Juan Ricardo López-Taylor¹⁰

Received: 12 August 2020 / Accepted: 25 November 2020 / Published online: 25 January 2021
© International Osteoporosis Foundation and National Osteoporosis Foundation 2021

Abstract

Summary Infection by SARS-Cov-2 (COVID-19) has affected practically all the world. This joint position statement of Latin American Medical Societies provides an updated guide for the prevention, diagnosis, and treatment of osteoporotic patients in the face of possible clinical scenarios posed by the COVID-19 health crisis.

Background Infection by SARS-Cov-2 (COVID-19) has affected practically all the world. Characterized by high contagiousness, significative morbidity, and mortality in a segment of those infected, it has overwhelmed health services and forced to redirect resources to the emergency while impacting the attention of acute non-COVID-19 and many chronic conditions.

Objective The objective of this study is to provide an updated guide for the prevention, diagnosis, and treatment of osteoporotic patients in the face of possible clinical scenarios posed by the COVID-19 health crisis.

Methods A task force, of bone specialists with a wide range of disciplines in the field of osteoporosis and fragility fracture, was convened with the representation of several professional associations, namely, the Mexican Association of Bone and Mineral Metabolism (AMMOM), the National College of Geriatric Medicine (CONAMEGER), the Latin American Federation of Endocrinology (FELAEN), the Mexican Federation of Colleges of Obstetrics and Gynecology (FEMECOG), the Mexican Federation of Colleges of Orthopedics and Traumatology (FEMECOT), and the Institute of Applied Sciences for Physical Activity and Sports of the University of Guadalajara (ICAAFYD). Clinical evidence was collated, and an evidence report was rapidly generated and disseminated. After finding the gaps in the available evidence, a consensus opinion of experts was made. The resulting draft was reviewed and modified accordingly, in 4 rounds, by the participants.

✉ Francisco Torres-Naranjo
dr.francisco.torres@icloud.com

¹ Centro de Investigación Ósea y de la Composición Corporal, Guadalajara, Mexico

² Mexican Association of Bone and Mineral Metabolism (AMMOM), Mexico City, Mexico

³ Mexican Federation of Colleges of Obstetrics and Gynecology (FEMECOG), Mexico City, Mexico

⁴ Servicios Médicos De la Peña, Guadalajara, Mexico

⁵ Clínica de Fracturas por Osteoporosis, Hospital de Especialidades San Francisco de Asís, Guadalajara, Mexico

⁶ Mexican Federation of Colleges of Orthopedics and Traumatology (FEMECOT), Guadalajara, Mexico

⁷ Morales Vargas Centro de Investigación SC, Leon, Mexico

⁸ Hospital Aranda de la Parra, Leon, Mexico

⁹ National College of Geriatric Medicine (CONAMEGER), Mexico City, Mexico

¹⁰ Institute of Applied Sciences for Physical Activity and Sport, Department of Human Movement Sciences, Education, Sport, Recreation and Dance, University Health Sciences Center, University of Guadalajara, Guadalajara, Jalisco, Mexico

¹¹ Instituto Endocrinológico de Colombia, Bogota, Colombia

¹² Latin American Federation of Endocrinology (FELAEN), Bogota, Colombia

¹³ Departamento de Medicina Interna, Hospital Civil de Guadalajara, Guadalajara, Mexico

¹⁴ Departamento de Reumatología, Instituto Nacional de Rehabilitación, Mexico City, Mexico

¹⁵ University of Guadalajara, Guadalajara, Jalisco, Mexico

Results The task force approved the initial guidance statements, with moderate and high consensus. These were combined, resulting in the final guidance statements on the (1) evaluation of fracture risk; (2) stratification of risk priorities; (3) indications of bone density scans and lab tests; (4) initiation and continuation of pharmacologic therapy; (5) interruptions of therapy; (6) treatment of patients with incident fracture; (7) physical therapy and fall prevention; and (8) nutritional interventions.

Conclusion These guidance statements are provided to promote optimal care to patients at risk for osteoporosis and fracture, during the current COVID-19 pandemic. However, given the low level of available evidence and the rapidly evolving literature, this guidance is presented as a “living document” and future updates are anticipated.

Keywords Osteoporosis · COVID-19 · SARS-CoV-2 · Fragility fracture · Statement

Introductory notes

A new disease named COVID-19 caused by a novel coronavirus called SARS-CoV-2 has affected millions of persons worldwide.

SARS-CoV-2

In December 2019, a possible coronavirus outbreak was reported in Wuhan, China. The causative virus was initially identified as 2019-nCoV and then officially named SARS-CoV-2 [1]. The disease quickly spread from Asia to Europe, America, and virtually the entire world. The World Health Organization (WHO) formally declared the pandemic COVID-19 by March 11, 2020. And Latin America was declared as the new epicenter of the coronavirus pandemic by June 2020, when most new cases and deaths were in the Americas, and several Latin American countries were among the higher ranks of cumulative cases and deaths by the infection [2].

SARS-CoV-2 is easily transmitted; the estimated basic reproduction number (R_0) ranges from 2.24 to 6.47 [3–5]. COVID-19 clinical features range from asymptomatic or very mild disease to severe cases with elevated mortality. This group of patients requires attention in hospitals, bringing health systems to their limits. Extraordinary measures have been taken to mitigate its transmission, through the implementation of social distancing for long periods and redirecting health resources to focus on the attention of COVID-19 cases.

Healthcare systems have been overwhelmed and have issued general recommendations urging people to avoid hospitals and doctor’s offices unless strictly necessary. These measures represent a challenge to provide continuity of healthcare for other significant public health problems such as chronic degenerative diseases, including osteoporosis and fragility fractures. Some pragmatic adaptations have been proposed for continued delivery of medical attention [6].

Osteoporosis

Osteoporosis is a chronic disease characterized by loss of bone mass and structural deterioration, lessens bone strength, and increases fracture risk. Osteoporosis is a frequent cause of

disability and death [7–9]. Given the chronic nature of osteoporosis, patients require regular evaluation and long-term treatment, especially those who have suffered fragility fractures. Those patients are at a very high risk of suffering one or more new fractures in the following 24 months [10–12], increasing the risk of disability and death [13]. The situation of Latin America and the Caribbean, with a multi-ethnic population above 669 million persons, increasing elderly population; high incidence and prevalence of osteoporosis and fractures, and limited access to diagnostic tools and therapy, has led this condition to be an important personal, social, and economic burden. Guidelines for diagnosis and treatment of osteoporosis exist in many countries, but in many of them, health authorities’ endorsement and adherence are limited. FRAX calculator is available for seven countries, and adaptations to use in neighboring countries are common [14, 15].

It is estimated that due to social distancing measures across the countries, many patients will stop treatment and delay diagnostic studies. At the same time, hospital reconversion, confinement, and postponement of non-urgent medical consultations may put the continuity of medical care at risk, including possible interruptions in the treatment of patients with osteoporosis. Stopping administration of some treatment modalities [16] and discontinuation of secondary fracture prevention services predispose to an increased risk of fracture. Since the epidemic curves of COVID-19 can be “propagated curves” lasting longer than 6 months, mitigation strategies could be prolonged and particularly affect adults 60 years of age and older, in whom the prevalence of osteoporosis and the incidence of fragility fracture are high.

It is essential to establish measures to continue with patients’ care at imminent fracture risk, avoiding further saturation at the hospital emergency services. An analysis of the impact of different epidemiological scenarios on medical care for osteoporosis and fragility fractures should be considered and, consequently, issue recommendations for adjustments in care processes. Those recommendations should address patients continuing their treatment without exposing them to unnecessary risks of contracting COVID-19. It is also necessary to recognize that the burden of COVID-19 in a country can vary significantly between regions, mainly in large

territory countries with marked differences in the social determinants of health among members of the population.

Objective

The objective of this study is to provide an updated guide for the prevention, diagnosis, and treatment of osteoporotic patients in the face of possible clinical scenarios posed by the COVID-19 health crisis.

Methodology

The Mexican Association of Bone and Mineral Metabolism (AMMOM), the National College of Geriatric Medicine (CONAMEGER), the Latin American Federation of Endocrinology (FELAEN), the Mexican Federation of Colleges of Obstetrics and Gynecology (FEMECOG), the Mexican Federation of Colleges of Orthopedics and Traumatology (FEMECOT), and the Institute of Applied Sciences for Physical Activity and Sports of the University of Guadalajara (ICAAFYD) jointed an expert panel of bone specialists with a wide range of disciplines in the field of osteoporosis and fragility fracture (endocrinologists, gynecologists, geriatricians, orthopedic surgeons, rheumatologists, and exercise and nutrition scientists), to focus in establishing an organized response position statement during the COVID-19 contingency. The members of this panel were invited based on their knowledge and expertise in diagnosing and treating osteoporosis and their ability to critically evaluate and analyze the available published information for application in decision-making in a clinical setting.

Before their inclusion in the panel of experts, the summoned members declared their possible conflicts of interest. The majority were free of conflict of interest during the last 2 years. The expert panel members did not receive financial compensation or were informed about the sources of financing before or during the document's analysis or preparation.

The panel of experts evaluated the factors that could negatively affect osteoporotic patients' care during the COVID-19 pandemic, based on the best evidence available at the time of this analysis. Possible adjustments were proposed for standard practice as well as its potential effects on the patient's general health, considering scientific support, relevance, and the feasibility of its application in general clinical practice.

Clinical evidence was collated, and an evidence report was rapidly generated and disseminated. After finding the gaps in the available evidence, a consensus opinion of experts was made. After the first draft was made, an anonymous peer review was made in four rounds. However, given the low level of available evidence and the rapidly evolving literature, the

expert's panel will continue with periodical reviews to update these guidelines.

The guidelines for osteoporosis management in the COVID-19 health crisis of the American Society for Bone and Mineral Research (ASBMR), American Association of Clinical Endocrinologists (AACE), Endocrine Society, European Calcified Tissue Society (ECTS), and the National Osteoporosis Foundation (NOF) [17], as well as the recommendations of the Capture the Fracture initiative of the International Osteoporosis Foundation (CTF-IOF) [18] and the International Society of Clinical Densitometry (ISCD) [19], were included in the reasoning.

Recommendations

Whenever possible, existing current clinical practice guidelines should be observed during COVID-19 contingency. Otherwise, we recommend the following.

Stratify the need for care

During the COVID-19 pandemic, stratify the need for care based on the severity of the clinical situation, the impact of a delay of health interventions on the prognosis and evolution, and the absolute risk of suffering a major fragility fracture in the next 12 months.

Based on these factors, guide the most appropriate behaviors and identify the need for immediate attention and the possibility of postponing some measures or procedure.

We recommend stratifying the need for care at three priority levels:

- Priority A. Patients have a condition that puts the patient's life at imminent risk, is clinically unstable, or where a short delay would significantly alter their prognosis. Those patients require prioritizing immediate care and not deferring attention.
- Priority B. These are clinically stable patients with a very high risk of fracture, in whom study and treatment should not be delayed, given the high morbidity and mortality associated with fractures and the possible need to require hospitalization for an incident fracture, although, if necessary, some interventions could be postponed for 1 to 2 months during the critical period of the pandemic.
- Priority C. These are clinically stable patients, with no recent history of fragility fracture and no clinical suspicion of life-threatening conditions in the short- or medium-term, in whom specific treatments or services can be deferred for some time until sanitary conditions allow, without negatively

affecting the results as long as the patient's clinical conditions remain unchanged. It is essential to consider these factors even if the patient is already under osteoporosis treatment since compliance and duration of therapy may modify the probability of new fragility fractures in a short period (less than 12 months) (Table 1).

Absolute fracture risk assessment

The absolute risk of fragility fracture in patients with osteoporosis can vary widely from a probability of less than one fragility fracture in every 100 patients in 10 years to more than one fracture for every two patients in the next 12 months [10–12].

It is advisable to determine the absolute risk of fracture of each patient, particularly the immediate risk, and assess the factors associated with fracture susceptible to modification, including those extra-osseous factors.

DXA bone scans

DXA bone scans should be considered an elective procedure, and all patients with a recent fragility fracture should be considered being at intermediate or high risk of fracture even without DXA or FRAX assessment.

As DXA services are frequently performed in hospital facilities, and many of those facilities are dedicated to the COVID-19 crisis or emergency management, “elective” radiology may be severely limited. Therefore, it may be necessary to postpone DXA scans when health authorities recommend suspending elective imaging procedures. If a DXA bone scan, vertebral morphometry, or a bilateral full-length femur image (FFI) is required, determine the priority of the need for care. If a priority A or B patient is identified, it is preferable to choose DXA centers geographically located outside of COVID-19 hospitals or laboratory facilities. If it is not possible to perform a DXA scan, it is feasible to stratify the risk of fracture in adults without prior treatment using the Fracture Risk Assessment Tool (FRAX). However, FRAX may underestimate the risk of new fractures in patients with an incident or recent fracture [10, 20]; therefore, patients with a recent fragility fracture should be considered patients at imminent fracture risk [10–12], regardless of FRAX assessment. It is essential to keep a list of patients without DXA to perform it ideally within the next 6 months or once the care services unit restart their activities.

Clinical laboratory

It is necessary for all candidates for the treatment of osteoporosis to determine the etiology, rule out secondary causes, and

Table 1 Need for care priority levels

Priority	Decision-making	Appropriate interventions	Representative case
A	<i>Patient is clinically unstable</i> <i>(OR)</i> <i>Has a condition that puts the patient's life at imminent risk</i> <i>(OR)</i> <i>A short delay would significantly alter the prognosis</i>	<i>It requires prioritizing immediate care and not deferring attention</i>	<i>A 78-year-old patient with an incident hip fracture</i> <i>A 70-year-old patient with an incident clinical vertebral fracture</i> <i>An 82-year-old patient on antiresorptive treatment with evidence of hypocalcemia</i>
B	<i>Patient is clinically stable</i> <i>(AND)</i> <i>Has very high fracture risk^a</i>	<i>Treatment should not be delayed, although if necessary, some interventions could be postponed for 1 to 2 months during the critical period of the pandemic</i>	<i>73-year-old patient with a major osteoporotic fracture during the previous year.</i> <i>A 65-year-old patient with a fracture while on chronic use of glucocorticoids at high doses</i> <i>A 68-year-old patient with several prevalent vertebral fractures on denosumab treatment during the 6 months since the previous denosumab dose</i>
C	<i>The patient is clinically stable</i> <i>(AND)</i> <i>Has risk factors or diagnosed osteoporosis, without recent history of fragility fracture, or other risk factors for imminent fracture</i>	<i>Some specific treatments or services can be deferred for some time until sanitary conditions allow, without negatively affecting the results as long as the patient's clinical conditions remain unchanged</i>	<i>A 65-year-old patient, without previous fracture and a DXA scan acquired during the previous year with low bone mineral density values</i> <i>A 62-year-old osteoporotic patient on bisphosphonates treatment, without previous fracture</i>

^a Consider patients with a recent fragility fracture (e.g., within the past 12 months), fractures while on approved osteoporosis therapy, multiple fractures, fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids), very low T-score (less than -3.0), or a high risk for falls or history of recent injurious falls

factors that contribute to bone loss, and identify conditions that may contraindicate specific pharmacological therapy.

In candidates who receive a potent antiresorptive such as denosumab or an IV bisphosphonate, and who have never received antiresorptive treatment, it is necessary to measure serum calcium, creatinine, and vitamin D to identify patients at high risk of hypocalcemia; this is particularly important in the presence of impaired kidney function.

According to previous year's laboratory test results (if available), new lab procedures could be delayed in clinically stable patients with normal kidney function.

In patients with clinical conditions associated with hypocalcemia (e.g., hypoparathyroidism, use of loop diuretics, malabsorption), impaired renal function, or clinically unstable, it is necessary to determine serum calcium and vitamin D levels before receiving any pharmacological treatment [21].

During the COVID-19 pandemic, the risk of fracture, the possible etiology of bone loss, and the patient's clinical condition will determine the priority to attend or not a clinical laboratory facility.

If a laboratory test is necessary, explore if home sampling could be an option, or check if the laboratory facility has a recommended schedule and a non-COVID area in order to avoid cross-contamination as much as possible.

Pharmacologic treatment

Patients with osteoporosis should start, continue, or adjust treatment without delay. To date, there is no evidence that any osteoporosis drug affects the risk or susceptibility to contagion of SARS-CoV-2 virus or affects in some way the clinical course of COVID-19. In the case of considering it strictly necessary to postpone a medical intervention due to the pandemic, take into account the prioritization recommended above.

Pharmacologic treatment initiation

Patients with priority A and B need to start treatment immediately, considering their high risk of fracture. For those patients in whom it is not possible to start pharmacological treatment, it is necessary to make a register and determine a strategy to start it as soon as possible.

The choice of appropriate pharmacological therapy should be consistent with the elements of efficacy and safety of the drug, the specific clinical characteristics of each patient, and their preferences when possible.

Osteoporosis medications approved for use in Latin-American countries include denosumab, zoledronate, alendronate, risedronate, ibandronate, raloxifene, bazedoxifene, tibolone, and teriparatide. They have no contraindications associated with COVID-19. However, it is convenient to keep in mind the following considerations:

The use of zoledronate and denosumab may require a visit to the doctor for its application. Intravenous bisphosphonates occasionally produce moderate to severe inflammatory reactions [22–24] that may require medical observation and could be confused with COVID-19 symptoms. Therefore, in areas with a high COVID-19 burden and limited availability of diagnostic tests, it may be appropriate to consider other therapies, particularly in naive patients, in whom the inflammatory reactions associated with bisphosphonates are more common and intense. This concern about the acute reaction with IV bisphosphonates has to be balanced with both the convenience and the possible logistical problem of an IV dose and the strong advantage of long-lasting effectiveness without frequent return for another dose [25].

Selective estrogen receptor modulators (SERMS) are associated with a slight risk of thromboembolic events [26–29]. As some cases of COVID-19 have a hypercoagulability component [30], it is convenient to evaluate the risk-benefit relationship carefully.

All osteoporosis treatment patients should receive an adequate supply of 1200 mg of calcium and 1500–2000 IU of vitamin D daily. Patients at risk for D deficiency (e.g., obese, impaired kidney function) may require a personalized vitamin D dose, ideally according to laboratory results when available [31, 32].

It should be considered that some services and supplies, such as specialized drug parcel services, medications, and supplements, may have limited availability during the pandemic.

Treatment continuity and temporary interruption of pharmacologic treatment

All patients at high risk for fragility fracture and those with prior hip or vertebral fracture should continue treatment without interruption. “Drug holiday” is not recommended, especially in patients with several fracture risk factors, because anti-fracture efficacy is not maintained. They could increase the risk of new vertebral and non-vertebral fragility fractures [16, 33].

During the phases of increased transmission of SARS-CoV-2 disease, it is likely that there are some significant limitations to administering or taking medications for osteoporosis. Access to parenteral drug dosing centers may be limited, or there may be disruptions in some supply chains.

For most patients, the interruption of 1–3 months in the pharmacological treatment of osteoporosis may not imply significant risks, except for those receiving denosumab, as patients who discontinue its administration show a rapid rebound of bone remodeling markers and a decrease in BMD

[34, 35], and vertebral fractures have been reported to occur as early as the seventh month after the last denosumab injection, particularly in patients with previous fractures [36, 37]. Therefore, in patients receiving denosumab treatment, a new dose should be guaranteed within 4 weeks after the end of the 6-month period after its last dose [21, 38]. If arrangements for the next dose of denosumab cannot be made, consider switching to a bisphosphonate.

In patients under teriparatide regimens who have difficulty continuing their treatment, it is possible that 3 months without daily application does not significantly affect their evolution and prognosis, since no rebound effect has been observed when stopping teriparatide application, and the pivotal study suggests that some anti-fracture effect persists for 18 months after stopping treatment [39, 40]. Additionally, the gain in BMD achieved with intermittent regimens of 3 months of application, alternated with three free months for 4 years, is similar to that obtained with the standard approved daily application of teriparatide in 2 years [41, 42]. However, intermittent application should not be considered a recommendation, and a prolonged interruption of treatment should be avoided since it may entail a progressive decrease in BMD and a possible increase in fracture risk. Therefore, in patients who cannot continue teriparatide therapy, and in those who completed 18 or 24 months on teriparatide, sequential therapy is recommended either with denosumab or bisphosphonates [21, 38].

In the case of oral bisphosphonates, and perhaps IV ibandronate, a delay up to 6 months does not seem to affect the patient's evolution significantly [16, 43]. However, stopping treatment for more than 3 months on any osteoporosis medication is not recommended. A possible exception is IV zoledronate since data from the extension to the HORIZON PFT study indicate that although persistence with annual dosing between years 3 and 6 results in fewer vertebral fractures than does drug discontinuation at year 3, fracture rates remain much lower in the discontinuation group than in the original placebo group [44]. While those patients at high risk for vertebral fracture, including those with recent incident morphometric vertebral fracture and/or hip BMD T-score ≤ -2.5 , will benefit most from continuing of osteoporosis therapy, consider selecting medical facilities different and geographically distant to hospitals, applying the drug at home, drive-through application modules, delivering the medications and self-application at home, and temporary or definitive substitution of a parenteral drug for oral medications.

Clinicians must consider that the temporary or definitive substitution of a parenteral drug for oral drugs requires ascertainment that there is no contraindication (e.g., esophagitis or renal deterioration associated with bisphosphonates).

It is necessary to recognize that self-application may need training and monitoring. For that purpose, we can take advantage of available technological resources, such as video training.

Management of osteoporosis in the patient with incident fragility fracture

Despite confinement, and the relative decline in mobility, during the COVID-19 pandemic, the incidence of hip fracture has not decreased [45], and it is still among the leading causes of care in emergency services.

Due to the saturation of medical services and hospital re-conversion, several fracture liaison services (FLS) have been disrupted, compromising secondary fracture prevention. Since patients with incident fragility fractures have a very high risk of a new fracture during the first year following the fracture [46], detection and management of osteoporosis, including rehabilitation and fall prevention, are essential parts of treatment that should not be overlooked [47, 48]. In these patients, when a DXA bone scan is not possible, the diagnosis of osteoporosis can be preliminarily made based on clinical and radiographic findings [18], so that DXA scan can be postponed until epidemiological conditions allow it or requested in an out-of-hospital DXA center. Laboratory tests should be performed before starting osteoporosis medication [49]. Calcium and vitamin D supplementation must be prescribed before hospital discharge [47, 50]. It is also advisable to make the appropriate arrangements to reduce the hospital stay and transfers within the hospital for patients and staff, to reduce cross-contamination risk.

When it is not feasible to complete the clinical and laboratory evaluation or start treatment during the hospitalization, for the fractured patient, it is necessary to complete the study protocol and start treatment later. In that case, telemedicine can be useful [48, 51].

In hospitalized COVID-19 positive patients, special care should be taken when prescribing antiresorptive treatment, as it could worsen the hypocalcemia that has been associated with COVID-19 infection [52–54]. It could be considered either to defer the treatment initiation or to stop it temporarily during hospitalization.

Patients with suspected or confirmed coronavirus disease (COVID-19)

Subjects with suspected or confirmed COVID-19 should be treated according to their care priority. However, we recommend that the osteoporosis diagnostic approach and treatment be deferred until the SARS-CoV-2 infection has been ruled out, or the patient has been fully recovered.

In the event of an incident fragility fracture, the patient must receive trauma care, following strict personal protection measures.

In patients in osteoporosis treatment who are infected by SARS-CoV-2, it is possible to discontinue the osteoporosis drugs during several weeks until complete recovery from COVID-19, particularly in those with moderate to severe disease-

Physical activity, exercise, and nutrition

Regular physical activity and good nutrition are important components of osteoporosis management and should be continued during COVID-19.

Physical activity and exercise

Because some precautionary measures for COVID-19 may affect the amount of physical activity that patients perform, patients and caregivers must pay special attention to physical activity and the time spent by patients sitting or reclining [55] since rapid bone and muscular detriment on mass and function are associated with unloading bone and muscle [56], and inactivity is associated not only with the loss of bone and muscle but a decrease in motor skills. It is strongly recommended that patients engage in a regular physical activity plan and avoid sedentary behaviors. Patients should be as physically active as their abilities and conditions allow, breaking up sedentary time [57].

It is recommended that the exercise program for osteoporosis patients be designed and supervised by a clinical exercise expert. Those programs should consider specific exercises for static and dynamic balance, motor coordination, proprioception, and muscular strength [58]. A correct body posture during the execution of exercise routines, and activities at home reduces fracture risk, so it is essential to prioritize the postural training [59].

In those patients who already had an exercise program, it is necessary to adjust it to the available resources in confinement. If close supervision is not possible, clinical exercise programs should be adjusted based on patient safety, and telerehabilitation may play a role in some instances.

Calcium and vitamin D

During confinement, eating habits can vary significantly and affect the intake of certain essential nutrients. Therefore, it is advisable to guarantee a daily calcium intake of 1000 to 1500 mg [60], preferably through the diet, mainly from fermented dairy (like yogurt) and cheese, as their consumption, beyond calcium and vitamin D content, is associated with a lower risk of fractures [61–64]. Subjects with insufficient calcium intake or who do not include dairy products in the diet, or those in treatment with an antiresorptive agent, should receive a calcium supplement [62].

Since low serum calcidiol levels are common worldwide, including Latin American countries [65–67], and self-confinement probably decreases sun exposure, a vitamin D dose of 1500 to 2000 IU/day is recommended to reach or maintain adequate serum calcidiol levels, even without a recent determination of vitamin D levels [21, 31, 68]. However,

if a marked vitamin D deficiency is observed (≤ 20 ng/mL), a higher supplementation pattern may apply [69].

During lockdown, consuming vitamin D-rich foods (like high-fat fish, red meat, and egg yolks) and fortified ones is highly advised, due to their overall nutrient contribution [70–72]. Similarly, sufficient and safe sun exposure should be advised when possible, for instance, 5 to 15 min of sun exposure (arms and legs uncovered) between 10 am and 3 pm.[73].

Protein

Due to self-confinement, the decreased physical activity might lead to a reduced anabolic sensitivity to protein intake [74]. Older adults should consume 1.0 to 1.2 g of protein, per kg of body weight, per day [61–63, 75–80], following an evenly distributed pattern [81]. Each meal should contain at least 30 g of protein or the equivalent to 0.4 g protein/kg [82–85], and most of the protein should come from animal sources, because of their availability and essential amino acid profile (including leucine) [86, 87].

The patient and family members should be committed to nutritional surveillance to identify changes in weight or eating behaviors that could be considered risky and may harm bone mass, body composition, and functional capacity and increase the risk of falls. If nutritional risk factors or lack of compliance are detected, it is advisable to request the nutrition professional's assessment, which could be through the tele-nutrition modality. This form of remote care aims to provide nutritional care to patients in contexts in which, for security reasons, they cannot go to receive in-person consultation [88].

Conclusion

Infection by SARS-Cov-2 (COVID-19) keeps rising worldwide, and mitigation measures are expected to continue for several more months. Healthcare systems will likely be affected for a long time. Consequently, an organized response for delivering medical attention to the patients at risk of fragility fracture is necessary. This joint position of Mexican and Latin American Medical Societies provides an updated guide for the prevention, diagnosis, and treatment of osteoporotic patients in the face of possible clinical scenarios posed by the COVID-19 crisis.

We recommend that whenever possible, existing osteoporosis guidelines should be observed during COVID-19 contingency. Otherwise, stratify the need for care at three priority levels based on patient absolute fracture risk and clinical stability. Therefore, patients with intermediate and high fracture risk must receive immediate attention, although in some epidemic scenarios, selected laboratory tests and intervention could be delayed 1 or 2 months with relatively low risk of

complications. Patients with an incident fragility fracture are at imminent risk of a new fracture and must receive treatment even without a DXA scan.

Osteoporosis pharmacologic treatment must be based on safety and efficacy. Although there is no evidence that any osteoporosis drug affects the risk or susceptibility for the contagion of SARS-CoV-2 virus or affects the clinical course of COVID-19, some specific observations are made to aid clinicians to select the optimal treatment for each patient.

Finally, we hope that this guideline is useful in a wide variety of possible Latin America scenarios. We recognize we are learning more about COVID-19, and an update of these guides may be necessary.

Disclaimer

Disruption of health services caused by the COVID-19 pandemic, the urgency of having adequate recommendations in the care of patients at high risk of fracture, and the low level of evidence available on multiple aspects of the behavior of the SARS-CoV-2 imply that most of the recommendations are based on levels of evidence III and IV and expert opinions. Since information may undergo changes in the coming months with the advent of new knowledge, the experts continue to meet and evaluate new information to update the recommendations if necessary.

Acknowledgments We are thankful to the members of expert panel, Maria de los Angeles Aguilera-Barreiro, PhD; Jose Manuel Aguilera Zepeda, MD; Hilario Avila Armengol, MD; Cuauhtemoc Celis Gonzalez MD; Rolando Espinosa Morales, MD, MSc; Pedro Alberto Garcia Hernandez, MD; Alejandro Gaytan-Gonzalez, BSc(NutrSc); Jose Maximo Gomez Acevedo, MD; Miguel Flores Castro, PhD; Roberto Gabriel Gonzalez-Mendoza, MSc; Hugo Gutierrez Hermosillo, PhD; Alan Christopher Guzman-Rico, BSc(PHS); Rafael Alfonso Jimenez Umbarila, MD; Roberto Enrique Lopez Cervantes, MD; Juan Ricardo Lopez Taylor, MD; MSc; Pedro Nel Rueda Plata, MD; Jose Manuel Perez Atanasio, MD; Victor Mercado Cardenas, MD; Jorge Morales Torres, MD; Jorge Morales Vargas, MSc; Alfonso Murillo Uribe, MD; Maria del Pilar De la Peña-Rodriguez, MD; Juan Carlos Perez Barba, MD, MSc; Sergio Quintero Hernandez, MD; Douglas Fernando Solorzano Moreno; Valeria Taylor Sanchez, MD; Jose Francisco Torres-Naranjo, MD, MSc; and Juan Carlos Viveros Garcia, MD.

We want to extend our sincere appreciation to Dr. Michael McClung for his valuable assistance and advice in drafting the manuscript.

Endorsements This consensus report is endorsed by the following societies:

Mexican Association of Bone and Mineral Metabolism (AMMOM)
National College of Geriatric Medicine (CONAMEGER)
Latin American Federation of Endocrinology (FELAEN)
Mexican Federation of Colleges of Obstetrics and Gynecology (FEMECONG)

Mexican Federation of Colleges of Orthopedics and Traumatology (FEMEOT)

Institute of Sciences Applied for Physical Activity and Sports of the University of Guadalajara (ICAAFYD)

Authors' contributions FTN, PPR, RLC, JMT, and JMV conceived the idea of the statement and drafted the original manuscript. All authors (FTN, PPR, RLC, JMT, JMV, HGH, AGR, RGM, PRP, MFC, CCG, REM, SQH, and JLT) critically revised the manuscript for intellectual content, language, and presentation. All authors approved the final version of the article.

Compliance with ethical standards

Conflicts of interest FTN has nothing to declare for the context of this paper, but several ad hoc consultancies/speaking honoraria and/or research funding from Amgen Eli Lilly, GSK, Sanofi-Aventis.

PPR has nothing to declare for the context of this paper, but several ad hoc consultancies/speaking honoraria and/or research funding from Amgen, Eli Lilly.

JMT has nothing to declare for the context of this paper, but several ad hoc consultancies/speaking honoraria and/or research funding from AbbVie, Amgen, Eli Lilly, Merck; Roche; and Sanofi-Aventis.

PRP has nothing to declare for the context of this paper, but several ad hoc consultancies/speaking honoraria and/or research funding from Amgen, Eli Lilly, Sanulac, Fama de Colombia. Position on advisory medical board 6 years ago.

RLC, JMT, JMV, HGH, AGR, RGM, MFC, CCG, REM, SQH, and JLT have no conflicts of interest.

Informed consent For this type of study, an informed consent is not required.

Statement of human and animal rights This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Gorbalenya AE, Baker SC, Baric RS, de Groot RJ, Drosten C, Gulyaeva AA, Haagmans BL, Lauber C, Leontovich AM, Neuman BW, Penzar D, Perlman S, Poon LLM, Samborskiy DV, Sidorov IA, Sola I, Ziebuhr J (2020) The species severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* 5:536–544. <https://doi.org/10.1038/s41564-020-0695-z>
- WHO (2020) Coronavirus disease (COVID-19) situation report – 189. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200727-covid-19-sitrep-189.pdf?sfvrsn=b93a6913_2
- Li X, Wang W, Zhao X, Zai J, Zhao Q, Li Y, Chaillon A (2020) Transmission dynamics and evolutionary history of 2019-nCoV. *J Med Virol* 92:501–511. <https://doi.org/10.1002/jmv.25701>
- Zhang S, Diao MY, Yu W, Pei L, Lin Z, Chen D (2020) Estimation of the reproductive number of novel coronavirus (COVID-19) and the probable outbreak size on the Diamond Princess cruise ship: a data-driven analysis. *Int J Infect Dis* 93:201–204. <https://doi.org/10.1016/j.ijid.2020.02.033>
- Tang B, Wang X, Li Q, Bragazzi NL, Tang S, Xiao Y, Wu J (2020) Estimation of the transmission risk of the 2019-nCoV and its implication for public health interventions. *J Clin Med* 9:462. <https://doi.org/10.3390/jcm9020462>
- Morales-Torres J, Aceves-Ávila FJ (2020) Rheumatologists in the COVID-19 era: will there be a new role for the rheumatologist in the care of rheumatic patients? *Clin Rheumatol* 39:3177–3183. <https://doi.org/10.1007/s10067-020-05380-1>

7. Kanis JA, Odén A, McCloskey EV, Johansson H, Wahl DA, Cooper C (2012) A systematic review of hip fracture incidence and probability of fracture worldwide. *Osteoporos Int* 23:2239–2256
8. Clark P, Lavielle AEP, Esperanza FFÆ, John AÆ, Steven AKÆ, Salmero J (2005) Incidence rates and life-time risk of hip fractures in Mexicans over 50 years of age: a population-based study. 2025–2030. <https://doi.org/10.1007/s00198-005-1991-4>
9. Gonzalez EDDL, Mendivil LLL, Garza DPS, Hermosillo HG, Chavez JHM, Corona RP (2018) Low handgrip strength is associated with a higher incidence of pressure ulcers in hip fractured patients. *Acta Orthop Belg* 84:284–291
10. Pinedo-Villanueva R, Charokopou M, Toth E, Donnelly K, Cooper C, Prieto-Alhambra D, Libanati C, Javai MK (2019) Imminent fracture risk assessments in the UK FLS setting: implications and challenges. *Arch Osteoporos* 14. <https://doi.org/10.1007/s11657-019-0569-2>
11. Johansson H, Siggeirsdóttir K, Harvey NC, Odén A, Gudnason V, McCloskey E, Sigurdsson G, Kanis JA (2017) Imminent risk of fracture after fracture. *Osteoporos Int* 28:775–780. <https://doi.org/10.1007/s00198-016-3868-0>
12. Laurs-Van Geel TACM, Center JR, Geusens PP, Dinant GJ, Eisman JA (2010) Clinical fractures cluster in time after initial fracture. *Maturitas* 67:339–342. <https://doi.org/10.1016/j.maturitas.2010.09.002>
13. Bliuc D, Nguyen ND, Milch VE, Nguyen TV, Eisman JA, Center JR (2009) Mortality risk associated with low-trauma osteoporotic fracture and subsequent fracture in men and women. *JAMA J Am Med Assoc* 301:513–521. <https://doi.org/10.1001/jama.2009.50>
14. Morales-Torres J, Gutiérrez-Ureña S (2004) The burden of osteoporosis in Latin America. *Osteoporos Int* 15. <https://doi.org/10.1007/s00198-004-1596-3>
15. Zanchetta J (2012) The Latin America Regional Audit. Epidemiology, cost & burden of osteoporosis in 2012. http://www.iofbonehealth.org/sites/default/files/media/PDFs/RegionalAudits/2012-Latin_America_Audit_0_0.pdf
16. Dennison EM, Cooper C, Kanis JA, Bruyère O, Silverman S, McCloskey E, Abrahamson B, Prieto-Alhambra D, Ferrari S, IOF Epidemiology/Quality of Life Working Group (2019 Sep) Fracture risk following intermission of osteoporosis therapy. *Osteoporos Int* 30(9):1733–1743. <https://doi.org/10.1007/s00198-019-05002-w>
17. American Society for Bone and Mineral Research (ASBMR) AA of, Clinical Endocrinologists (AACE), Endocrine Society, European Calcified Tissue Society (ECTS) and NOF (NOF) (2020) joint guidance on osteoporosis management in the era of COVID-19. <https://www.endocrine.org/-/media/endocrine/files/membership/joint-statement-on-covid19-and-osteoporosis-final.pdf>. Accessed 10 Jun 2020
18. Capture the Fracture (CTF) IOF (IOF), International Osteoporosis Foundation (2020) COVID-19 para todos los centros de CTF/FLS. In: COVID statement SP. http://www.worldosteoporosisday.org/CTF_website/COVid-statement-SP.pdf. Accessed 10 Jun 2020
19. International Society of Clinical Densitometry (ISCD) (2020) ISCD Initial COVID-19 Statement March 30, 2020. https://www.iscd.org/about-iscd/iscd-covid-19-dxa-guidance/?utm_source=Informz&utm_medium=Email&utm_campaign=ISCD. Accessed 15 Jun 2020
20. Kanis JA, Johansson H, Harvey NC, Gudnason V, Sigurdsson G, Siggeirsdóttir K, Lorentzon M, Liu E, Vandenput L, McCloskey EV (2020) Adjusting conventional FRAX estimates of fracture probability according to the recency of sentinel fractures. *Osteoporos Int*. <https://doi.org/10.1007/s00198-020-05517-7>
21. Camacho PM, Petak SM, Binkley N, Diab DL, Eldeiry LS, Farooki A, Harris ST, Hurley DL, Kelly J, Lewiecki EM, Pessah-Pollack R, McClung M, Wimalawansa SJ, Watts NB (2020) American Association of Clinical Endocrinologists/American College of Endocrinology Clinical Practice Guidelines for the Diagnosis and Treatment of Postmenopausal Osteoporosis- 2020 Update Executive Summary. *Endocr Pract* 26:564–570. <https://doi.org/10.4158/GL-2020-0524>
22. Reid IR, Gamble GD, Mesenbrink P, Lakatos P, Black DM (2010) Characterization of and risk factors for the acute-phase response after zoledronic acid. *J Clin Endocrinol Metab* 95:4380–4387. <https://doi.org/10.1210/jc.2010-0597>
23. Adami S, Bhalla AK, Dorizzi R, Montesanti F, Rosini S, Salvagno G, Lo Cascio V (1987) The acute-phase response after bisphosphonate administration. *Calcif Tissue Int* 41:326–331. <https://doi.org/10.1007/BF02556671>
24. Schweitzer DH, Oostendorp-van de Ruit M, van der Pluijm G, Löwik CWGM, Papapoulos SE (1995) Interleukin-6 and the acute phase response during treatment of patients with Paget's disease with the nitrogen-containing bisphosphonate dimethylaminohydroxypropylidene bisphosphonate. *J Bone Miner Res* 10:956–962. <https://doi.org/10.1002/jbmr.5650100617>
25. Reid IR, Black DM, Eastell R, Bucci-Rechtweg C, Su G, Hue TF, Mesenbrink P, Lyles KW, Boonen S (2013) Reduction in the risk of clinical fractures after a single dose of zoledronic acid 5 milligrams. *J Clin Endocrinol Metab* 98:557–563. <https://doi.org/10.1210/jc.2012-2868>
26. Barrett-Connor E, Mosca L, Collins P, Geiger MJ, Grady D, Kornitzer M, McNabb MA, Wenger NK (2006) Effects of raloxifene on cardiovascular events and breast cancer in postmenopausal women. *N Engl J Med* 355:125–137. <https://doi.org/10.1056/NEJMoa062462>
27. Mosca L, Grady D, Barrett-Connor E, Collins P, Wenger N, Abramson BL, Paganini-Hill A, Geiger MJ, Dowsett SA, Amewou-Atisso M, Kornitzer M (2009) Effect of raloxifene on stroke and venous thromboembolism according to subgroups in postmenopausal women at increased risk of coronary heart disease. *Stroke* 40:147–155. <https://doi.org/10.1161/STROKEAHA.108.518621>
28. Christiansen C, Chesnut CH, Adachi JD, Brown JP, Fernandes CE, Kung AWC, Palacios S, Levine AB, Chines AA, Constantine GD (2010) Safety of bazedoxifene in a randomized, double-blind, placebo- and active-controlled phase 3 study of postmenopausal women with osteoporosis. *BMC Musculoskelet Disord* 11: <https://doi.org/10.1186/1471-2474-11-130>
29. Silverman SL, Chines AA, Kendler DL, Kung AWC, Tegljbærg CS, Felsenberg D, Mairon N, Constantine GD, Adachi JD (2012) Sustained efficacy and safety of bazedoxifene in preventing fractures in postmenopausal women with osteoporosis: results of a 5-year, randomized, placebo-controlled study. *Osteoporos Int* 23: 351–363. <https://doi.org/10.1007/s00198-011-1691-1>
30. Cao W, Li T (2020) COVID-19: towards understanding of pathogenesis. *Cell Res* 30:367–369. <https://doi.org/10.1038/s41422-020-0327-4>
31. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM (2011) Evaluation, treatment, and prevention of vitamin D deficiency: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* 96:1911–1930
32. Shoback D, Rosen CJ, Black DM, Cheung AM, Murad MH, Eastell R (2020) Pharmacological management of osteoporosis in postmenopausal women: an endocrine society guideline update. *J Clin Endocrinol Metab* 105
33. McClung MR, Wagman RB, Miller PD, Wang A, Lewiecki EM (2017) Observations following discontinuation of long-term denosumab therapy. *Osteoporos Int* 28:1723–1732. <https://doi.org/10.1007/s00198-017-3919-1>
34. Bolognese MA, Yuen CK, Kendler DL, Miller PD, Yang YC, Grazette L, San Martin JGJ (2011) Effects of denosumab treatment and discontinuation on bone mineral density and bone turnover

- markers in postmenopausal women with low bone mass. *J Clin Endocrinol Metab* 96:972–980
35. Popp AW, Varathan N, Buffat H, Senn C, Perrelet R, Lippuner K (2018) Bone mineral density changes after 1 year of denosumab discontinuation in postmenopausal women with long-term denosumab treatment for osteoporosis. *Calcif Tissue Int* 103:50–54. <https://doi.org/10.1007/s00223-018-0394-4>
 36. Cummings SR, Ferrari S, Eastell R, Gilchrist N, Jensen JEB, McClung M, Roux C, Törring O, Valter I, Wang AT, Brown JP (2018) Vertebral fractures after discontinuation of denosumab: a post hoc analysis of the randomized placebo-controlled FREEDOM trial and its extension. *J Bone Miner Res* 33:190–198. <https://doi.org/10.1002/jbmr.3337>
 37. Gonzalez-Rodriguez E, Aubry-Rozier B, Stoll D, Zaman K, Lamy O (2020) Sixty spontaneous vertebral fractures after denosumab discontinuation in 15 women with early-stage breast cancer under aromatase inhibitors. *Breast Cancer Res Treat* 179:153–159. <https://doi.org/10.1007/s10549-019-05458-8>
 38. Kanis JA, Cooper C, Rizzoli R, Reginster JY (2019) Executive summary of the European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Calcif Tissue Int* 104:235–238. <https://doi.org/10.1007/s00223-018-00512-x>
 39. Neer RM, Arnaud CD, Zanchetta JR, Prince R, Gaich GA, Reginster J-Y, Hodsman AB, Eriksen EF, Ish-Shalom S, Genant HK, Wang O, Mitlak BH (2001) Effect of parathyroid hormone (1–34) on fractures and bone mineral density in postmenopausal women with osteoporosis. *Obstet Gynecol Surv.* <https://doi.org/10.1097/00006254-200110000-00018>
 40. Lindsay R, Scheele WH, Neer R, Pohl G, Adami S, Mautalen C, Reginster JY, Stepan JJ, Myers SL, Mitlak BH (2004) Sustained vertebral fracture risk reduction after withdrawal of teriparatide in postmenopausal women with osteoporosis. *Arch Intern Med* 164:2024–2030. <https://doi.org/10.1001/archinte.164.18.2024>
 41. Cosman F, Nieves JW, Zion M, Garrett P, Neubort S, Dempster D, Lindsay R (2015) Daily or cyclical teriparatide treatment in women with osteoporosis on no prior therapy and women on alendronate. *J Clin Endocrinol Metab* 100:2769–2776. <https://doi.org/10.1210/jc.2015-1715>
 42. Cosman F, Nieves JW, Roimisher C, Neubort S, McMahon DJ, Dempster DW, Lindsay R (2019) Administration of teriparatide for four years cyclically compared to two years daily in treatment naïve and alendronate treated women. *Bone* 120:246–253. <https://doi.org/10.1016/j.bone.2018.10.020>
 43. Watts NB, Chines A, Olszynski WP, McKeever CD, McClung MR, Zhou X, Grauer A (2008) Fracture risk remains reduced one year after discontinuation of risedronate. *Osteoporos Int* 19:365–372. <https://doi.org/10.1007/s00198-007-0460-7>
 44. Compston JE, Bilezikian JP (2012) Bisphosphonate therapy for osteoporosis: the long and short of it. *J Bone Miner Res* 27:240–242. <https://doi.org/10.1002/jbmr.1542>
 45. James N, Hadfield ACG (2020) The evolving COVID-19 effect on hip fracture patients. *Injury* 51(7):1411–1412
 46. Banefelt J, Åkesson KE, Spångéus A, Ljunggren O, Karlsson L, Ström O, Orsäter G, Libanati C, Toth E (2019) Risk of imminent fracture following a previous fracture in a Swedish database study. *Osteoporos Int.* <https://doi.org/10.1007/s00198-019-04852-8>
 47. Napoli N, Elderkin AL, Kiel DP, Khosla S (2020) Managing fragility fractures during the COVID-19 pandemic. *Nat Rev Endocrinol*:5–6. <https://doi.org/10.1038/s41574-020-0379-z>
 48. Upadhyaya GK, Iyengar K, Jain VK, Vaishya R (2020) Challenges and strategies in management of osteoporosis and fragility fracture care during COVID-19 pandemic. *J Orthop* 21:287–290. <https://doi.org/10.1016/j.jor.2020.06.001>
 49. Girgis CM, Clifton-Bligh RJ (2020) Osteoporosis in the age of COVID-19. *Osteoporos Int* 31:1189–1191. <https://doi.org/10.1007/s00198-020-05413-0>
 50. Chakhtoura M, Napoli N, El Hajj G (2020) Commentary: myths and facts on vitamin D amidst the COVID-19 pandemic. *Metab Clin Exp* 109:154276. <https://doi.org/10.1016/j.metabol.2020.154276>
 51. Joseph B, Hadeed G, Sadoun M, Rhee PM, Weinstein RS (2012) Video consultation for trauma and emergency surgical patients. *Crit Care Nurs Q* 35:341–345. <https://doi.org/10.1097/CNQ.0b013e318266c2f2>
 52. Di Filippo L, Formenti AM, Rovere-Querini P, Carlucci M, Conte C, Ciceri F, Zangrillo A, Giustina A (2020) Hypocalcemia is highly prevalent and predicts hospitalization in patients with COVID-19. *Endocrine* 68:475–478. <https://doi.org/10.1007/s12020-020-02383-5>
 53. Lippi G, South AM, Henry BM (2020) Electrolyte imbalances in patients with severe coronavirus disease 2019 (COVID-19). *Ann Clin Biochem* 57:262–265. <https://doi.org/10.1177/0004563220922255>
 54. Cappellini F, Brivio R, Casati M, Cavallero A, Contro E, Brambilla P (2020) Low levels of total and ionized calcium in blood of COVID-19 patients. *Clin Chem Lab Med* 1–3. <https://doi.org/10.1515/cclm-2020-0611>
 55. Chen P, Mao L, Nassiss GP, Harmer P, Ainsworth BE, Li F (2020) Coronavirus disease (COVID-19): the need to maintain regular physical activity while taking precautions. *J Sport Health Sci* 9: 103–104. <https://doi.org/10.1016/j.jshs.2020.02.001>
 56. Goodman CA, Hornberger TA, AGR (2015) Bone and skeletal muscle: key players in mechanotransduction and potential overlapping mechanisms. *Bone* 80:24–36. <https://doi.org/10.1016/j.bone.2015.04.014>
 57. Dalgleish TW, Mark J, Golden G, Perkins A-MJ, Barrett N, Lisa Feldman Barnard PJ (2010) World Health Organization: global recommendations on physical activity for health: WHO 2010. *J Exp Psychol Gen* 136:23–42
 58. Sinaki M, Brey RH, Hughes CA, Larson DR, Kaufman KR (2005) Significant reduction in risk of falls and back pain in osteoporotic-kyphotic women through a spinal proprioceptive extension exercise dynamic (SPEED) program. *Mayo Clin Proc* 80:849–855. <https://doi.org/10.4065/80.7.849>
 59. Sinaki M, Itoi E, Wahner HW, Wollan P, Gelzcer R, Mullan BP, Collins DA, Hodgson SF (2002) Stronger back muscles reduce the incidence of vertebral fractures: a prospective 10 year follow-up of postmenopausal women. *Bone* 30:836–841. [https://doi.org/10.1016/S8756-3282\(02\)00739-1](https://doi.org/10.1016/S8756-3282(02)00739-1)
 60. Lozano M, Manyes L, Peiró J, Ramada JM (2018) Nutrients associated with diseases related to aging: a new healthy aging diet index for elderly population. *Nutr Hosp* 35:1287–1297. <https://doi.org/10.20960/nh.1946>
 61. Stránský M, Ryšavá L (2009) Nutrition as prevention and treatment of osteoporosis. *Physiol Res* 58:S7–S11
 62. Torres-Morales J (2019) Osteoporosis: the other interventions. *Reumatol Clin* 15:185–187. <https://doi.org/10.1016/j.reuma.2019.05.002>
 63. Camacho PM, Petak SM, Binkley N, Clarke BL, Harris ST, Hurley DL, Kleerekoper M, Lewiecki EM, Miller PD, Narula HS, Pessah-Pollack R, Tangpricha V, Wimalawansa SJ, Watts NB (2016) American association of clinical endocrinologists and American college of endocrinology clinical practice guidelines for the diagnosis and treatment of postmenopausal osteoporosis - 2016. *Endocr Pract* 22:1–42. <https://doi.org/10.4158/EPI161435.GL>
 64. Geiker NRW, Mølgaard C, Iuliano S, Rizzoli R, Manios Y, van Loon LJC, Lecerf JM, Moschonis G, Reginster JY, Givens I, Astrup A (2020) Impact of whole dairy matrix on musculoskeletal health and aging—current knowledge and research gaps. *Osteoporos Int* 31:601–615. <https://doi.org/10.1007/s00198-019-05229-7>

65. Holick MF (2017) The vitamin D deficiency pandemic: approaches for diagnosis, treatment and prevention. *Rev Endocr Metab Disord* 18:153–165. <https://doi.org/10.1007/s11154-017-9424-1>
66. Arabi A, El Rassi R, El-Hajj Fuleihan G (2010) Hypovitaminosis D in developing countries-prevalence, risk factors and outcomes. *Nat Rev Endocrinol* 6:550–561. <https://doi.org/10.1038/nrendo.2010.146>
67. Carrillo-Vega MF, García-Peña C, Gutiérrez-Robledo LM, Pérez-Zepeda MU (2017) Vitamin D deficiency in older adults and its associated factors: a cross-sectional analysis of the Mexican Health and Aging Study. *Arch Osteoporos* 12:8. <https://doi.org/10.1007/s11657-016-0297-9>
68. Glade MJ (2012) A 21st century evaluation of the safety of oral vitamin D. *Nutrition* 28:344–356. <https://doi.org/10.1016/j.nut.2011.11.006>
69. Narvaez J, Maldonado G, Guerrero R, Messina OD, Ríos C (2020) Vitamin D megadose: definition, efficacy in bone metabolism, risk of falls and fractures. *Open Access Rheumatol Res Rev* 12:105–115. <https://doi.org/10.2147/OARRR.S252245>
70. Maurya VK, Aggarwal M (2017) Factors influencing the absorption of vitamin D in GIT: an overview. *J Food Sci Technol* 54:3753–3765. <https://doi.org/10.1007/s13197-017-2840-0>
71. Pilz S, März W, Cashman KD, Kiely ME, Whiting SJ, Holick MF, Grant WB, Pludowski P, Hilgsmann M, Trummer C, Schwetz V, Lerchbaum E, Pandis M, Tomaschitz A, Grübler MR, Gaksch M, Verheyen N, Hollis BW, Rejnmark L, Karras SN, Hahn A, Bischoff-Ferrari HA, Reichrath J, Jorde R, Elmadfa I, Vieth R, Scragg R, Calvo MS, van Schoor NM, Bouillon R, Lips P, Itkonen ST, Martineau AR, Lamberg-Allardt C, Zittermann A (2018) Rationale and plan for vitamin D food fortification: a review and guidance paper. *Front Endocrinol (Lausanne)* 9:373. <https://doi.org/10.3389/fendo.2018.00373>
72. Aguilera JM (2019) The food matrix: implications in processing, nutrition and health. *Crit Rev Food Sci Nutr* 59:3612–3629. <https://doi.org/10.1080/10408398.2018.1502743>
73. Holick MF (2005) The influence of vitamin D on bone health across the life cycle. *J Nutr* 135:2726S–2727S. <https://doi.org/10.1093/jn/135.11.2726S>
74. Oikawa SY, Holloway TM, Phillips SM (2019) The impact of step reduction on muscle health in aging: protein and exercise as countermeasures. *Front Nutr* 6:1–11. <https://doi.org/10.3389/fnut.2019.00075>
75. Rizzoli R, Biver E, Bonjour JP, Coxam V, Goltzman D, Kanis JA, Lappe J, Rejnmark L, Sahni S, Weaver C, Weiler H, Reginster JY (2018) Benefits and safety of dietary protein for bone health—an expert consensus paper endorsed by the European Society for Clinical and Economical Aspects of Osteoporosis, Osteoarthritis, and Musculoskeletal Diseases and by the International Osteoporosis Foundation. *Osteoporos Int* 29:1933–1948. <https://doi.org/10.1007/s00198-018-4534-5>
76. Bauer J, Biolo G, Cederholm T, Cesari M, Cruz-Jentoft AJ, Morley JE, Phillips S, Sieber C, Stehle P, Teta D, Visvanathan R, Volpi E, Boirie Y (2013) Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE study group. *J Am Med Dir Assoc* 14:542–559. <https://doi.org/10.1016/j.jamda.2013.05.021>
77. Coelho-Júnior HJ, Milano-Teixeira L, Rodrigues B, Bacurau R, Marzetti E, Uchida M (2018) Relative protein intake and physical function in older adults: a systematic review and meta-analysis of observational studies. *Nutrients* 10:1–16. <https://doi.org/10.3390/nu10091330>
78. Traylor DA, Gorissen SHM, Phillips SM (2018) Perspective: protein requirements and optimal intakes in aging: are we ready to recommend more than the recommended daily allowance? *Adv Nutr* 9:171–182. <https://doi.org/10.1093/advances/nmy003>
79. Rizzoli R, Bonjour JP (2004) Dietary protein and bone health. *J Bone Miner Res* 19:527–531. <https://doi.org/10.1359/JBMR.040204>
80. Darling AL, Manders RJF, Sahni S, Zhu K, Hewitt CE, Prince RL, Millward DJ, Lanham-New SA (2019) Dietary protein and bone health across the life-course: an updated systematic review and meta-analysis over 40 years. *Osteoporos Int* 30:741–761. <https://doi.org/10.1007/s00198-019-04933-8>
81. Hudson JL, Bergia RE, Campbell WW (2020) Protein distribution and muscle-related outcomes: does the evidence support the concept? *Nutrients* 12:1–22. <https://doi.org/10.3390/nu12051441>
82. Gaytán-González A, Ocampo-Alfaro MDJ, Arroniz-Rivera M, Torres-Naranjo F, González-Mendoza RG, Gil-Barreiro M, López-Taylor JR (2019, 2019) Inadequate protein intake at specific meals is associated with higher risk of impaired functionality in middle to older aged Mexican adults. *J Aging Res*. <https://doi.org/10.1155/2019/6597617>
83. Gaytán-González A, de Jesús Ocampo-Alfaro M, Torres-Naranjo F, Arroniz-Rivera M, González-Mendoza RG, Gil-Barreiro M, López-Taylor JR (2020) The consumption of two or three meals per day with adequate protein content is associated with lower risk of physical disability in Mexican adults aged 60 years and older. *Geriatr* 5:1–18. <https://doi.org/10.3390/geriatrics5010001>
84. Moore DR, Churchward-Venne TA, Witard O, Breen L, Burd NA, Tipton KD, Phillips SM (2015) Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. *J Gerontol Ser A Biol Sci Med Sci* 70:57–62. <https://doi.org/10.1093/gerona/glu103>
85. Loenneke JP, Loprinzi PD, Murphy CH, Phillips SM (2016) Per meal dose and frequency of protein consumption is associated with lean mass and muscle performance. *Clin Nutr* 35:1506–1511. <https://doi.org/10.1016/j.clnu.2016.04.002>
86. Gorissen SHM, Witard OC (2018) Characterising the muscle anabolic potential of dairy, meat and plant-based protein sources in older adults. *Proc Nutr Soc* 77:20–31. <https://doi.org/10.1017/S002966511700194X>
87. Berrazaga I, Micard V, Gueugneau M, Walrand S (2019) The role of the anabolic properties of plant-versus animal-based protein sources in supporting muscle mass maintenance: a critical review. *Nutrients* 11:1825. <https://doi.org/10.3390/nu11081825>
88. Peregrin T (2019) Telehealth is transforming health care: what you need to know to practice telenutrition. *J Acad Nutr Diet* 119:1916–1920. <https://doi.org/10.1016/j.jand.2019.07.020>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.