



CORRECTION

Correction To: Epidemiological derivation of flux-based critical levels for visible ozone injury in European forests

Pierre Sicard¹ · Alessandra De Marco² · Elisa Carrari³ · Laurence Dalstein-Richier⁴ · Yasutomo Hoshika³ · Ovidiu Badea⁵ · Diana Pitar⁵ · Silvano Fares⁶ · Adriano Conte⁶ · Ionel Popa⁵ · Elena Paoletti³

Published online: 7 January 2021
© Northeast Forestry University 2021

Correction to:

J. For. Res.

<https://doi.org/10.1007/s11676-020-01191-x>

In the Original publication of the article, the authors found an error in the text “We recommend CLef of 5 and 12 mmol m⁻² POD1 for broadleaved species and conifers, respectively” under the section abstract and conclusion. The corrected text is given below:

“We recommend CLef of 5 and 12 mmol m⁻² POD1 for conifers and broadleaved species, respectively”.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s11676-020-01191-x>.

✉ Pierre Sicard
psicard@argans.eu

¹ ARGANS, 260 route du Pin Montard, 06410 Biot, France

² ENEA, Via Anguillarese 301, 00123 Santa Maria di Galeria, Italy

³ IRET-CNR, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy

⁴ GIEFS, 69 avenue des Hespérides, 06300 Nice, France

⁵ INCDS, 128 Eroilor Blvd., 077030 Voluntari, Romania

⁶ CREA, Viale S. Margherita 80, 52100 Arezzo, Italy