CORRECTION

Correction to: Maternal environment and seed size are important for successful germination and seedling establishment of *Pterocarpus erinaceus* (Fabaceae)

Beda Innocent Adji^{1,2} · Doffou Sélastique Akaffou¹ ·
Philippe De Reffye² · Sylvie Sabatier²

Published online: 8 November 2021
© Northeast Forestry University 2021

Correction to: J. For. Res.

<https://doi.org/10.1007/s11676-021-01412-x>

In the original publication, the year mentioned in the references 1 and 4 of the document have two date errors.

The first reference is 2021 and not 2020: Adji BI, Akaffou SD, Kouassi KH, Houphouet YP, Duminil J, Sabatier S (2021) Bioclimate influence on seed germination and seedling morphology parameters in *Pterocarpus erinaceus* Poir., 1804 (Fabaceae). International Journal of Environment, Agriculture and Biotechnology 6(3):2456–1878.

Then the fourth reference is 2019 and not 2021: Akaffou SD, Kouamé KA, Gore BBN, Abessika YG, Kouassi KH,

Hamon P, Sabatier S, Duminil J (2019) Effect of the seeds provenance and treatment on the germination rate and plants growth of four forest trees species of Côte d'Ivoire. Journal of Forestry Research 32:161–169.

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s11676-021-01412-x>.

✉ Beda Innocent Adji
adjibedainnocent@gmail.com; beda-innocent.adji@cirad.fr

¹ Department for Seeds and Seedlings Production, Université Jean Lorougnon Guédé, Agroforestry UFR, BP 150, Daloa, Côte d'Ivoire

² UMR-botAnique Et Modélisation de L'Architecture Des Plantes Et Des Végétations (AMAP), CIRAD, Université Montpellier, CNRS, INRAE, 34398 Montpellier, IRD, France