



Ozone pollution threatens bird populations to collapse: an imminent ecological threat?

Evgenios Agathokleous¹ · Pierre Sicard² ·
Zhaozhong Feng¹ · Elena Paoletti³

Received: 31 May 2023 / Accepted: 20 June 2023 / Published online: 5 August 2023
© The Author(s) 2023

Abstract While bird populations are declining, the factors associated with this decline are unclear. Based on laboratory experiments, air pollution has long been recognized as a factor causing oxidative stress and adversely affecting bird health. Recently, studies employing an epidemiological approach have reported significant declines in avian populations in Central Europe and the United States due to air pollution, and ozone in particular. We advocate that urgent actions are needed to mitigate these effects, which threaten biodiversity and environmental health, and propose a series of measures which can enlighten the path toward mitigating air pollution effects on avian populations.

Keywords Air pollution · Biosecurity · Bird population · Biodiversity · Ornithology · Ozone risk

Project funding: This study was supported by the National Natural Science Foundation of China (No. 4210070867 and 42130714).

The online version is available at <http://www.springerlink.com>.

Corresponding editor: Yu Lei.

✉ Evgenios Agathokleous
evgenios@nuist.edu.cn

¹ School of Applied Meteorology, Nanjing University of Information Science & Technology (NUIST), Nanjing 210044, People's Republic of China

² ARGANS, Sophia Antipolis, France

³ IRET-CNR, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy


Introduction

Worldwide ground-level ozone (O₃) exposures display a multi-fold increase, e.g. about 35–50 nmol mol⁻¹ average O₃ concentration in the northern hemisphere, compared to the pre-industrial levels (Agathokleous et al. 2020). This increase is attributed to anthropogenic inputs that change the balance between volatile organic compounds (VOCs) and nitrogen oxides (NO_x) (O₃ precursors) under ideal weather conditions (e.g., temperature, sunlight). Due to its complex chemistry and the influence of natural emitters of VOCs (e.g. vegetation) even when NO_x are decreased, mediation of O₃ pollution is challenging and expected to persist for several decades to come (Agathokleous et al. 2020).

Bird populations are declining but the driving factors are unclear (Lehikoinen et al. 2019), and air pollution has long been considered as a potential factor (Sanderfoot and Holloway 2017). Specifically, an abundance of controlled and in situ studies show that birds experience adverse effects upon a variety of air pollutants including O₃, carbon monoxide (CO), heavy metals, smoke, sulfur dioxide (SO₂), and mixtures of emissions (Sanderfoot and Holloway 2017), of which O₃ exposures are predominant at high elevations (Sicard et al. 2016). When exposed to toxic levels of such air pollutants, birds undergo oxidative stress and allocate resources/energy to scavenge harmful reactive chemical species and detoxify (Sanderfoot and Holloway 2017) (Fig. 1). As the stress elevates, birds experience distress of the respiratory system, immunosuppression, illness, behavioral alterations, and potentially impaired reproductive success (Sanderfoot and Holloway 2017).

While ongoing O₃ pollution is well-known to adversely affect human health and vegetation (Otu-Larbi et al. 2020; Sicard et al. 2020; Holm and Balmes 2022), new studies employing a thorough, epidemiological-like approach

Fig. 1 Tropospheric ozone (O_3) pollution threatens bird populations. VOCs: volatile organic compounds; NO_x : nitrogen oxides



have reported significant related declines in birds populations in the United States and Central Europe (Liang et al. 2020; Reif et al. 2023), now demonstrating the real-world impacts of air pollution on avian populations and further supporting the expectations based on stress biology and the toxicological literature (Sanderfoot and Holloway 2017). For example, an analysis of 25-year data from a Central European mountain range (Giant Mountains, Czech), which also considered the impact of weather conditions, revealed that O_3 exposure had a significant negative effect on the population growth rates of upland bird species in the alpine zone above treeline (Reif et al. 2023). The effect was non-significant across all species, and the O_3 risks become higher along an altitudinal gradient (Reif et al. 2023). A further analysis of data from the United States revealed an estimated averted loss of approximately 20% (1.5 billion birds) of current total bird populations due to improvements of air quality in the last four decades (Liang et al. 2020). Massive deaths of birds, facilitated by emissions of primary pollutants boosting O_3 formation, threaten diminishing alpha diversity, and undermine biodiversity and conservation targets (Agathokleous et al. 2020) and, thus, must be averted.

In accordance with the Convention on Biological Diversity (<https://www.un.org/en/observances/biological-diversity-day/conventio>), more actions are needed to aid a sustainable future. Immediate actions by individual countries and unions are needed to comply with the effective laws. Existing laws include the European Council Directive 2009/147/EC on the conservation of wild birds (aka Birds Directive) and the Migratory Bird Treaty Act of 1918, a US's federal

law and convention with countries like Canada, Mexico, Japan, and Soviet Union.

A solution to mediate the ongoing decline in bird populations is to limit O_3 exposure, especially at high elevation areas where O_3 levels are higher, creating ideal conditions for chronic high exposures of birds (Lelieveld and Dentener 2000; Agathokleous et al. 2023; Reif et al. 2023). Limiting O_3 precursor emissions in urban areas could lead to lower transport of O_3 and its precursors to high-elevation areas, and weakened O_3 exposures (Cristofanelli et al. 2021; Rizos et al. 2022; Wang et al. 2022). However, this countermeasure alone may not be expected to suffice, since local O_3 formation is largely regulated by the VOCs-to- NO_x ratio (Calatayud et al. 2023; Guo et al. 2023; Latha et al. 2023; Lee et al. 2023; Lu et al. 2023). Hence, while NO_x can be decreased by regulating primary air pollutant emissions, VOCs are mostly biogenic and increase with climate warming (Bourtsoukidis et al. 2012), and, under ideal conditions, reduction of anthropogenic precursor emissions could increase O_3 exposures (Querol et al. 2021; Calatayud et al. 2023). Importantly, at high-elevation areas, there is a considerable input from stratospheric O_3 , further increasing O_3 exposures (Li et al. 2018; Hu et al. 2022). Hence, limiting O_3 exposures is challenging, policies to reduce air pollution are often unsuccessful to decrease O_3 exposures, and decreasing O_3 precursor emissions overall without accounting for atmospheric chemistry reactions may fail to decrease O_3 exposures. More measures are needed besides aiming at merely decreasing pollutant emissions.

There are many technologies allowing for monitoring of O_3 and other air pollutant concentrations, some of which are

low-cost (Idrees and Zheng 2020; Saitanis et al. 2020). Due to lack of electricity and difficulty to access, there is lack of monitoring stations on mountainous forests and especially at high-elevation areas. This is a fundamental issue because O₃ risk assessments and exposure standards are mainly based on urban and suburban stations, while high heterogeneity and insufficiency of monitoring stations to represent air pollutant exposures still exists in large areas of the world (De Marco et al. 2022). However, technological advances, such as the use of solar panels, provide an opportunity to develop and expand monitoring networks to record exposures (Paoletti et al. 2019) and evaluate risks to wildlife. Hence, to limit O₃ exposures on mountainous forests, it is essential to greatly expand the coverage of air pollution monitoring stations across such areas.

Another solution is a holistic approach to increase bird health. High chronic O₃ exposures cause oxidative stress in birds (Sanderfoot and Holloway 2017) while decreasing yield and alter the nutritional quality of plant edible products (Hoshika et al. 2022; Sahoo et al. 2023). The nutraceutical quality of fruits may increase in some tolerant plants but it may decrease in susceptible plants, suggesting bird species-specific risks depending on their feeding behavior. Local programs are needed to assist birds such as by providing antioxidant supplements during feedings and watering, especially in urban areas where supervised citizens can support such actions. Moreover, further developments are needed to increase the tolerance of plants to O₃ such as via bioengineering and by applying approved plant protectants that alleviate O₃ stress; however, the applications of such approaches in the real-world require extensive assessments to determine whether they may pose a threat to biosecurity, including ecological health as well as the health of animals that may consume such bioengineered products or products treated with synthetic chemicals. If such holistic measures are not taken, the global populations of birds are at risk of collapse and biodiversity threatened.

Acknowledgements This paper was prepared within the International Union of Forest Research Organizations (IUFRO) Research Group 8.04.00 “Air Pollution and Climate Change” and under the Working Party 8.04.05 “Ground-level Ozone.”

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Agathokleous E, Feng Z, Oksanen E, Sicard P, Wang Q, Saitanis CJ, Araminiene V, Blande JD, Hayes F, Calatayud V, Domingos M, Veresoglou SD, Peñuelas J, Wardle DA, De Marco A, Li Z, Harmens H, Yuan X, Vitale M, Paoletti E (2020) Ozone affects plant, insect, and soil microbial communities: a threat to terrestrial ecosystems and biodiversity. *Sci Adv* 6:eabc1176. <https://doi.org/10.1126/sciadv.abc1176>
- Agathokleous S, Saitanis CJ, Savvides C, Sicard P, Agathokleous E, De Marco A (2023) Spatiotemporal variations of ozone exposure and its risks to vegetation and human health in Cyprus: an analysis across a gradient of altitudes. *J for Res* 34:579–594. <https://doi.org/10.1007/s11676-022-01520-2>
- Bourtsoukidis E, Bonn B, Dittmann A, Hakola H, Hellén H, Jacobi S (2012) Ozone stress as a driving force of sesquiterpene emissions: a suggested parameterisation. *Biogeosciences* 9:4337–4352. <https://doi.org/10.5194/bg-9-4337-2012>
- Calatayud V, Diéguez JJ, Agathokleous E, Sicard P (2023) Machine learning model to predict vehicle electrification impacts on urban air quality and related human health effects. *Envir Res* 228:115835. <https://doi.org/10.1016/j.envres.2023.115835>
- Cristofanelli P, Arduni J, Serva F, Calzolari F, Bonasoni P, Busetto M, Maione M, Sprenger M, Trisolino P, Putero D (2021) Negative ozone anomalies at a high mountain site in northern Italy during 2020: a possible role of COVID-19 lockdowns? *Environ Res Lett* 16:074029. <https://doi.org/10.1088/1748-9326/ac0b6a>
- De Marco A, Sicard P, Feng Z, Agathokleous E, Alonso R, Araminiene V, Augustatis A, Badea O, Beasley JC, Branquinho C, Bruckman VJ, Collalti A, David-Schwartz R, Domingos M, Du E, Garcia Gomez H, Hashimoto S, Hoshika Y, Jakovljevic T, McNulty S, Oksanen E, Omid Khaniabadi Y, Prescher AK, Saitanis CJ, Sase H, Schmitz A, Voigt G, Watanabe M, Wood MD, Kozlov MV, Paoletti E (2022) Strategic roadmap to assess forest vulnerability under air pollution and climate change. *Glob Chang Biol* 28:5062–5085. <https://doi.org/10.1088/1748-9326/ac0b6a>
- Guo J, Zhang X, Gao Y, Wang Z, Zhang M, Xue W, Herrmann H, Brasseur GP, Wang T, Zhe W (2023) Evolution of ozone Pollution in China: What track will it follow? *Environ Sci Technol* 57:109–117. <https://doi.org/10.1021/acs.est.2c08205>
- Holm SM, Balmes JR (2022) Systematic review of ozone effects on human lung function, 2013 through 2020. *Chest* 161:190–201. <https://doi.org/10.1016/j.chest.2021.07.2170>
- Hoshika Y, Cotrozzi L, Marchica A, Carrari E, Lorenzini G, Nali C, Paoletti E, Pellegrini E (2022) Season-long exposure of bilberry plants to realistic and future ozone pollution improves the nutraceutical quality of fruits. *Sci Total Environ* 822:153577. <https://doi.org/10.1016/j.scitotenv.2022.153577>
- Hu J, Zhao T, Liu J, Cao L, Wang C, Li Y, Shi C, Tan C, Sun X, Shu Z, Li J (2022) Exploring the ozone pollution over the western Sichuan Basin, southwest China: the impact of diurnal change in mountain-plains solenoid. *Sci Total Environ* 839:156264. <https://doi.org/10.1016/j.scitotenv.2022.156264>
- Idrees Z, Zheng L (2020) Low cost air pollution monitoring systems: a review of protocols and enabling technologies. *J Ind Inf Integr* 17:100123. <https://doi.org/10.1016/j.jii.2019.100123>
- Latha R, Bano S, More D, Ambulkar R, Mondal T, Maurya P, Murthy BS (2023a) On the transition of major pollutant and O₃ production regime during Covid-19 lockdowns. *J Environ Manage* 328:116907. <https://doi.org/10.1016/j.jenvman.2022.116907>
- Lee HJ, Kuwayama T, FitzGibbon M (2023) Trends of ambient O₃ levels associated with O₃ precursor gases and meteorology in California: synergies from ground and satellite observations. *Remote Sens Environ* 284:113358. <https://doi.org/10.1016/j.rse.2022.113358>

- Lehikoinen A, Brotons L, Calladine J, Campedelli T, Escandell V, Flousek J, Grueneberg C, Haas F, Harris S, Herrando S, Husby M, Jiguet F, Kälås JA, Lindström Å, Lorrillière R, Molina B, Pladevall C, Calvi G, Sattler T, Schmid H, Sirkkiä PM, Teufelbauer N, Trautmann S (2019) Declining population trends of European mountain birds. *Glob Chang Biol* 25:577–588. <https://doi.org/10.1111/gcb.14522>
- Lelieveld J, Dentener FJ (2000) What controls tropospheric ozone? *J Geophys Res* 105(D3):3531–3551. <https://doi.org/10.1029/1999JD901011>
- Li S, Wang T, Huang X, Pu X, Li M, Chen P, Yang XQ, Wang M (2018) Impact of east Asian summer monsoon on surface ozone pattern in China. *J Geophys Res Atmos* 123:1401–1411. <https://doi.org/10.1002/2017JD027190>
- Liang Y, Rudik I, Zou EY, Johnston A, Rodewald AD, Kling CL (2020) Conservation cobenefits from air pollution regulation: evidence from birds. *Proc Natl Acad Sci* 117(49):30900–30906. <https://doi.org/10.1073/pnas.2013568117>
- Lu B, Zhang Z, Jiang J, Meng X, Liu C, Herrmann H, Chen J, Xue L, Li X (2023) Unraveling the O₃-NO_x-VOCs relationships induced by anomalous ozone in industrial regions during COVID-19 in Shanghai. *Atmos Environ* 308:119864. <https://doi.org/10.1016/j.atmosenv.2023.119864>
- Otu-Larbi F, Conte A, Fares S, Wild O, Ashworth K (2020) Current and future impacts of drought and ozone stress on northern hemisphere forests. *Glob Chang Biol* 26:6218–6234. <https://doi.org/10.1111/gcb.15339>
- Paoletti E, Alivernini A, Anav A, Badea O, Carrari E, Chivulescu S, Conte A, Ciriani ML, Dalstein-Richier L, De Marco A, Fares S, Fasano G, Giovannelli A, Lazzara M, Leca S, Materassi A, Moretti V, Pitar D, Popa I, Sabatini F, Salvati L, Sicard P, Sorgi T, Hoshika Y (2019) Toward stomatal-flux based forest protection against ozone: the MOTTLES approach. *Sci Total Environ* 691:516–527. <https://doi.org/10.1016/j.scitotenv.2019.06.525>
- Querol X, Massagué J, Alastuey A, Moreno T, Gangoiti G, Mantilla E, Duéñez JJ, Escudero M, Monfort E, Pérez García-Pando C, Petetin H, Jorba O, Vázquez V, de la Rosa J, Campos A, Muñoz M, Monge S, Hervás M, Javato R, Cornide MJ (2021) Lessons from the COVID-19 air pollution decrease in Spain: now what? *Sci Total Environ* 779:146380–146380. <https://doi.org/10.1016/j.scitotenv.2021.146380>
- Reif J, Gamero A, Flousek J, Hůnová I (2023b) Ambient ozone—new threat to birds in mountain ecosystems? *Sci Total Environ* 876:162711. <https://doi.org/10.1016/j.scitotenv.2023.162711>
- Rizos K, Logothetis I, Koukouli ME, Meleti C, Melas D (2022) The influence of the summer tropospheric circulation on the observed ozone mixing ratios at a coastal site in the eastern Mediterranean. *Atmos Pollut Res* 13:101381. <https://doi.org/10.1016/j.apr.2022.101381>
- Sahoo JP, Mishra P, Mishra AP, Panda KK, Samal KC (2023) Physiological, biochemical, and molecular responses of rice (*Oryza sativa* L.) towards elevated ozone tolerance. *Cereal Res Commun* 51:315–324. <https://doi.org/10.1007/s42976-022-00316-8>
- Saitanis CJ, Sicard P, De Marco A, Feng ZZ, Paoletti E, Agathokleous E (2020) On the atmospheric ozone monitoring methodologies. *Curr Opin Environ Sci Heal* 18:40–46. <https://doi.org/10.1016/j.coesh.2020.07.004>
- Sanderfoot OV, Holloway T (2017) Air pollution impacts on avian species via inhalation exposure and associated outcomes. *Environ Res Lett* 12:083002. <https://doi.org/10.1088/1748-9326/aa8051>
- Sicard P, De Marco A, Carrari E, Dalstein-Richier L, Hoshika Y, Badea O, Pitar D, Fares S, Conte A, Popa I, Paoletti E (2020) Epidemiological derivation of flux-based critical levels for visible ozone injury in European forests. *J for Res* 31:1509–1519. <https://doi.org/10.1007/s11676-020-01191-x>
- Sicard P, Serra R, Rossello P (2016) Spatiotemporal trends in ground-level ozone concentrations and metrics in France over the time period 1999–2012. *Envir Res* 149:22–144. <https://doi.org/10.1016/j.envres.2016.05.014>
- Wang L, Liu D, Yan W, Kang Z, Liu R, Zhang J, Li Z (2022) Spatiotemporal distribution, transport characteristics and synoptic patterns of ozone pollution near surface in Jiangsu province, China. *Atmos Pollut Res* 13:101616. <https://doi.org/10.1016/j.apr.2022.101616>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.