Acupuncture-related techniques for the treatment of opiate addiction: a case of translational medicine

Jisheng Han (✉)^{1,2,3,4}, Cailian Cui^{1,2,3,4}, Liuzhen Wu^{1,2,3,4}

¹ Neuroscience Research Institute, Peking University, Beijing 100191, China; ² Department of Neurobiology, School of Basic Medical Sciences, Peking University, Beijing 100191, China; ³ Key Laboratory for Neuroscience, Ministry of Education, Beijing 100191, China; ⁴ Key Laboratory for Neuroscience, Ministry of Health, Beijing 100191, China

© Higher Education Press and Springer-Verlag Berlin Heidelberg 2011

Frontiers of Medicine 2011, 5(2): 141–150

A mistake appeared in the sentence on lines 4–6 on the right column of page 148.

The sentence “Low frequency (represented by 2 Hz) EA is most effective for suppressing the withdrawal syndromes and high frequency (100 Hz) EA is best for alleviating craving.” should be changed to “Low frequency (represented by 2 Hz) EA is most effective for alleviating craving and high frequency (100 Hz) EA is best for suppressing the withdrawal syndromes.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11684-011-0136-8>

Received March 11, 2011; accepted April 11, 2011
Correspondence: hanjisheng@bjmu.edu.cn